



## Biofilm formation of *Flavobacterium psychrophilum* on various substrates

Journal:	<i>Aquaculture Research</i>
Manuscript ID	ARE-OA-18-Mar-251.R1
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Ríos Castillo, Abel G.; Universidad Autònoma de Barcelona, Department of Animal and Food Science Thompson, Kim; Moredun Research Institute, Aquaculture Research Group Adams, Sandra; University of Stirling, Institute of Aquaculture Marín de Mateo, Mercedes; Universitat Autònoma de Barcelona, Department of Animal and Food Science Rodríguez Jerez, José J.; Universitat Autònoma de Barcelona, Department of Animal and Food Science
Keywords:	Flavobacterium psychrophilum, biofilm formation, fluorescent microscopy, substrates, fish farming systems, rainbow trout farms

SCHOLARONE™  
Manuscripts

This is the peer reviewed version of the following article: Ríos-Castillo AG, Thompson KD, Adams A, Marín de Mateo M, Rodríguez-Jerez JJ. Biofilm formation of *Flavobacterium psychrophilum* on various substrates. *Aquac Res.* 2018;49:3830–3837, which has been published in final form at <https://doi.org/10.1111/are.13849>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

1     **Biofilm formation of *Flavobacterium psychrophilum* on various substrates**

2

3     Abel **Guillermo** Ríos-Castillo<sup>a</sup>, Kim D. Thompson<sup>b</sup>, Alexandra Adams<sup>c</sup>, Mercedes Marín de

4     Mateo<sup>a</sup>, José **Juan** Rodríguez-Jerez<sup>a</sup>

5

6     <sup>a</sup>Departament de Ciència Animal i dels Aliments, Facultat de Veterinària, Universitat

7     Autònoma de Barcelona, CP 08193, Barcelona, Spain.

8     <sup>b</sup>Moredun Research Institute, Pentlands Science Park, Bush Loan, Penicuik EH26 0PZ, UK.

9     <sup>c</sup>Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling FK9

10    4LA, UK.

11

12    Corresponding author: Abel **Guillermo** Ríos-Castillo. Facultat de Veterinària, Universitat

13    Autònoma de Barcelona, Travessera dels Turons s/n. Bellaterra, 08193, Barcelona, Spain.

14    Phone: +34 935811448. E-mail: [abelguillermo.rios@uab.cat](mailto:abelguillermo.rios@uab.cat)

15

16

## Abstract

The ability of *Flavobacterium psychrophilum* to adhere to and form biofilms on different types of materials used on rainbow trout (*Oncorhynchus mykiss*) farms was evaluated in this study. *F. psychrophilum* NCIMB 1947<sup>T</sup>, was inoculated onto a variety of different surfaces, including stainless steel, plastic, glass, wood and zinc pyrithione encapsulated antibacterial plastic. The samples were then cultured in a humidified chamber or transferred into fish tanks containing either (1) freshwater or (2) filtered lake water. The formation of biofilms was quantified by fluorescent microscopy. *F. psychrophilum* formed biofilms on all of the surfaces tested, however, the adherence of the bacterium to the antibacterial plastic was much lower than the attachment observed on the other surfaces, illustrating the bacteriostatic properties of this material for *F. psychrophilum*. Moreover, bacterial numbers were greater on the surfaces maintained in lake water compared to those maintained in freshwater. The mineral composition of the lake water may have been responsible for the increased bacterial adherence observed between the two types of water. Treatment of the water, regular cleaning of equipment and the use of antimicrobial material to house the fish may help reduce biofilm formation by *F. psychrophilum* in fish farming systems.

## Key words

*Flavobacterium psychrophilum*, biofilm formation, fluorescent microscopy, substrates, fish farming systems, rainbow trout farms

## 1. Introduction

The continuous use of water within a fish farm can act as a reservoir for pathogenic bacteria, and this can be a factor in the spread of disease within the farm (King, 2001; Cai & Arias, 2017). In the aquatic environment, bacteria rarely occur in a planktonic form, and their presence is more likely to be associated with surface-associated microbial communities known as biofilms (Huq, Whitehouse, Grim, Alam & Colwell, 2008; Nocker, Burr & Camper, 2014; Satpathy, Sen, Pattanaik & Raut, 2016). Biofilms are structured communities of bacterial cells adhered to inert or living surfaces and enclosed in a polymeric matrix produced by the bacteria, referred to as an extracellular polymer substance (EPS). Biofilm formation is beneficial to the bacteria, providing them with protection against desiccation, increased nutrient availability and is more resistant to antimicrobial agents than planktonic bacteria (Costerton, Stewart & Greenberg, 1999; Srey, Jahid & Ha, 2013; Satpathy et al., 2016). Biofilms are composed of a variety of microflora present in the water, capable of colonizing surfaces, which can then act as a reservoir for pathogenic bacteria. Pathogenic microorganisms within the biofilm can be shed from the biofilm and are able to cause a reoccurrence of disease in fish (King, 2001; Branda, Vik, Friedman & Kolter, 2005; Nocker et al., 2014). Biofilm formation is important to many pathogenic bacterial species, especially those living in water, giving them a selective advantage by increasing their ability to persist under adverse environmental conditions (Duchaud et al., 2007). They can form on many of the materials found within aquaculture systems, appearing on the surfaces of water pipes and fish tanks, suspended matter, incubators, bio-filtration systems and even on the internal and external surfaces of the fish (King et al., 2004).

Bacteria belonging to genus *Flavobacterium* have been identified as a group of bacteria able to persist in a latent form in the aquatic environment (Waśkiewicz & Irzykowska, 2014). *F. psychrophilum* is a Gram-negative, yellow-pigmented bacterium,

67 responsible for causing cold water disease (CWD) (Borg, 1948; Holt, Rohovec & Fryer,  
68 1993), or rainbow trout fry syndrome (RTFS) (Lorenzen, 1994; Rangdale, 1995) in salmonids  
69 and other freshwater fish. The bacterium not only infects farmed fish, but can also affect wild  
70 fish, although disease outbreaks are apparently less severe in non-salmonids (Nematollahi,  
71 Decostere, Pasmans & Haesebrouck, 2003). It is, currently, one of the main bacterial  
72 pathogens in reared and wild salmonids, causing substantial economic losses in salmonid fish  
73 farms worldwide, and hindering expansion of the salmonid aquaculture industry (Nematollahi  
74 et al., 2003; Bernardet & Bowman, 2006). The bacterium grows in the aquatic environment in  
75 temperatures ranging between 4°C to 23°C (Holt, 1988). It has the ability to adhere to the  
76 skin, gut and eggs of fish and disease transmission studies suggest that reservoirs of the  
77 bacterium can be found within the water system of the fish farm (Madetoja, Dalsgaard &  
78 Wiklund, 2002). It also has the ability to adhere to surfaces forming biofilms, and has been  
79 detected in sediment, river water, especially near outlet water from infected fish farms  
80 (Amita, Hoshino, Honma & Wakabayashi, 2000; Álvarez, Secades, Prieto, McBride &  
81 Guijarro, 2006; Sundell & Wiklund, 2011).

82 As aquaculture facilities are particularly prone to the development of biofilms by *F.*  
83 *psychrophilum*, understanding the factors that influence biofilm formation could reduce the  
84 presence of this pathogenic bacterium within the fish farming system (Huq et al., 2008;  
85 Wietz, Hall & Høj, 2009; Srey et al., 2013). Thus, the purpose of this study was to gain a  
86 better understanding of the survival of this bacterium in the aquatic environment and to  
87 examine the ability of *F. psychrophilum* to adhere to and form biofilms on different types of  
88 materials used by the salmonid aquaculture industry. Biofilm formation by *F. psychrophilum*  
89 was examined in the presence of tryptone yeast extract salts (TYES) broth, freshwater taken  
90 from the aquarium or water from a freshwater lake.

91

## 2. Materials and methods

### 2.1. Bacterial culture

*F. psychrophilum*, strain NCIMB 1947<sup>T</sup>, was obtained from a stock of cryopreservation beads (Cryoprotect; Technical Service Consultants Service Ltd. Lancashire, UK) stored at -70°C. The bacterium was grown in TYES broth (tryptone, 4.0 g; yeast extract, 0.4 g; MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.5 g; CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.2 g; distilled water, 1000 mL; pH 7.2; autoclaved for 20 min at 121°C) under constant agitation at 140 rpm (Kühner Shaker LT-W, Adolf Kühner AG, Switzerland) for 72-96 h at 15°C. This culture was subsequently cultured on TYES agar plates (TYES broth with 15.0 g/L bacteriological agar) and incubated for 96 h at 15°C, from which bacterial colonies were taken and cultivated in TYES broth under agitation for 72-96 h at 15°C to obtain the bacterial culture used in the analysis.

### 2.2. Test materials

Four different types of material, e.g. stainless steel (type 1.4301, also known as grade 304) used as positive control (Fuster-Valls, Hernández, Marín de Mateo & Rodríguez-Jerez, 2008), polyethylene (PE) plastic, silica glass, wood (*Pinus* sp.) and an antibacterial plastic [polypropylene (PP) containing micro-encapsulated zinc pyrithione] (Microlitix, Sant Cugat del Valles, Spain) were selected to assess their ability to support *F. psychrophilum* biofilm formation. The size of the material used was 4.0 x 4.0 cm<sup>2</sup>, while the stainless steel surfaces consisted of discs with a diameter of 2.0 cm and a thickness of 1.2 mm. Prior to performing the study, the surfaces were cleaned, disinfected and autoclaved at 121°C for 15 min (Ríos-Castillo, González-Rivas & Rodríguez-Jerez, 2017), except the antibacterial plastic, which was only cleaned with 70% isopropyl alcohol (2-propanol) before use so as not to interfere with its antibacterial properties.

### 2.3. Experimental design

The formation of biofilms on the different test materials was performed by incubating the supports in a *F. psychrophilum* suspension under two different sets of test conditions.

(a) In the first test, all surfaces were inoculated with  $16 \mu\text{L}/\text{cm}^2$ . The number of live and dead bacteria was determined using the LIVE/DEAD staining kit with a bacterial concentration on the surfaces of  $2.01 \times 10^6$  live cells/ $\text{cm}^2$  and  $4.79 \times 10^1$  of dead/damaged cells/ $\text{cm}^2$ . The surfaces inoculated were placed in Petri dishes, which were then placed into a humidified chamber (30 x 22 x 14 cm) maintained at a saturated relative humidity of  $\geq 90\%$  using pieces of paper towel moistened with sterile distilled water (Wiklund & Dalsgaard, 2003; Fuster-Valls et al., 2008; International Organization for Standardization, 2011). The bacteria were incubated on these surfaces for 48 h at  $15^\circ\text{C}$ . After this time, the excess liquid was removed from each surface and  $50 \mu\text{L}$  of TYES broth was added to the surface of the stainless steel disc and  $255 \mu\text{L}$  to the other surfaces with a sterile pipette. These were then incubated for a further 48 h at  $15^\circ\text{C}$  before the degree of biofilm formation on each support was assessed using fluorescence microscopy.

(b) The ability of *F. psychrophilum* to form biofilms on the different surfaces (stainless steel, plastic, glass, antibacterial plastic and wood) was also assessed using freshwater collected from either a freshwater aquarium [Aquaculture Research Facility (ARF), Institute of Aquaculture, University of Stirling], or from a freshwater lake (Airthrey Loch, University of Stirling). The water samples taken from the aquarium and the lake were filtered through a  $0.45 \mu\text{m}$  filter (Millipore Co., Billerica, Ma, U.S.A.) prior to use. The various supports were incubated with *F. psychrophilum* ( $2.14 \times 10^6$  of viable cells/ $\text{cm}^2$  and  $7.41 \times 10^1$  cells/ $\text{cm}^2$  of dead or injured cells) for 72 h. After this time, the excess liquid was removed from each surface and the surfaces were attached onto the side of plastic fish tanks (20 x 40 x 22 cm) with Blu-Tack™ malleable rubber adhesive (Bostik Ltd, Leicester, UK). The tanks contained:

(i) 100 mL of TYES broth with 10 litres of freshwater from the aquarium (dechlorinated, mains water), or (ii) 100 mL of TYES broth with 10 litres of freshlake water. The surfaces were maintained in the tanks for 96 h at 15-16°C. After this incubation period, the test materials were removed from the tanks, washed carefully with distilled water, taking care not to disturb the biofilm on the surfaces and these were then examined by fluorescence microscopy using the LIVE/DEAD staining kit.

#### 2.4. Assessing the number of live bacteria by fluorescence microscopy

**Live/Dead BacLight Bacteria Viability Kit** (Molecular Probes, Europe BV) was used to determine the initial number of live bacteria present in the bacterial cultures used for the studies and to assess the number of live bacteria present on surfaces after 96 h of incubation with *F. psychrophilum*. The LIVE/DEAD staining kit is composed of two nucleic acid-binding stains: SYTO<sup>®</sup> 9, penetrating all bacterial membranes and stains the cells green, and propidium iodide which only penetrates cells with damaged membranes, producing red fluorescing cells when the cells are damaged or dead. The kit was used according to manufacturer's instructions.

The stainless steel discs were stained with 20 µL and plastic, glass, wood and the antibacterial plastic with 100 µL. Stained surfaces were left in the dark for 15 min at 22°C to allow the stains to penetrate. Eight images were acquired from every surface evaluated using a fluorescence microscope IX70 (Olympus Optical, Tokyo, Japan) equipped with a mercury lamp, and two filters: (1) filter A (excitation 470–490 nm, emission 515–550 nm) and (2) filter B (excitation 510–550 nm, emission > 590 nm). The same **microscopic** parameters, input calibration and image acquisition were used throughout and images were analysed using Cytovision<sup>®</sup> software, version 2.51 (Applied Imaging, Sunderland, Tyne & Wear, UK). Cell counts and bacteria size (i.e. minimum and maximum diameter and area) were automatically



measured as a colour scale interpretation using Soft Imaging System™ program, AnalySIS® version 3.2 (GmbH, Munich, Germany).

## 2.5. Water analysis

Prior to performing the fresh water studies, the mineral composition of the water was analysed from both water sources. To do this, 10 litres of freshwater was obtained from the aquarium facility and directly from the **freshwater lake**, both of which were filtered through a 0.45 µm filter. The samples of water were then analysed by inductively coupled plasma mass spectrometry (ICP-MS) to determine the concentration of their mineral content. This analysis was performed by the Water Quality Laboratory, Institute of Aquaculture, University of Stirling.

## 2.6. Statistical analysis

Each analysis comparing biofilm formation on the various supports was repeated three times and each test material surface was analysed in triplicate (n = 9). The statistical software package SAS® v 9.1.3.4 (Institute Inc, North Carolina, USA) was used for the statistical analysis. The assumption of normality of the data was carried out using the Shapiro–Wilk test. Statistical analysis was performed on data between cells counted on the various surfaces under the different test condition using an analysis of variance (ANOVA). Student-Newman-Keuls *post hoc* test was used to test the significance of differences between live and dead or injured bacteria on the surfaces, where  $p \leq 0.05$  was considered significant.

## 3. Results

### 3.1. Biofilm formation by *F. psychrophilum*

In the first study, in which a suspension of *F. psychrophilum* in TYES was incubated onto the various supports in a humidified chamber, the live-cell counts obtained using the LIVE/DEAD kit showed a similar amount of live and dead bacteria attached to the various supports (Table 1). There was no statistical difference obtained in the level of adherence or biofilm formation on stainless steel ( $1.41 \times 10^6$  live cells/cm<sup>2</sup>), plastic ( $9.12 \times 10^5$  live cells/cm<sup>2</sup>) or glass ( $8.32 \times 10^5$  live cells/cm<sup>2</sup>) after 96 h of incubation. Whereas adherence of the bacterium to the antibacterial plastic surface was significantly lower ( $p < 0.05$ ) than observed on the other materials with respect to both live ( $6.46 \times 10^3$  cells/cm<sup>2</sup>) and dead/injured cells ( $1.10 \times 10^4$  cells/cm<sup>2</sup>). Also, an increase in the number of cells/cm<sup>2</sup> of dead or injured cells was observed on all test materials, increasing from an initial concentration of  $4.79 \times 10^1$  cells/cm<sup>2</sup> to  $10^4$  or  $10^5$  cells/cm<sup>2</sup> after 96 h of being introduced on to the support.

### 3.2. Biofilm formation of *F. psychrophilum* in aquarium or lake water

The degree of bacterial attachment to the various supports (stainless steel, plastic, glass, wood and antibacterial plastic) after 96 h at 15 - 16°C, in either the aquarium water or the lake water, is presented in Table 2. Higher levels of live bacteria ( $p < 0.05$ ) were present in the freshwater from the aquarium adhering to stainless steel ( $8.68 \times 10^4$  cells/cm<sup>2</sup>), plastic ( $1.09 \times 10^5$  cells/cm<sup>2</sup>) (Figure 1c), glass surfaces ( $8.52 \times 10^4$  cells/cm<sup>2</sup>) (Figure 1d) and wood ( $1.11 \times 10^5$  cells/cm<sup>2</sup>) compared to the antibacterial plastic ( $2.88 \times 10^2$  cells/cm<sup>2</sup>). The results with the water obtained from freshwater lake showed significantly higher levels of live-cells ( $p < 0.05$ ) attached to stainless steel ( $2.30 \times 10^5$  cells/cm<sup>2</sup>), plastic ( $2.98 \times 10^5$  cells/cm<sup>2</sup>) and glass ( $2.41 \times 10^5$  cells/cm<sup>2</sup>) compared with the wood ( $1.38 \times 10^5$  cells/cm<sup>2</sup>) (Figure 1e) or the antibacterial plastic surface ( $6.03 \times 10^3$  cells/cm<sup>2</sup>) (Figure 1f). Figure 2 shows the values in parts per billion (ppb) of sodium, magnesium, potassium and calcium of freshwater aquarium and in lake water used to evaluate the biofilm formation of *P. psychrophilum*. According to

these results, the concentrations of all minerals analysed in the lake water were higher than those obtained from the freshwater aquarium. The highest mineral element concentration for the lake water was calcium (13760.0 ppb), and the lowest magnesium (1022.0 ppb). In the case of aquarium water, the highest value, although lower than that found in the lake water was also calcium (7118.5 ppb), while the potassium concentration was the lowest (211.0 ppb).

### 3.3. Antimicrobial properties of antibacterial plastic [poly-propylene (PP) containing micro-encapsulated zinc pyrithione]

The results of this study indicate that the antibacterial properties of zinc pyrithione under saturated relative humidity of  $\geq 90\%$  had  $2.34 \log_{10}$  cells/cm<sup>2</sup> fewer bacteria attached relative to the positive control,  $2.48 \log_{10}$  cells/cm<sup>2</sup> in freshwater and  $1.58 \log_{10}$  cells/cm<sup>2</sup> in lake water, while only a minimal reduction was observed for other surfaces under the various conditions.

## 4. Discussion

When environmental conditions are unfavourable, aquatic bacteria are subjected to a rapid change in nutrient availability and must therefore adapt accordingly in order to be able to survive under these adverse conditions. For example, cells undergo reduced cell division, with the resulting cells having an overall reduction in size and typically become rounder and coccus in morphology, in what is known as a 'rounding up' strategy (Arias, LaFrentz, Cai & Olivares-Fuster, 2012).

In our study, where *F. psychrophilum* cells were incubated on the various supports in the humidity chamber, some of the dead/injured bacteria became rounded in appearance. The morphological changes in *F. psychrophilum* cells observed here have also been reported by Vatsos, Thompson & Adams (2003), for bacteria maintained in a broth culture for four weeks.

In this study, these changes were observed after only 96 h incubation suggesting that this adaptation may be accelerated during the growth of the bacteria on the surfaces compared to growth in TYES broth, reflecting the environment stress experience by the bacteria during biofilm formation. This reduction in bacterial size during biofilm formation, which in mature stages of biofilm contained more damaged cells (dead or non-viable) than live cells, has also been reported by Roszak & Colwell (1987); Boulos, Prevost, Barbeau, Coallier & Desjardins, (1999); Chmielewski & Frank (2003); and Fuster-Valls et al. (2008). The results also suggest that high levels of humidity and the use of TYES broth could favour the adhesion and biofilm formation of *F. psychrophilum*. According to Ehrlich, Miller & Walker (1970), the survival of *Flavobacterium* sp. is not affected by high conditions of humidity (up to 99%), but can be affected by a lack of nutrients. Under dry conditions, Fuster-Valls et al. (2008) observed a considerable reduction in the level of bacterial attachment by the cells, with some cells appearing injured, and non-culturable in culture medium. They were still considered to have the potential to cause disease outbreaks, however. Humid areas within the fish farming system, ideal for bacterial growth, can favour the adhesion and biofilm formation by *F. psychrophilum*, and microorganisms present on equipment and surfaces within the fish farm, may survive there for prolonged periods of time (Lee Wong, 2004).

The results of the mineral analysis may explain the high levels of bacteria seen adhering to the surfaces in the presence of the lake water compared to the aquarium water (Figure 2). These values were statistically different ( $p < 0.05$ ) for the live cell counts attached to the stainless steel, glass and antibacterial zinc pyrithione surfaces. These results are in accord with Fletcher (1988), who observed that cationic metal concentrations of sodium, calcium, magnesium minimize the repulsive forces between the bacterial cell and surfaces, having an influence on the ability of bacteria to adhere to surfaces and form biofilms. The lower concentration of minerals in the aquarium freshwater may also influence bacterial

attachment; in fact, a deficiency of certain nutrients may increase the ability of bacteria to form biofilms, though the concentration of nutrients necessary for bacterial development is low (Mattila-Sandholm & Wirtanen, 1992; Percival & Walker, 1999). The presence of organic and inorganic material can also influence biofilm formation by bacteria within the *Flavobacterium* genus. Staroscik, Hunnicutt & Nelson (2007) observed that the addition of  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  or glucose to the culture medium, or the presence of mucus from salmon skin induced the formation of biofilms by *Flavobacterium columnare*. Likewise, the environment can represent a reservoir of *F. psychrophilum*, since the ability of this microorganism to adhere to surfaces could explain the bacterium's survival under adverse conditions. The fact that water can act as a source of infection implies that *F. psychrophilum* is able to survive outside its host under conditions of starvation (Vatsos, Thompson & Adams, 2001). Madetoja, Nystedt & Wiklund (2003) found that the virulence of *F. psychrophilum* was maintained for at least seven days after transferring the bacteria to freshwater, and the bacterium's survival increased with the addition of nutrient-containing sediments; thereby *F. psychrophilum* can readily spread from infected fish to uninfected ones in recirculating aquaculture systems. The differences in bacterial counts (expressed in decimal logarithms,  $\log_{10}$ ) of live-cells adhered to plastic, glass, wood or antibacterial plastic were compared with the number of live-cells attached to the stainless steel surfaces, used as a control (Table 3). According to Japanese Standard JIS Z 2801 (Japanese Standards Association, 2010) and ISO 22196 (International Organization for Standardization, 2011) surfaces with antibacterial properties must demonstrate a reduction in bacterial attachment equal to or higher than 2  $\log_{10}$  of that determined for the control surface. The zinc pyrithione antibacterial plastic showed a high efficiency in preventing bacterial adherence when it was tested under the humidity conditions (2.34 log) or under the aquarium water condition (2.48 log). On the other hand, when it was tested under the lake water condition, the efficiency was lower (1.58 log) (Table 3). This

could be explained because the high mineral concentration of sodium, magnesium, potassium, and calcium in the lake water may prevent the adequate action of zinc-pyrithione. It has been earlier reported that higher level of minerals favour the adherence of *Flavobacterium* and biofilm formation (Madetoja et al., 2003; Staroscik et al., 2007). The antibacterial action of zinc pyrithione in preventing the adherence of cells is favoured by the use freshwater used in fish farms and is partly inhibited by the presence of water with a high mineral content. Zinc pyrithione interacts with the membrane phospholipids in bacteria, inhibiting membrane transport of substrates and decreasing intracellular ATP levels by inhibiting ATP synthesis causing a lethal toxicity of bacterial cells (Qian, Chen & Xu, 2013).

As established from the genome analysis of *F. psychrophilum*, the bacterium has the ability to form biofilms and store cyanophycin, which could explain the bacterium's prolonged survival outside its host (Duchaud et al., 2007) and the spread of disease by this bacterium through the aquatic environment (Madetoja et al., 2002; Nematollahi et al., 2003). The ability of this bacterium to adhere to surfaces and form biofilms may explain why it is less susceptible to antimicrobial treatment. Sundell & Wiklund (2011) observed an increased antimicrobial resistance in *F. psychrophilum* biofilms containing high bacterial cell densities ( $> 10^7$  CFU/mL). These characteristics, together with adherent properties of *F. psychrophilum* may explain the subsequent transmission of this bacterium to fish, and probably contribute to its dissemination in salmonid fish farms, representing a significant risk in the development of the salmonid aquaculture (Nematollahi et al., 2003; Barnes & Brown, 2011). Zinc pyrithione, is widely used as an antifouling agent in paints and exhibit a high antimicrobial effects against biofilm bacteria (Konstantinou & Albanis, 2004; Ciriminna, Bright & Pagliaro, 2015). The commercial cost of zinc pyrithione, used in a concentration of 2.0% as biocide and antifouling is approximately US\$ 2.50 - US\$ 3.50 to cover each 100 m<sup>2</sup> of fish-farm environments. Although this is an added expense to the fish farms, the reduced mortality caused by disease

from this bacterium justifies the investment. Thus, the use of materials that inhibit bacterial growth such as zinc pyrithione may offer alternative ways to reduce the spread of *F. psychrophilum* within the fish farming system as well as other bacterial species involved in disease outbreaks.

## 5. Conclusions

This study suggests that *F. psychrophilum* has the ability to adhere to and form biofilms on materials used within aquaculture systems such as stainless steel, plastic, glass and wood at saturated relative humidity levels of  $\geq 90\%$  and in freshwater aquarium or lake water.

Procedures such as water treatment, regular sanitation of equipment, and the use of antimicrobial surfaces may be useful in preventing biofilm formation in fish farming systems, and in turn preventing disease outbreaks caused by this bacterium.

## Acknowledgments

The authors wish to acknowledge technical support in the laboratory and in the Aquarium Research Facilities at the Institute of Aquaculture, University of Stirling, where this work was carried out.

## Conflict of interest

The authors declare no conflicts of interest.

## References

- 340 Álvarez, B., Secades, P., Prieto, M., McBride, M.J., & Guijarro, J.A. (2006). A mutation in  
341 *Flavobacterium psychrophilum tlpB* inhibits gliding motility and induces biofilm  
342 formation. *Applied and Environmental Microbiology*, 72, 4044–4053.  
343 doi:10.1128/AEM.00128-06
- 344 Amita, K., Hoshino, M., Honma, T., & Wakabayashi, H. (2000). An investigation on the  
345 distribution of *Flavobacterium psychrophilum* in the Umikawa River. *Fish Pathology*, 35,  
346 193–197. doi:10.3147/jsfp.35.193
- 347 Arias, C.R., LaFrentz, S., Cai, W., & Olivares-Fuster, O. (2012). Adaptive response to  
348 starvation in the fish pathogen *Flavobacterium columnare*: cell viability and  
349 ultrastructural changes. *BMC Microbiology*, 12, 266. doi:10.1186/1471-2180-12-266
- 350 Barnes, M.E., & Brown, M.L. (2011). A review of *Flavobacterium psychrophilum* biology,  
351 clinical signs, and bacterial cold water disease prevention and treatment. *The Open Fish*  
352 *Science Journal*, 4, 40–48. doi:10.2174/1874401X01104010040
- 353 Bernardet, J.-F., & Bowman, J. P. (2006). The genus *Flavobacterium*. In M. Dworkin, S.  
354 Falkow, E. Rosenberg, K.-H. Schleifer & E. Stackebrandt (Eds.), *The Prokaryotes: a*  
355 *Handbook on the Biology of Bacteria* (pp. 481–531). New York: Springer.
- 356 Borg, A.F. (1948). Studies on myxobacteria associated with diseases in salmonid fishes.  
357 Ph.D. Thesis. University of Washington. Seattle, WA, USA.
- 358 Boulos, L., Prévost, M., Barbeau, B., Coallier, J., & Desjardins, R. (1999). LIVE/DEAD®  
359 *BacLight*™: application of a new rapid staining method for direct enumeration of viable  
360 and total bacteria in drinking water. *Journal of Microbiological Methods*, 37, 77–86.  
361 doi:10.1016/S0167-7012(99)00048-2
- 362 Branda, S.S., Vik, A., Friedman, L., & Kolter, R. (2005). Biofilms: the matrix revisited.  
363 *Trends in Microbiology*, 13, 20–26. doi:10.1016/j.tim.2004.11.006



- 364 Cai, W., & Arias, C.R. (2017). Biofilm Formation on Aquaculture Substrates by Selected  
365 Bacterial Fish Pathogens. *Journal of Aquatic Animal Health*, 29, 95–104.  
366 doi:10.1080/08997659.2017.1290711
- 367 Chmielewski, R., & Frank, J. (2003). *Comprehensive Reviews in Food Science and Food*  
368 *Safety*, 2, 22–32. doi:10.1111/j.1541-4337.2003.tb00012.x
- 369 Ciriminna, R., Bright, F.V., & Pagliaro, M. (2015). Ecofriendly Antifouling Marine Coatings.  
370 *ACS Sustainable Chemistry & Engineering*, 3, 559–565. doi:10.1021/sc500845n
- 371 Costerton, J.W., Stewart, P.S., & Greenberg, E.P. (1999). Bacterial biofilms: A common  
372 cause of persistent infections. *Science*, 284, 1318–1322.  
373 doi:10.1126/science.284.5418.1318
- 374 Duchaud, E., Boussaha, M., Loux, V., Bernardet, J.F., Michel, C., Kerouault, B., Mondot, S.,  
375 Nicolas, P., Bossy, R., Caron, C., Bessi eres, P., Gibrat, J., Claverol, S., Dumetz, F.,  
376 Henaff, M., & Benmansour, A. (2007). Complete genome sequence of the fish pathogen  
377 *Flavobacterium psychrophilum*. *Nature Biotechnology*, 25, 763–769.  
378 doi:10.1038/nbt1313
- 379 Ehrlich, R., Miller, S., & Walker, R.L. (1970). Effects of Atmospheric Humidity and  
380 Temperature on the Survival of Airborne *Flavobacterium*. *Applied Microbiology*, 20,  
381 884–887.
- 382 Fletcher, M. (1988). Attachment of *Pseudomonas fluorescens* to glass and influence of  
383 electrolytes on bacterium-substratum separation distance. *Journal of Bacteriology*, 170,  
384 2027–2030. doi:10.1128/jb.170.5.2027-2030.1988
- 385 Fuster-Valls, N., Hern andez-Herrero, M., Mar n de Mateo, M., & Rodr guez-Jerez, J.J.  
386 (2008). Effect of different environmental conditions on the bacteria survival on stainless  
387 steel surfaces. *Food Control*, 9, 308–314. doi:10.1016/j.foodcont.2007.04.013

- 388 Holt, R.A. (1988). *Cytophaga psychrophila*, the causative agent of bacterial cold-water  
389 disease in salmonid fish. Ph.D. Thesis. Oregon State University. Corvallis, OR, USA.
- 390 Holt, R.A., Rohovec, J.S., & Fryer, J.L. (1993). Bacterial cold-water disease. In V. Inglis, R.J.  
391 Roberts, N.R. Bromage (Eds.), *Bacterial Diseases of Fish* (pp. 3–23). Oxford: Blackwell  
392 Scientific Publications.
- 393 Huq, A., Whitehouse, C.A., Grim, C.J, Alam, M., & Colwell, R.R. (2008). Biofilms in water,  
394 its role and impact in human disease transmission. *Current Opinion in Biotechnology*, 19,  
395 244–247. doi:10.1016/j.copbio.2008.04.005
- 396 International Organization for Standardization (2011). Measurement of antibacterial activity  
397 on plastics and other non-porous surfaces, ISO 22196:2011.
- 398 Japanese Standards Association (2010). Antimicrobial products–Test for antimicrobial  
399 activity and efficacy. Japanese Industrial Standard, JIS Z 28010.
- 400 King, R.K. (2001). The Presence of Bacterial Pathogens in Recirculating Aquaculture System  
401 Biofilms and their Response to Various Sanitizers. Ph.D. Thesis. Virginia Polytechnic  
402 Institute and State University. Blacksburg, VA, USA.
- 403 King, R.K., Flick Jr., G.J., Pierson, M.D., Smith, S.A., Boardman, G.D., & Coale, C.W.J.  
404 (2004). Identification of bacterial pathogens in biofilms of recirculating aquaculture  
405 systems. *Journal of Aquatic Food Product Technology*, 13, 125–133.  
406 doi:10.1300/J030v13n01\_11
- 407 Konstantinou, I.K., & Albanis, T.A. (2004). Worldwide occurrence and effects of antifouling  
408 paint booster biocides in the aquatic environment: a review. *Environment International*,  
409 30, 235–248. doi:10.1016/S0160-4120(03)00176-4
- 410 Lee Wong, A.C. (2004). Biofilm formation and control in food processing environments. In  
411 R. Jordan, D. Williams, U. Charaf (Eds.), *Advances in Biofilm Science and Engineering*  
412 (pp. 245–281). Bozeman: Cytergy Publishing.

- 413 Lorenzen, E. (1994). Studies on *Flexibacter psychrophilus* in relation to rainbow trout fry  
414 syndrome (RTFS). Ph.D. Thesis. Royal Veterinary and Agricultural University.  
415 Copenhagen, Denmark.
- 416 Madetoja, J., Dalsgaard, I., & Wiklund, T. (2002). Occurrence of *Flavobacterium*  
417 *psychrophilum* in fish-farming environment. Diseases of Aquatic Organisms, 52, 109–  
418 118. doi:10.3354/dao052109
- 419 Madetoja, J., Nystedt, S., & Wiklund, T. (2003). Survival and virulence of *Flavobacterium*  
420 *psychrophilum* in water microcosms. FEMS Microbiology Ecology, 43, 217–223.  
421 doi:10.1111/j.1574-6941.2003.tb01061.x
- 422 Mattila-Sandholm, T., & Wirtanen, G. (1992). Biofilm formation in the industry: a review.  
423 Food Reviews International, 8, 573–603. doi:10.1080/87559129209540953
- 424 Nematollahi, A., Decostere, A., Pasmans, F., & Haesebrouck, F. (2003). *Flavobacterium*  
425 *psychrophilum* infections in salmonid fish. Journal of Fish Diseases, 26, 563–574.  
426 doi:10.1046/j.1365-2761.2003.00488.x
- 427 Nocker, A., Burr, M., & Camper, A. (2014). Pathogens in Water and Biofilms. In S.L.  
428 Percival, M.V. Yates, D.W. Williams, R.M. Chalmers, N.C. Gray (Eds.), Microbiology  
429 of Waterborne Diseases (pp. 3–32). London: Academic Press.
- 430 Percival, S.L., & Walker, J.T. (1999). Potable Water and Biofilms: A Review of the Public  
431 Health Implications. Biofouling, 14, 99–115. doi:10.1080/08927019909378402
- 432 Qian, P.Y., Chen, L., & Xu, Y. (2013). Mini-review: Molecular mechanisms of antifouling  
433 compounds. Biofouling, 29, 381–400. doi:10.1080/08927014.2013.776546
- 434 Rangdale, R.E. (1995). Studies on Rainbow Trout Fry Syndrome (RTFS). Ph.D. Thesis.  
435 University of Stirling. Scotland, UK.
- 436 Ríos-Castillo, A.G., González-Rivas, F., & Rodríguez-Jerez, J.J. (2017). Bactericidal Efficacy  
437 of Hydrogen Peroxide-Based Disinfectants against Gram-Positive and Gram-Negative

- 438 Bacteria on Stainless Steel Surfaces. *Journal of Food Science*, 82, 2351–2356.  
439 doi:10.1111/1750-3841.13790
- 440 Roszak, D.B., & Colwell, R.R. (1987). Survival strategies of bacteria in the natural  
441 environment. *Microbiological Reviews*, 51, 365–379.
- 442 Satpathy, S., Sen, S.K., Pattanaik, S., & Raut, S. (2016). Review on bacterial biofilm: An  
443 universal cause of contamination. *Biocatalysis and Agricultural Biotechnology*, 7, 56–66.  
444 doi:10.1016/j.bcab.2016.05.002
- 445 Srey, S., Jahid, I.K., & Ha, S.D. (2013). Biofilm formation in food industries: A food safety  
446 concern. *Food Control*, 31, 572–585. doi:10.1016/j.foodcont.2012.12.001
- 447 Staroscik, A.M., Hunnicutt, D.W., & Nelson, D.R. (2007). The influence of culture conditions  
448 on biofilm formation in *Flavobacterium columnare*. *Flavobacterium 2007 Workshop* (pp.  
449 21). National Conservation Training Center. Shepherdstown, WV, USA.
- 450 Sundell, K., & Wiklund, T. (2011). Effect of biofilm formation on antimicrobial tolerance of  
451 *Flavobacterium psychrophilum*. *Journal of Fish Diseases*, 34, 373–383.  
452 doi:10.1111/j.1365-2761.2011.01250.x
- 453 Vatsos, I.N., Thompson, K.D., & Adams, A. (2001). Adhesion of the fish pathogen  
454 *Flavobacterium psychrophilum* to unfertilized eggs of rainbow trout (*Oncorhynchus*  
455 *mykiss*) and n-hexadecane. *Letters in Applied Microbiology*, 33, 178–182.  
456 doi:10.1046/j.1472-765x.2001.00980.x
- 457 Vatsos, I.N., Thompson, K.D., & Adams, A. (2003). Starvation of *Flavobacterium*  
458 *psychrophilum* in broth, stream water and distilled water. *Diseases of Aquatic Organisms*,  
459 56, 115–126. doi:10.3354/dao056115
- 460 Waśkiewicz, A., & Irzykowska, L. (2014). *Flavobacterium* spp. – Characteristics,  
461 Occurrence, and Toxicity. In C.A. Batt, M.L. Tortorello (Eds.), *Encyclopedia of Food*  
462 *Microbiology* (pp. 938–942). Oxford: Academic Press.

- 463 Wietz, M., Hall, M.R., & Høj, L. (2009). Effects of seawater ozonation on biofilm  
464 development in aquaculture tanks. *Systematic and Applied Microbiology*, 32, 266–277.  
465 doi:10.1016/j.syapm.2009.04.001
- 466 Wiklund, T., & Dalsgaard, I. (2003). Association of *Flavobacterium psychrophilum* with  
467 rainbow trout (*Oncorhynchus mykiss*) kidney phagocytes *in vitro*. *Fish & Shellfish*  
468 *Immunology*, 15, 387–395. doi:10.1016/S1050-4648(03)00009-3
- 469

For Review Only

**Tables**

**Table 1.** Adherence and biofilm formation of *Flavobacterium psychrophilum* (cells/cm<sup>2</sup>) on stainless steel, plastic, glass, and antibacterial plastic surfaces after 96 h of incubation in humidity test condition.

Surfaces	Cells <sup>†</sup>	
	Live	Injured or dead
Stainless steel	1.41 x 10 <sup>6</sup> ± 0.37 <sup>a</sup>	3.47 x 10 <sup>5</sup> ± 0.45 <sup>a</sup>
Plastic	9.12 x 10 <sup>5</sup> ± 0.06 <sup>a</sup>	9.33 x 10 <sup>5</sup> ± 0.48 <sup>a</sup>
Glass	8.32 x 10 <sup>5</sup> ± 0.02 <sup>a</sup>	5.62 x 10 <sup>5</sup> ± 0.05 <sup>a</sup>
Antibacterial plastic	6.46 x 10 <sup>3</sup> ± 0.17 <sup>b</sup>	1.10 x 10 <sup>4</sup> ± 0.28 <sup>b</sup>

<sup>†</sup>Initial cells count/cm<sup>2</sup>: 2.01 x 10<sup>6</sup> live cells/cm<sup>2</sup> and 4.79 x 10<sup>1</sup> dead/injured cells/cm<sup>2</sup>.

<sup>a,b</sup> Values in columns for each surface are significantly different if the letters are different ( $p \leq 0.05$ ).

**Table 2.** Adherence and biofilm formation of *Flavobacterium psychrophilum* (cells/cm<sup>2</sup>) to stainless steel, plastic, glass, wood, and antibacterial plastic surfaces after 96 hours in aquarium freshwater and lake water conditions.

Surfaces	Aquarium water		Lake water	
	Cells <sup>†</sup>			
	Live	Injured or dead	Live	Injured or dead
Stainless steel	8.68 x 10 <sup>4</sup> ± 0.12 <sup>a A</sup>	1.46 x 10 <sup>5</sup> ± 0.12 <sup>ab</sup>	2.30 x 10 <sup>5</sup> ± 0.13 <sup>a A</sup>	1.54 x 10 <sup>5</sup> ± 0.39 <sup>a</sup>
Plastic	1.09 x 10 <sup>5</sup> ± 0.08 <sup>a B</sup>	5.87 x 10 <sup>4</sup> ± 0.11 <sup>ab</sup>	2.98 x 10 <sup>5</sup> ± 0.08 <sup>a A</sup>	3.09 x 10 <sup>5</sup> ± 0.50 <sup>a</sup>
Glass	8.52 x 10 <sup>4</sup> ± 0.11 <sup>a A</sup>	1.25 x 10 <sup>5</sup> ± 0.12 <sup>ab</sup>	2.41 x 10 <sup>5</sup> ± 0.09 <sup>a B</sup>	2.32 x 10 <sup>5</sup> ± 0.28 <sup>a</sup>
Wood	1.11 x 10 <sup>5</sup> ± 0.10 <sup>a A</sup>	2.80 x 10 <sup>5</sup> ± 0.39 <sup>a</sup>	1.38 x 10 <sup>5</sup> ± 0.07 <sup>b A</sup>	1.56 x 10 <sup>5</sup> ± 0.40 <sup>a</sup>
Antibacterial plastic	2.88 x 10 <sup>2</sup> ± 0.13 <sup>b B</sup>	1.39 x 10 <sup>4</sup> ± 0.34 <sup>c</sup>	6.03 x 10 <sup>3</sup> ± 0.17 <sup>c A</sup>	2.25 x 10 <sup>4</sup> ± 0.19 <sup>b</sup>

<sup>†</sup> Initial cells count/cm<sup>2</sup>:  $2.14 \times 10^6$  live cells and  $7.41 \times 10^1$  dead or injured cells. <sup>a-c</sup> Values in columns for each surface are significantly different if the letters are different ( $p \leq 0.05$ ). <sup>A-B</sup> Values in rows for each surface for live cells results are significantly different if the letters are different ( $p \leq 0.05$ ).

**Table 3.** The differences (represented in  $\log_{10}$  cells/cm<sup>2</sup>) in the live-cell counts of *F. psychrophilum* attached to plastic, glass, wood and antibacterial plastic compared with stainless steel used as a positive control. Values higher than 2  $\log_{10}$  represent surfaces with bacteriostatic properties according to the conditions evaluated.

Surfaces	Humidity condition	Freshwater or lake water conditions	
		Freshwater aquarium	Lake water
Stainless steel	6.15	4.94	5.36
Plastic	0.19	+ 0.1	+ 0.11
Glass	0.23	0.01	+ 0.02
Wood	-	+ 0.11	0.22
Antibacterial plastic	<u>2.34</u>	<u>2.48</u>	1.58

Positive signs (+) in  $\log_{10}$  values at plastic, glass or wood surfaces represent an increase of cells count respect to the stainless steel surface. No signs before the values represent a reduction respect the stainless steel as a control. Reductions with more than 2  $\log_{10}$  are underlined.



## Figure legends

**Figure 1.** Examples of the fluorescence microscopy images of *Flavobacterium psychrophilum* cells stained with the LIVE/DEAD<sup>®</sup> kit after 96 h of incubation. Live cells appeared green in colour and dead or injured cell appeared red. In humidity condition: (a) live cells forming biofilm, (b) round-shaped appearance of a dead cell indicated by an arrow on stainless steel surfaces. In freshwater condition: (c) high density of dead or injured cells on plastic surface, (d) the presence of live and dead or injured cells on glass surface. In lake water condition: (e) wood, (f) zinc pyrithione plastic surface. All scale bars: 10 µm.

**Figure 2.** Mineral concentration in parts per billion (ppb) of freshwater aquarium and lake water used to examine biofilm formation by *F. psychrophilum*. Abbreviations: Na-sodium, Mg-magnesium, K-potassium and Ca-calcium.




Figure 1. Examples of the fluorescence microscopy images of *Flavobacterium psychrophilum* cells stained with the LIVE/DEAD® kit after 96 h of incubation. Live cells appeared green in colour and dead or injured cell appeared red. In humidity condition: (a) live cells forming biofilm, (b) round-shaped appearance of a dead cell indicated by an arrow on stainless steel surfaces. In freshwater condition: (c) high density of dead or injured cells on plastic surface, (d) the presence of live and dead or injured cells on glass surface. In lake water condition: (e) wood, (f) zinc pyrithione plastic surface. All scale bars: 10 µm.

162x183mm (300 x 300 DPI)




Figure 2. Mineral concentration in parts per billion (ppb) of freshwater aquarium and lake water used to examine biofilm formation by *F. psychrophilum*. Abbreviations: Na-sodium, Mg-magnesium, K-potassium and Ca-calcium.

209x96mm (300 x 300 DPI)