

The effect of resource quality on the growth of *Holothuria scabra* during aquaculture waste bioremediation

Georgina Robinson^{1,2,*}, Gary S. Caldwell¹, Clifford L.W. Jones², Selina M. Stead¹

¹School of Natural and Environmental Sciences, Newcastle University, Newcastle, NE1 7RU, UK.

²Department of Ichthyology and Fisheries Science, Rhodes University, Grahamstown 6140, South Africa.

#Current address: Scottish Association for Marine Science, Scottish Marine Institute, PA37 1QA, Oban, UK

*Corresponding author. Tel +230 5982 4971; Email address Georgina.Robinson@sams.ac.uk (G. Robinson)

Accepted refereed manuscript of:

Robinson G, Caldwell GS, Jones CLW & Stead SM (2019) The effect of resource quality on the growth of *Holothuria scabra* during aquaculture waste bioremediation. *Aquaculture*, 499, pp. 101-108.

DOI: <https://doi.org/10.1016/j.aquaculture.2018.09.024>

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

Abstract

Reducing dependency on environmentally unsustainable formulated feeds, most of which include limited reserves of fishmeal as a protein source, is a priority for the aquaculture industry, particularly for intensive culture systems. One approach is to increase nitrogen reuse within the system by feeding nitrogen-rich aquaculture effluent to deposit feeders, thereby closing the aquaculture nitrogen-loop. This study, for the first time and on a laboratory-scale, has reared juveniles of the sea cucumber *Holothuria scabra* at high densities solely on particulate organic waste from a commercial-scale land-based abalone recirculating aquaculture system. Furthermore, growth rates and biomass yields were increased significantly by adjusting the effluent C:N from 5:1 to 20:1 by adding exogenous organic carbon sources (glucose, starch and cellulose), so fuelling accelerated heterotrophic bacterial production within the redox-stratified tank sediment. Sea cucumber juveniles reared solely on effluent had a biomass density of 711 g m⁻² after four months whereas animals reared on starch-amended effluent (the best performing treatment) had a final density of 1,011 g m⁻². Further optimisation of this approach could increase biomass yields and pave the way for the commercial cultivation of deposit feeding animals on waste streams, thus contributing to more environmentally sustainable aquaculture. Here, the nitrogen that originated from fishmeal is not lost through the discharge of aquaculture effluent; rather, it is immobilised into single cell biomass that is up-cycled into high-value secondary biomass. We demonstrate that sea cucumbers can be produced at high density through the manipulation of the C:N ratio of aquaculture effluent.

Keywords: C/N ratio; deposit feeder; stoichiometry; sustainable aquaculture; recirculating aquaculture system; sediment; sandfish

1. Introduction

Intensive aquaculture is generally characterised by the addition of nutritionally enriched diets that aim to satisfy the requirements of the culture species and completely replace natural food sources, a process not without its sustainability challenges (Naylor et al., 2009).

Intensive aquaculture is associated with an enrichment of toxic wastes such as ammonia and other nitrogenous species, which are treated using biological filtration. High-protein feed inputs thus lead to an inefficient use of nitrogen, and the waste of the natural resources, mostly fishmeal, from which this nitrogen originated.

As aquaculture has intensified, there has been a shift towards recirculating aquaculture systems (RAS) (Badiola et al., 2012), from which the effluent streams are typically separated into high volume flows of dissolved inorganic effluents and low volume flows of suspended solids that accumulate as sludge. To advance the sustainability agenda and strengthen the economics of intensive aquaculture, there is a clear case for the industry developing culture strategies based on nitrogen reuse rather than removal.

Detritivores such as sea cucumbers and polychaete worms are ideal candidates for nitrogen reuse, yielding an additional commercial crop whilst bioremediating nitrogenous effluent (Cubillo et al., 2016; Zamora et al., 2016). Sea cucumbers are highly prized in Far Eastern markets and aquaculture is considered the only means of meeting demand, with production growing to ~130,000 tonnes per annum (Han et al., 2016). Culture technologies include sea ranching, sea pen farming, pond farming, production in co-culture and integrated multi-trophic aquaculture systems, and in intensive RAS (Purcell, 2010; Robinson, 2013).

Microorganisms play pivotal roles in aquaculture bioremediation technologies, which have evolved from exploiting autotrophic microbes (photoautotrophs and chemolithoautotrophs) to fully heterotrophic systems (Ebeling et al., 2006). This transition emphasises the re-use and recycling of feed residues within the culture system thereby

reducing feed, space, and energy requirements (Chávez-Crooker and Obreque-Contreras, 2010). Sediment microbial communities are primarily net heterotrophic systems that link energy transfer to higher trophic levels; therefore recycling nutrients *in situ* may provide a viable means to intensively culture deposit feeders with a higher overall efficiency (Schroeder, 1987). Furthermore, prior research (Robinson et al., 2015; Robinson et al., 2016) demonstrated that redox-stratified sediment supported faster growth rates and a higher biomass yield of *Holothuria scabra* relative to fully oxic sediment, indicating that heterotrophic systems are more favourable for deposit feeder growth.

Heterotrophs fundamentally differ from autotrophs due to their metabolic requirement for an organic source of carbon. Heterotrophic bacteria assimilate organic carbon and nitrogen in a stoichiometric balance based on the carbon to nitrogen ratio (C:N) of the bacterial cytoplasm (Goldman et al., 1987; Herbert, 1967). The C:N of organic substrates is an important parameter determining the degree of nitrogen regeneration as carbon and nitrogen are incorporated into bacteria at a fixed rate (Tezuka, 1990). For bacteria grown in an environment with a C:N of 5:1 and an average growth efficiency (the quantity of biomass produced per unit of assimilated organic carbon) of 50% under aerobic conditions, the threshold between net release and net immobilisation of nitrogen is 10:1 (Azim et al., 2008; Rittmann and McCarty, 2001). Increasing C:N beyond 10:1 provides sufficient carbon for heterotrophic bacteria to assimilate ammonium (NH_4^+) into biomass, thus mediating a shift from net NH_4^+ release (ammonification) to net immobilisation (assimilation) (Avnimelech, 1999; Avnimelech, 2014; Azim et al., 2008; Ebeling et al., 2006).

From a thermodynamic perspective, heterotrophic bacteria preferentially utilise reduced inorganic forms of nitrogen such as NH_4^+ ; however, NH_4^+ assimilation is dependent on the availability of carbon substrates (Church, 2008; Fenchel and Blackburn, 1979). Particulate organic wastes from aquaculture primarily comprise waste food and faeces, and are generally

deficient in organic carbon, with an average C:N of 7:1; thus, there is a net release of NH_4^+ during decomposition, often measured alongside ammonia as total ammonia nitrogen (TAN) (Avnimelech, 1999; Mirzoyan et al., 2012; Schneider et al., 2006). Particulate organic waste recovered from mechanical filtration has been used as substrate to produce heterotrophic bacteria and deposit feeding macrofauna, including polychaetes and sea cucumbers (Brown et al., 2011; MacDonald et al., 2013; Palmer, 2010; Schneider et al., 2007a; Schneider et al., 2007b; Schneider et al., 2006). Raising secondary livestock on aquaculture waste can provide a direct means of assimilating a proportion of the effluent nitrogen (Erler et al., 2004); however, detritivores are predicted to have poor nitrogen retention compared to other trophic groups (Schneider et al., 2005). Erler et al. (Erler et al., 2004) hypothesized that stimulated bacterial nitrogen processing during the production of secondary livestock on RAS effluents may be more important than direct assimilation. Schneider et al. (Schneider et al., 2007b) demonstrated that adding a source of labile organic carbon, e.g. molasses, could increase the conversion of inorganic nitrogenous wastes to heterotrophic bacterial biomass. Stimulating heterotrophic bacteria by manipulating the C:N may therefore offer an indirect means of increasing nitrogen retention in macrofauna reared on aquaculture effluents.

In sediment-based sea cucumber culture, controlling inorganic nitrogen cycling by adding carbon may be particularly relevant due to the need to counteract additional sources of NH_4^+ within the system, including: i) net efflux from the sediment (Hargreaves, 1998); ii) excretion from the sea cucumbers (Uthicke and Klumpp, 1997); and, iii) decomposition of aquaculture waste and feeds (Avnimelech, 1999). Carbon supplementation may offer an indirect means to retain nitrogen safely within the system by immobilising NH_4^+ into microbial biomass that can be upcycled into high value secondary biomass. We therefore investigated whether, by carbon supplementation, the uptake of nitrogen from waste feed and faeces by sea cucumbers can be improved.

Applying C:N manipulation to sediment-based systems, where bacterial growth efficiencies are generally lower due to anoxia (Fenchel et al., 2012), is completely novel. As such, there is a need to test different carbon sources to determine their efficacy. This study compared a range of carbon sources of differing biochemical composition and degradation rates on the growth of *H. scabra* reared on particulate organic waste from an intensive abalone RAS.

2. Material and methods

2.1 Study site and experimental animals

The Ethics Panels of both Newcastle and Rhodes Universities approved the study, and no collections were made from wild populations to support it. The research was conducted in a purpose built, bio-secure, heated RAS between October 8th 2013 and January 28th 2014 at HIK Abalone Farm Pty (Ltd) in Hermanus, South Africa. The detailed system specifications are found in Robinson et al. (2015). Juvenile *Holothuria scabra* were imported from a commercial hatchery (Research Institute for Aquaculture III, Vietnam) on September 5th 2013, and quarantined in a bio-secure facility for six weeks in accordance with South African importation and scientific investigations licenses. Following the quarantine period and prior to experimentation, the sea cucumber juveniles were held in the RAS in tanks filled with 10 cm of calcium carbonate sand and were fed a 34% protein commercial abalone weaning diet (Abfeed®-S34, 1.0 mm sugar grain pellet; Marifeed Pty Ltd, South Africa). The proximate nutritional composition of the feed was; crude protein 33%, crude lipid 3.0%, vitamins (0.1%), gross energy 15.6 kJ g⁻¹.

2.2 Experimental design

The experiment was designed to test a range of carbon sources of differing biochemical composition on *H. scabra* growth in a sediment-based culture system. Two

soluble (D-glucose and starch) and one insoluble carbon sources (microcrystalline cellulose) representing a range of first-order decomposition rate constants and labilities were purchased from Merck Millipore, South Africa (Table 1). The carbon sources were tested in conjunction with aquaculture waste (comprising uneaten feed and faeces recovered by mechanical filtration from a land-based abalone RAS) at a C:N of 20:1; a fourth treatment receiving aquaculture waste only (C:N of 5:1) was included as a control (Table 1). The average C:N of the dried particulate aquaculture waste was 5.21 ± 0.55 . The quantity of carbon necessary to increase the ratio to 20:1 was calculated from the proportion of carbon in the molecular composition of each compound. Carbon additions were standardised between treatments based on a daily addition of $200 \text{ mmol C m}^{-2} \text{ d}^{-1}$, which is within the upper range tolerated by benthic animals under eutrophic conditions of $100\text{-}400 \text{ mmol C m}^{-2} \text{ d}^{-1}$ (Lehtoranta et al., 2009). At a C:N of 20:1, this equates to $2.4 \text{ g m}^{-2} \text{ d}^{-1}$ of carbon and $0.12 \text{ g m}^{-2} \text{ d}^{-1}$ of nitrogen, which is between a ‘mid’ and ‘high’ ration of 100 and $150 \text{ mg m}^{-2} \text{ d}^{-1}$ of nitrogen respectively for deposit feeders (Alongi and Hanson, 1985; Tenore and Chesney, 1985).

Abalone (*Haliotis midae*) waste comprising mainly uneaten feed and faeces was collected every morning at 09:00 from the backwash of a grow-out RAS sand filter over five days prior to the experiment. The total system volume was approximately 32,000 L, holding a maximum of 800 kg of adult abalone (Yearsley et al., 2009). The abalone were fed to satiation daily with a 34% protein commercial abalone weaning diet (S34 Abfeed®, $10 \times 10 \times 1.2 \text{ mm}$ pellet; Marifeed Pty Ltd, South Africa). The water leaving the production tanks drained through a 500 L sand filter filled with BS8:16 silica filter media for particulate organic waste removal (1-2 mm) prior to foam fractionation and biofiltration. The sand filter was fitted with a Jetco 5-port valve to change the water flow direction and permit backwashing. A 50 mm flexihose pipe was connected from the sand filter outflow to a 100 L conical fibreglass tank to collect the waste during the first 30 seconds of back washing. The

waste was settled for one hour and the supernatant discarded. The particulate waste was then concentrated by centrifuging in 50 mL at 10,000 g for 10 minutes (compact centrifuge Z 206 A, Hermle Labortechnik, Germany). The organic carbon and total nitrogen content of triplicate pre-weighed and dried (105 °C, 24 h) waste samples were analysed on a LECO TruSpec CHN elemental analyser.

The four experimental treatments were randomly allocated to one of sixteen tanks using a randomised block design of four blocks with one replicate from each treatment represented in each block. As such, there were four replicates per treatment and each replicate consisted of one laboratory-scale tank with four sea cucumbers per tank (i.e. sixteen cucumbers per treatment). These experimental tanks had an area of 0.125 m², which resulted in a sea cucumber stocking density equivalent to those used commercially. The tanks were filled with calcium carbonate sand sourced from a commercial dune quarry (SSB Mining, Macassar, South Africa) sieved to 125-250 µm. Tanks were supplied with heated (29.18 ± 0.26 °C) recirculating seawater and aeration as described in Robinson et al. (2015).

Feed additions (aquaculture waste with or without carbon sources) were made to all tanks at 16:00 hours daily. All tanks received 2.41 g of concentrated aquaculture waste on a wet weight basis per day. The experimental treatments received either 0.68 g of starch or cellulose or 0.75 g of glucose per day on a dry weight basis respectively (Table 1). Prior to feeding, the abalone waste was mixed with ambient seawater from the RAS into a wet slurry. Similarly, the carbon sources were prepared by dissolving the soluble carbon sources in beakers of ambient seawater while the insoluble cellulose was mixed in beakers to facilitate even dispersal. Aeration was supplied continuously, except during feeding when the air and water supplies were interrupted for 45 minutes. Polyvinylchloride end-caps (25 mm) were placed over the tank standpipes and maintained in position for one hour to prevent the waste and carbon sources from being washed out. The tank outflows were adapted to enable the tank

water to run to waste. This prevented the soluble carbon sources from entering the biological filter.

Tanks were cleaned every two weeks to remove floating cyanobacteria colonies (*Oscillatoria* sp.), epiphytic algae and biofilm. Tanks were subject to a natural photoperiod which increased from 12.41: 11.19 L: D (06:10 hours to 18:51 hours, sunrise to sunset) to 13.53: 10.07 L: D (05:59 hours to 19:52 hours, sunrise to sunset) as day length increased during the austral summer.

2.3 Water and sediment quality and environmental variables

Light readings (aerial) were taken using a portable light meter (LX-107, Lutron Electronic Enterprise Co. Ltd, Taipei, Taiwan) positioned 10 cm directly above the tank outflow. Water quality parameters including salinity, temperature, pH, dissolved oxygen and total ammonia nitrogen (NH₄-N; TAN) were recorded weekly as described in Robinson et al. (2015). Nitrate concentration ($\pm 0.01 \text{ mg L}^{-1}$) was measured weekly using a commercial test kit (Merck Nitrate Test Kit, 109713, Merck, South Africa) and a spectrophotometer (Prim Light, Secomam, 30319 Ales, France).

Sediment quality was monitored monthly to coincide with monthly sea cucumber growth assessments. Sediment reduction-oxidation (redox) potential was measured with a redox probe (Eutech Instruments pH 6+ portable meter) to the base of the sediment. Readings were taken following stabilization (typically five minutes). Composite samples of the sediment surface layers (upper 2-3 mm) were collected from all replicate tanks to determine sediment characteristics. Chlorophyll *a* and phaeopigment concentrations were measured using the spectrophotometric method described in Robinson et al. (2015). The organic content measured as particulate organic carbon and total nitrogen was determined on an elemental analyser after removal of carbonates by fuming with HCl. Total sediment carbohydrates ($\mu\text{g g}^{-1}$) were measured using the phenol-sulphuric acid method (Underwood et

al., 1995). The absorbance of the supernatant was measured at 485 nm, quantified against a glucose standard and converted into the concentration of total carbohydrates using coefficients derived from the standard curves.

2.4 Sea cucumber growth

At the start of the experiment juvenile sea cucumbers ($n = 64$; 4.08 ± 0.58 g; mean \pm standard deviation) were gut evacuated for 24 hours whilst suspended in mesh bags. They were drained on a damp cloth for one minute, weight and photographed, before being randomly allocated to tanks (four per tank), photo-identified and weighed. Each individual was re-weighed every four weeks (28 days) over the four-month experiment. Wet weight data were used to calculate biomass density (g m^{-2}) and growth rate (g d^{-1} ; Robinson et al., 2015).

2.5 Statistical analyses

For all measured parameters, the data recorded per replicate tank were averaged and the mean value per tank was used. The light and water quality data were averaged to provide a mean value per month to give five time intervals for repeated measures analysis of variance (repeated measures ANOVA) and ensure consistency with the sediment quality and sea cucumber weight data. Units of pH were transformed prior to averaging using the antilog function in Microsoft Excel. Results are expressed as mean \pm standard error unless otherwise stated.

Growth and environmental (light, water and sediment quality) data were tested for homogeneity of variance and for the normal distribution of the residuals using Levene's (Levene, 1960) and Shapiro Wilk's (Shapiro and Wilk, 1965) tests respectively. Initial weight data did not meet the assumptions of homogeneity of variance, therefore a non-parametric Kruskal-Wallis one-way ANOVA was used to test for significant differences between treatment medians. Data that met the test assumptions were analysed using repeated measures

ANOVA with treatment (carbon source) as the main factor and sampling time (month) as the repeated measure. Tukey's post-hoc honest significant difference (HSD) tests were used to compare differences among means. Multiple regression analysis was used to identify significant categorical predictors of sea cucumber biomass density. Differences were considered significant at $\alpha < 0.05$. All statistical analyses were performed using Statistica version 13.

3. Results

3.1 Water and sediment quality and environmental variables

There were no significant differences in light, salinity, temperature, dissolved oxygen concentration, total ammonia or nitrate among treatments (repeated measures ANOVA: $p > 0.05$; Table 2 and Table S1). Salinity maintained a constant 35 g L^{-1} and dissolved oxygen concentration varied between 6.05 and 8.95 mg L^{-1} ($7.26 \pm 0.07 \text{ mg L}^{-1}$). The mean water temperature increased over the course of the experiment with the onset of the austral summer from a mean temperature of $25.49 \pm 0.14 \text{ }^{\circ}\text{C}$ at the start, increasing to a mean of $32.30 \pm 0.34 \text{ }^{\circ}\text{C}$ during the final month. pH differed significantly between treatments (repeated measures ANOVA; $F_{(12, 48)} = 2.79$, $p = 0.006$). The starch-amended treatment had the highest seawater pH between months one and three, peaking at 9.12 ± 0.05 in month three. In all treatments, pH tended to increase with time until the final month when values returned to those measured at the start of the experiment (Fig. 1A).

Ammonia concentrations increased in all treatments from $0.026 \pm 0.001 \text{ mg L}^{-1}$ at the start, peaking at $0.84 \pm 0.15 \text{ mg L}^{-1}$ in month two, and decreasing during the remaining two months to a mean of $0.19 \pm 0.01 \text{ mg L}^{-1}$ (Fig. 1B). Nitrate levels increased significantly over the experiment from $1.05 \pm 0.08 \text{ mg L}^{-1}$ at the start to $4.90 \pm 0.23 \text{ mg L}^{-1}$ in the final month

(repeated measures ANOVA; $F_{(4, 48)} = 84.45$, $p < 0.001$), although there were no significant differences between treatments (combined mean: $3.92 \pm 0.20 \text{ mg L}^{-1}$; Table 2 and Table S1).

3.2 Sediment characteristics

The type of carbon source significantly affected sediment redox potential (repeated measures ANOVA, $F_{(12, 48)} = 4.76$, $p = <0.001$; Fig. 2A and Table S2). There were no significant differences between treatments at the start of the experiment with a positive reading of $173.06 \pm 5.99 \text{ mV}$ indicating fully oxic conditions. The redox potential decreased over time in all treatments; however, there was an apparent relatedness to the lability of the carbon source. The more refractory cellulose and starch had the greatest impact on redox potential, decreasing to $-179.75 \pm 18.93 \text{ mV}$ and $-109.50 \pm 6.14 \text{ mV}$ in the final month respectively. In the final month, the redox potential in tanks solely receiving aquaculture waste was marginally negative ($-22.00 \pm 12.19 \text{ mV}$) and not significantly different to glucose-amended tanks, the most labile carbon source ($-45.75 \pm 8.47 \text{ mV}$).

There were no significant differences in the levels of organic carbon, total nitrogen, C:N, total carbohydrate concentration, chlorophyll *a* or phaeopigment between treatments (repeated measures ANOVA: $p > 0.05$; Tables S2 and S3). Levels of organic carbon were initially low and increased in all treatments with time, with the highest levels of $0.20 \pm 0.05\%$ recorded in the cellulose-amended treatment (Fig. 2B and Table 2). Levels of total nitrogen were generally low ($0.05 \pm 0.00\%$) with minor fluctuations between treatments over time (Fig. 2C and Table 2). The surface sediment C:N of the starch-amended tanks was relatively constant over the experiment with a mean of $7.41 \pm 0.23\%$. The C:N in the other three treatments fluctuated over time and between treatments with no clear trend, although there was a C:N increase in these treatments in the final month (Fig. 2D).

3.3 Growth and survival

There were no significant differences in mean wet weight or biomass density between treatments at the start of the experiment ($130.59 \pm 1.92 \text{ g m}^{-2}$; Kruskal-Wallis, $H_{(3, 16)} = 1.53$, $p = 0.68$ (Fig 3A and 3C). Survival was 100% in all treatments. Over the course of the growth trial, the mean growth rate ($0.25 \pm 0.02 \text{ g d}^{-1}$) and biomass density ($1,011.46 \pm 75.58 \text{ g m}^{-2}$) of *H. scabra* reared in the starch treatment was significantly higher than those reared on the waste alone ($0.16 \pm 0.01 \text{ g d}^{-1}$ and $702.12 \pm 35.93 \text{ g m}^{-2}$) repeated measures ANOVA, $F_{(12, 48)} = 2.49$, $p = 0.013$; Fig 3). Sea cucumbers in the starch treatment reached a final biomass density of $1,011.46 \pm 75.58 \text{ g m}^{-2}$ by month-4 compared to $702.12 \pm 35.93 \text{ g m}^{-2}$ in the control tanks. There were no significant differences in growth rate or biomass densities between the three carbon sources.

3.4 Multiple regression analysis

Sediment redox potential, light intensity and nitrate concentration were significant predictors of *H. scabra* biomass density (multiple regression, $F_{(13, 60)} = 36.26$, $r^2 = 0.89$; $p < 0.001$). Light and nitrate showed a positive relationship while sediment redox potential had a negative relationship with sea cucumber density.

4. Discussion

The current study indicated that increasing the C:N from 5:1 to 20:1 through carbon supplementation successfully increased the growth rate and biomass density of *H. scabra* juveniles. It is possible that increasing the C:N improved the nutritional value of the waste by increasing the quantity of organic carbon. It is commonly thought that deposit feeders gain energy for maintenance from carbon while nitrogen is predominantly used for growth (Lopez and Levinton, 1987). Deposit feeders are assumed to metabolise organic carbon and nitrogen in a 17:1 molar ratio (Russell-Hunter, 1970); therefore, carbon addition may have improved

310 growth rates by balancing the elemental stoichiometry of the food (Rice and Rhoads, 1989).
 311 An alternate explanation is that *H. scabra* was able to assimilate the carbon sources or their
 312 degradation products directly or indirectly through collaboration with microbial communities.
 313 Mechanisms for the uptake of organic molecules such as amino acids and carbohydrates are
 314 thought to exist in tissues such as the respiratory trees, body wall or tentacles, thereby
 315 enabling holothurians to assimilate dissolved organic matter directly from the water column
 316 (Fontaine and Chia, 1968; Jaeckle and Strathmann, 2013; Jangoux and Lawrence, 1982;
 317 Krishnan and Krishnaswamy, 1970; Lawrence, 1982).
 318 Direct utilisation of starch for enhanced growth may not have been limited to the sea
 319 cucumbers. Although there was no significant difference in either chlorophyll *a* or
 320 phaeopigment concentrations, there were apparent differences (albeit qualitative) in the
 321 extent of phototroph biomass production between treatments. Carbohydrates such as glucose
 322 and polysaccharides can also be photo-oxidised with high quantum efficiencies in illuminated
 323 habitats leading to concomitant increases in microbial and algal productivity (Krishnan and
 324 Krishnaswamy, 1970; Stuart et al., 2016). The quality and quantity of organic matter supply
 325 to the sediment is modulated by sunlight, temperature and nutrient availability (Huettel et al.,
 326 2014). Light and nitrate were significant positive predictors of sea cucumber biomass density.
 327 Cyanobacteria may have supplied additional carbon to heterotrophic sediment bacteria and
 328 sea cucumbers as extracellular polymeric substances from photosynthesis (Stuart et al.,
 329 2016).
 330 Avnimelech (1999) hypothesised that manipulating C:N may change nitrogen cycling
 331 pathways in aquaculture systems by mediating a shift from ammonification (net release) to
 332 assimilation (net uptake) of NH_4^+ by heterotrophic bacteria. For aerobic heterotrophic
 333 bacteria, the removal of NH_4^+ by incorporation in cells is enhanced by carbohydrate addition
 334 (Ebeling et al., 2006). It was therefore expected in these experiments that carbon

335 supplementation would decrease total ammonia nitrogen (TAN) concentrations compared to
336 the control. However, this was not the case, although levels were generally low throughout
337 the study (averaging $0.35 \pm 0.04 \text{ mg L}^{-1}$). This may have been due to assimilation by
338 heterotrophic bacteria, which are responsible for a large fraction of NH_4^+ assimilation in
339 marine waters (Kirchman, 2012); however, the trends in the ammonia and nitrate data seem
340 to indicate that nitrification rather than assimilation was the dominant NH_4^+ conversion
341 pathway (Wu et al., 2013). The increasing nitrate concentration from the start of the
342 experiment to month two is consistent with an increasing nitrification capacity with the
343 establishment of a community of nitrifying bacteria in the predominately oxic sediment
344 layers (Wu et al., 2013). Autotrophic nitrifying bacteria use little energy for cell synthesis;
345 consequently, nitrifiers are typically slow to establish due to slow growth rates compared to
346 heterotrophic bacteria (Ebeling et al., 2006). The trends in the sediment redox data support
347 this hypothesis, since the redox potential in the first two months of the experiment reflected
348 conditions suitable for nitrification (Gerardi, 2002). The stabilization of nitrate concentrations
349 between months two and three may indicate that the nitrification capacity reached steady
350 state with the onset of coupled nitrification-denitrification as the redox potential indicated
351 conditions suitable for denitrification (+50 mV to -50 mV) in month two (Gerardi, 2002). The
352 quantity of organic carbon (i.e. carbon loading) is one of the main controls on the
353 denitrification efficiencies of sediments since this is a heterotrophic pathway of anaerobic
354 nitrate respiration (Blackburn and Blackburn, 1992; Joye and Anderson, 2008). The
355 decreasing nitrate concentrations in the final month are consistent with the conversion to di-
356 nitrogen gas by denitrifying bacteria under anoxic conditions indicated by the negative
357 sediment oxidation potentials recorded at the end of the trial (Blackburn and Blackburn,
358 1992; Seitzinger, 1988).

359 The conditions under which the organic carbon source is degraded (oxic or anaerobic) will
360 also affect the decomposition rate since bacterial growth efficiencies are a direct function of
361 oxygen availability (Fenchel et al., 2012; Goldman et al., 1987; Tezuka, 1990). In the present
362 study, the C:N of the aquaculture waste was increased from 5:1 to 20:1. For heterotrophic
363 bacteria (average C:N of 5:1), a substrate C:N of 20:1 represents the threshold for net
364 removal or net regeneration of ammonium at a bacterial growth efficiency of 25%. Under
365 anaerobic conditions, such as redox-stratified sediments, bacterial growth efficiencies range
366 from 5–30% (Goldman et al., 1987). It is possible that the C:N of 20:1 was not sufficiently
367 high to mediate a shift in nitrogen cycling pathways to promote net NH_4^+ assimilation due to
368 lower bacterial growth efficiencies under reducing conditions. Knowledge of the sediment
369 redox potential and how it changes over time in response to waste and carbon addition is
370 therefore a pre-requisite for determining the optimal C:N to assure the assimilation of NH_4^+
371 into bacterial biomass.

372 The biochemical composition of substrate is an important factor in defining resource quality
373 since the degradability is linked to its structural complexity. The carbon sources tested in this
374 study differed in their biodegradability; from glucose, a labile carbon source with a first-order
375 degradation rate constant of 1.15 d^{-1} , to cellulose, a complex structural polysaccharide with a
376 slow degradation rate of 0.05 d^{-1} (Avnimelech et al., 1995; Reddy et al., 1986). Readily
377 biodegradable substrates such as glucose are effective in promoting heterotrophic bacterial
378 growth under aerated conditions in biofloc systems (Crab et al., 2010); however, in this study
379 glucose had little impact on sea cucumber growth. The fundamental concept of C:N control
380 underpins biofloc technology where carbon-rich substrates are added to aquaculture
381 production tanks to overcome carbon limitation in heterotrophic bacteria and control the
382 build-up of inorganic nitrogen in the water column (Avnimelech, 1999). Biofloc uses high
383 aeration rates to maintain dense flocs in suspension (Crab et al., 2012). For this reason,

carbon sources tend to be highly labile and soluble and include molasses, glucose, sucrose, glycerol and acetate (Avnimelech, 2014; Crab et al., 2010). In contrast, the biomass density curve of sea cucumber juveniles reared in tanks amended with cellulose did not show any signs of plateauing, indicating that the maximum system carrying capacity was not reached. Had the experiment continued for a further two months the cellulose treatment may have outperformed the starch treatment. In a redox-stratified sediment-based system where heterotrophic bacteria have lower growth efficiencies, more complex (i.e. refractory) polysaccharides may be a more suitable long-term substrate. Cellulose is more resistant to hydrolysis as its β -1,4 bonds form rigid, ribbon-like chains with crystalline structures (Fenchel et al., 2012). Since the slow hydrolysis of more stable polysaccharides such as cellulose, demands an incorporation of nitrogen and phosphorus into microbial cells (Schroeder, 1987), the use of a more complex compound may offer a more stable means of ensuring sufficient carbon availability to bacteria in the long-term. This is particularly relevant in microbial-deposit feeder aquaculture bioremediation systems (see (Robinson et al., 2018)), due to the need to balance the constant efflux of NH_4^+ issuing from the redox-stratified sediment, sea cucumber excretion and ammonification following waste addition. The addition of cellulose would also approximate the natural habitat of *H. scabra* that comprises seagrass beds in areas with high terrigenous input (Hamel et al., 2001). This laboratory-scale observation highlights the potential to further increase the environmental and possibly the economic sustainability of integrated aquaculture production and bioremediation systems by utilising cheaper sources of complex carbon, including agricultural by-products such as bagasse or biochar (Srinivasva, 1987).

5. Conclusions

Carbon addition is utilised in other aquaculture treatment technologies including biofloc, denitrifying reactors and treatment of saline sludge by anaerobic digestion (Avnimelech, 2014; Hamlin et al., 2008; Luo et al., 2015; van Rijn et al., 2006). While these technologies utilise low cost carbon sources, they focus on the permanent removal of nitrogen through denitrification, or generate additional solid wastes that require disposal. Carbon supplementation may offer a more sustainable alternative to retain nitrogen in the system by promoting the net immobilisation of NH_4^+ into single cell biomass that can be up-cycled into high value secondary biomass. The potential for high-density culture is therefore an attractive approach to closing the nitrogen loop, especially if waste streams from aquaculture and agriculture are combined.

Acknowledgements: This research was funded by a Biotechnology and Biological Sciences Research Council (BBSRC) Industrial CASE Studentship to G.R. (Grant Code BB/J01141X/1) with HIK Abalone Farm Pty Ltd as the CASE partner, and with additional contributions from the THRIP program of the National Research Foundation, South Africa (Grant Number TP2011070800007). The work was conceptualised and funding was secured by G.R., C.L.W.J. and S.M.S. Experiments were performed by G.R. Data were analysed by G.R., and G.S.C. The manuscript was written by G.R. and G.S.C. and edited by C.L.W.J. and S.M.S. All authors have approved the final article.

The authors declare no competing financial interests.

References

Alongi, D.M., Hanson, R.B., 1985. Effect of detritus supply on trophic relationships within experimental benthic food webs. II. Microbial responses, fate and composition of decomposing detritus. *J. Exp. Mar. Biol. Ecol.* 88, 167-182.

434 Avnimelech, Y., 1999. Carbon/nitrogen ratio as a control element in aquaculture systems.
435 Aquaculture 176, 227-235.

436 Avnimelech, Y., 2014. Biofloc Technology - A Practical Guidebook, 3rd Edition ed.

437 Avnimelech, Y., Mozes, N., Diab, S., Kochba, M., 1995. Rates of organic carbon and
438 nitrogen degradation in intensive fish ponds. Aquaculture 134, 211-216.

439 Azim, M.E., Little, D.C., Bron, J.E., 2008. Microbial protein production in activated
440 suspension tanks manipulating C:N ratio in feed and the implications for fish culture. Biores.
441 Technol. 99, 3590-3599.

442 Badiola, M., Mendiola, D., Bostock, J., 2012. Recirculating Aquaculture Systems (RAS)
443 analysis: main issues on management and future challenges. Aquacult. Eng. 51, 26-35.

444 Blackburn, T.H., Blackburn, N.D., 1992. Model of nitrification and denitrification in marine
445 sediments. FEMS Microbiol. Lett. 100, 517-521.

446 Bossers, S., 2015. The Sea Cucumber Microbiome. CCMAR, Universidade do Algarve and
447 University of Amsterdam., Amsterdam, p. 31.

448 Brown, N., Eddy, S., Plaud, S., 2011. Utilization of waste from a marine recirculating fish
449 culture system as a feed source for the polychaete worm, *Nereis virens*. Aquaculture 322-323,
450 177-183.

451 Chávez-Crooker, P., Obrique-Contreras, J., 2010. Bioremediation of aquaculture wastes.
452 Curr. Opin. Biotechnol. 21, 313-317.

453 Church, M.J., 2008. Resource control of bacterial dynamics in the sea, in: Kirchman, D.L.
454 (Ed.), Microbial Ecology of the Oceans, 2nd ed. John Wiley and Sons, USA, pp. 455-493.

455 Crab, R., Chielens, B., Wille, M., Bossier, P., Verstraete, W., 2010. The effect of different
456 carbon sources on the nutritional value of bioflocs, a feed for *Macrobrachium rosenbergii*
457 post larvae. Aquacult. Res. 41, 559-567.

458 Crab, R., Defoirdt, T., Bossier, P., Verstraete, W., 2012. Biofloc technology in aquaculture:
459 beneficial effects and future challenges. Aquaculture 356-357, 351-356.

460 Cubillo, A.M., Ferreira, J.G., Robinson, S.M.C., Pearce, C.M., Corner, R.A., Johansen, J.,
461 2016. Role of deposit feeders in integrated multi-trophic aquaculture — A model analysis.
462 Aquaculture 453, 54-66.

463 Ebeling, J.M., Timmons, M.B., Bisogni, J.J., 2006. Engineering analysis of the stoichiometry
464 of photoautotrophic, autotrophic, and heterotrophic removal of ammonia–nitrogen in
465 aquaculture systems. Aquaculture 257, 346-358.

466 Erler, D., Pollard, P.C., Knibb, W., 2004. Effects of secondary crops on bacterial growth and
467 nitrogen removal in shrimp farm effluent treatment systems. Aquacult. Eng. 30, 103-114.

468 Fenchel, T., Blackburn, T.H., 1979. Bacteria and Mineral Cycling. Academic Press, London,
469 UK.

470 Fenchel, T., King, G.M., Blackburn, T.H., 2012. Bacterial Biogeochemistry. The
471 Ecophysiology of Mineral Cycling, 3rd ed. Academic Press, San Diego.

472 Fontaine, A.R., Chia, F.S., 1968. Echinoderms: an autoradiographic study of assimilation of
473 dissolved organic molecules. Science (New York, N.Y.) 161, 1153-1155.

474 Gerardi, M.H., 2002. Nitrification and Denitrification in the Activated Sludge Process. John
475 Wiley and Sons, New York, USA.

476 Goldman, J.C., Caron, D.A., Dennett, M.R., 1987. Regulation of gross growth efficiency and
477 ammonium regeneration of bacteria by substrate C:N ratio. Limnol. Oceanogr. 32, 1239-
478 1252.

479 Hamel, J.-F., Mercier, A., Conand, C., Purcell, S., Toral-Granda, T.-G., Gamboa, R., 2013.
480 *Holothuria scabra*. The IUCN Red List of Threatened Species 2013.

481 Hamel, J.F., Conand, C., Pawson, D.L., Mercier, A., 2001. The sea cucumber *Holothuria*
482 *scabra* (Holothuroidea : Echinodermata): Its biology and exploitation as beche-de-mer. Adv.
483 Mar. Biol. 41, 129-223.

484 Hamlin, H.J., Michaels, J.T., Beaulaton, C.M., Graham, W.F., Dutt, W., Steinbach, P.,
485 Losordo, T.M., Schrader, K.K., Main, K.L., 2008. Comparing denitrification rates and carbon
486 sources in commercial scale upflow denitrification biological filters in aquaculture. Aquacult.
487 Eng. 38, 79-92.

488 Han, Q., Keesing, J.K., Liu, D., 2016. A review of sea cucumber aquaculture, ranching, and
489 stock enhancement in China. Rev. Fish. Sci. Aquacult. 24, 326-341.

490 Hargreaves, J.A., 1998. Nitrogen biogeochemistry of aquaculture ponds. Aquaculture 166,
491 181-212.

492 Herbert, D., 1967. Stoichiometric aspects of microbial growth, in: Dean, A.C.R., Ellwood,
493 D.C., Evans, C.G.T., Melling, J. (Eds.), Continuous culture 6: Applications and New Fields.
494 Horwood, Chichester, U.K., pp. 1-27.

495 Huettel, M., Berg, P., Kostka, J.E., 2014. Benthic exchange and biogeochemical cycling in
496 permeable sediments. Ann. Rev. Mar. Sci. 6, 23-51.

497 Jaeckle, W.B., Strathmann, R.R., 2013. The anus as a second mouth: anal suspension feeding
498 by an oral deposit-feeding sea cucumber. Invert. Biol. 132, 62-68.

499 Jangoux, M., Lawrence, J.M., 1982. Echinoderm Nutrition, Balkema, Rotterdam.

500 Joye, S.B., Anderson, I.C., 2008. Chapter 19 - Nitrogen Cycling in Coastal Sediments,
501 Nitrogen in the Marine Environment (2nd Edition). Academic Press, San Diego, pp. 867-915.

502 Kang, K.H., Kwon, J.Y., Kim, Y.M., 2003. A beneficial coculture: charm abalone *Haliotis*
503 *discus hannai* and sea cucumber *Stichopus japonicus*. Aquaculture 216, 87-93.

504 Kirchman, D.L., 2012. Processes in Microbial Ecology. Oxford University Press, USA, 326
505 p.

506 Krishnan, S., Krishnaswamy, S., 1970. Studies on the transport of sugars in the holothurian
507 *Holothuria scabra*. Mar. Biol. 5, 303-306.

508 Lawrence, J.M., 1982. Digestion, in: Jangoux, M., Lawrence, J.M. (Eds.), Echinoderm
509 Nutrition. Balkema, Rotterdam, Netherlands, pp. 283-316.

510 Lehtoranta, J., Ekholm, P., Pitkanen, H., 2009. Coastal eutrophication thresholds: a matter of
511 sediment microbial processes. Ambio 38, 303-308.

512 Levene, H., 1960. Robust tests for equality of variances, in: Olkin, I., Hotelling, H. (Eds.),
513 Contributions to Probability and Statistics: Essays in Honor of Harold Hotelling. Stanford
514 University Press.

515 Lin, G., Ji, Y., Gou, J., Sun, G., 1993. Studies on over-wintering technique of polyculture
516 abalone and sea cucumber by means of cooling seawater from coastal power plant. Mar. Sci.
517 1, 4-7.

518 Lopez, G.R., Levinton, J.S., 1987. Ecology of deposit-feeding animals in marine sediments.
519 Quart. Rev. Biol. 62, 235-260.

520 Luo, G.-z., Ma, N., Li, P., Tan, H.-x., Liu, W., 2015. Enhancement of anaerobic digestion to
521 treat saline sludge from recirculating aquaculture systems. Sci. World J. 2015, 479101.

522 MacDonald, C.L.E., Stead, S.M., Slater, M.J., 2013. Consumption and remediation of
523 European Seabass (*Dicentrarchus labrax*) waste by the sea cucumber *Holothuria forskali*.
524 Aquacult. Int. 21, 1279-1290.

525 Maxwell, K.H., Gardner, J.P.A., Heath, P.L., 2009. The effect of diet on the energy budget of
526 the brown sea cucumber, *Stichopus mollis* (Hutton). J. World Aquacult. Soc. 40, 157-170.

527 Mirzoyan, N., McDonald, R.C., Gross, A., 2012. Anaerobic treatment of brackishwater
528 aquaculture sludge: an alternative to waste stabilization ponds. J. World Aquacult. Soc. 43,
529 238-248.

530 Naylor, R.L., Hardy, R.W., Bureau, D.P., Chiu, A., Elliott, M., Farrell, A.P., Forster, I.,
531 Gatlin, D.M., Goldburg, R.J., Hua, K., Nichols, P.D., 2009. Feeding aquaculture in an era of
532 finite resources. Proc. Nat. Acad. Sci. U.S.A. 106, 15103-15110.

533 Palmer, P.J., 2010. Polychaete-assisted sand filters. Aquaculture 306, 369-377.

534 Palmer, P.J., Wang, S., Houlihan, A., Brock, I., 2014. Nutritional status of a nereidid
535 polychaete cultured in sand filters of mariculture wastewater. Aquacult. Nutrition 20, 675-
536 691.

537 Purcell, S.W., 2010. Managing sea cucumber fisheries with an ecosystem approach, FAO
538 Fisheries and Aquaculture Technical Paper. FAO, Rome, Italy, p. 157.

539 Purcell, S.W., 2014. Value, market preferences and trade of beche-de-mer from Pacific island
540 sea cucumbers. PLoS ONE 9, e95075.

541 Purcell, S.W., Hair, C.A., Mills, D.J., 2012a. Sea cucumber culture, farming and sea ranching
542 in the tropics: Progress, problems and opportunities. Aquaculture 368-369, 68-81.

543 Purcell, S.W., Samyn, Y., Conand, C., 2012b. Commercially important sea cucumbers of the
544 world. FAO, Rome, Italy, p. 150.

545 Reddy, K.R., Feijtel, T.C., W.H., P.J., 1986. Effect of soil redox conditions on microbial
546 oxidation of organic matter, in: Chen, Y., Avnimelech, Y. (Eds.), The role of organic matter
547 in modern agriculture. Martinus, Nijhoff, Dordrecht, pp. 117-156.

548 Rice, D.L., Rhoads, D.C., 1989. Early diagenesis of organic matter and the nutritional value
549 of sediment. *Lect. Notes Coast. Estuar. Stud.* 31, 59-97.

550 Rittmann, B.E., McCarty, P.L., 2001. *Environmental Biotechnology: Principles and*
551 *Applications*. McGraw-Hill, New York, USA.

552 Robinson, G., 2013. A bright future for sandfish aquaculture, *World Aquacult. Soc. Mag.* pp.
553 18-24.

554 Robinson, G., Caldwell, G.S., Jones, C.L.W., Slater, M.J., Stead, S.M., 2015. Redox
555 stratification drives enhanced growth in a deposit-feeding invertebrate: implications for
556 aquaculture bioremediation. *Aquacult. Environ. Interact.* 8, 1-13.

557 Robinson, G., Caldwell, G.S., Wade, M.J., Free, A., Jones, C.L.W., Stead, S.M., 2016.
558 Profiling bacterial communities associated with sediment-based aquaculture bioremediation
559 systems under contrasting redox regimes. *Sci. Rep.* 6, 38850.

560 Robinson, G., MacTavish, T., Savage, C., Caldwell, G. S., Jones, C. L. W., Probyn, T., Eyre,
561 B. D., and Stead, S. M. (2018). Carbon amendment stimulates benthic nitrogen cycling
562 during the bioremediation of particulate aquaculture waste, *Biogeosciences*, 15, 1863-1878,
563 2018.

564

565 Russell-Hunter, W.D., 1970. *Aquatic Productivity*. Macmillan Publishers, New York.

566 Schneider, O., Chabrilion-Popelka, M., Smidt H., Haenen, O., Sereti, V., Eding, E.H.,
567 Verreth, J.A.J., 2007a. HRT and nutrients affect bacterial communities grown on
568 recirculation aquaculture system effluents. *FEMS Microbiol. Ecol.* 60, 207-219.

569 Schneider, O., Sereti, V., Eding, E.H., Verreth, J.A., 2007b. Heterotrophic bacterial
570 production on solid fish waste: TAN and nitrate as nitrogen source under practical RAS
571 conditions. *Biores. Technol.* 98, 1924-1930.

572 Schneider, O., Sereti, V., Eding, E.H., Verreth, J.A.J., 2005. Analysis of nutrient flows in
573 integrated intensive aquaculture systems. *Aquacult. Eng.* 32, 379-401.

574 Schneider, O., Sereti, V., Eding, E.H., Verreth, J.A.J., 2006. Molasses as C source for
575 heterotrophic bacteria production on solid fish waste. *Aquaculture* 261, 1239-1248.

576 Schroeder, G.L., 1987. Carbon pathways in aquatic detrital systems, in: Moriarty, D.J.W.,
577 Pullin, R.S.V. (Eds.), *Detritus and microbial ecology in aquaculture*. International Center for
578 Living Aquatic Resources Management, Manila, Philippines, p. 420.

579 Seitzinger, S.P., 1988. Denitrification in freshwater and coastal marine ecosystems:
580 ecological and geochemical significance. *Limnol. Oceanogr.* 33, 702-724.

581 Shapiro, S.S., Wilk, M.B., 1965. An analysis of variance test for normality (complete
582 samples). *Biometrika* 52, 591-611.

583 Shi, C., Dong, S., Wang, F., Gao, Q., Tian, X., 2013. Effects of four fresh microalgae in diet
584 on growth and energy budget of juvenile sea cucumber *Apostichopus japonicus* (Selenka).
585 *Aquaculture* 416-417, 296-301.

586 Srinivasva, R.N., 1987. Convemion of cellulosic and other organic wastes into microbial
587 proteins, in: Moriarty, D.J.W., Pullin, R.S.V. (Eds.), *Detritus and microbial ecology in*
588 *aquaculture*. International Center for Living Aquatic Resources Management, Manila,
589 Philippines, pp. 181-191.

590 Stuart, R.K., Mayali, X., Lee, J.Z., Craig Everroad, R., Hwang, M., Bebout, B.M., Weber,
591 P.K., Pett-Ridge, J., Thelen, M.P., 2016. Cyanobacterial reuse of extracellular organic carbon
592 in microbial mats. *ISME J* 10, 1240-1251.

593 Tenore, K.R., Chesney, E.J., 1985. The effects of interaction of rate of food supply and
594 population density on the bioenergetics of the opportunistic polychaete, *Capitella capitata*
595 (type 1). *Limnol. Oceanogr.* 30, 1188-1195.

596 Tezuka, Y., 1990. Bacterial regeneration of ammonium and phosphate as affected by the
597 carbon:nitrogen:phosphorus ratio of organic substrates. *Microb. Ecol.* 19, 227-238.

598 Underwood, G.J.C., Paterson, D.A., Parkes, R.J., 1995. The measurement of microbial
599 carbohydrate exopolymers from intertidal sediments. *Limnol. Oceanogr.* 40, 1243-1253.

600 Uthicke, S., Klumpp, D.W., 1997. Ammonium excretion by holothurians enhances
601 production and turnover of benthic diatom communities, *Proceedings of the 8th International*
602 *Coral Reef Symposium*, Smithsonian Tropical Research Institute, Panama, pp. 873-876.

603 van Rijn, J., Tal, Y., Schreier, H.J., 2006. Denitrification in recirculating systems: theory and
604 applications. *Aquacult. Eng.* 34, 364-376.

605 Wu, B., Xia, S., Rahman, M.M., Rajkumar, M., Fu, Z., Tan, J., Yang, A., 2015. Substituting
606 seaweed with corn leaf in diet of sea cucumber (*Apostichopus japonicus*): Effects on growth,
607 feed conversion ratio and feed digestability. *Aquaculture* 444, 88-92.

608 Wu, Y., Ke, X., Hernández, M., Wang, B., Dumont, M.G., Jia, Z., Conrad, R., 2013.
609 Autotrophic Growth of Bacterial and Archaeal Ammonia Oxidizers in Freshwater Sediment
610 Microcosms Incubated at Different Temperatures. *Appl. Environ. Microbiol.* 79, 3076-3084.

611 Yearsley, R.D., Jones, C.L.W., Britz, P.J., 2009. Effect of settled sludge on dissolved
612 ammonia concentration in tanks used to grow abalone (*Haliotis midae* L.) fed a formulated
613 diet. *Aquacult. Res.* 40, 166-171.

614 Zamora, L.N., Yuan, X., Carton, A.G., Slater, M.J., 2016. Role of deposit-feeding sea
615 cucumbers in integrated multitrophic aquaculture: progress, problems, potential and future
616 challenges. *Rev. Aquacult.* doi: 10.1111/raq.12147
617

618

619

620 **Figure legends.**

621 **Figure 1.** The mean (\pm standard error) (A) pH, (B) total ammonia-nitrogen (TAN) and (C)
622 nitrate concentration of the water in sea cucumber tanks dosed with aquaculture waste only
623 (none) or with aquaculture waste amended with various carbon sources (i.e. glucose, starch or
624 cellulose).

625 **Figure 2.** The mean (\pm standard error) (A) reduction-oxidation (redox) potential at the base
626 of the sediment (4 cm deep), (B) organic carbon content (%) measured in the surface
627 sediment, (C) total nitrogen content (%) measured in the surface sediment and (D) carbon to
628 nitrogen ratios (C:N) measured in the surface sediment in sea cucumber tanks dosed with
629 aquaculture waste only (none) or with aquaculture waste amended with various carbon
630 sources (i.e. glucose, starch or cellulose).

631 **Figure 3.** The mean (\pm standard error) (A) growth rate, (B) cumulative biomass density and
632 (C) wet weight of *Holothuria scabra* juveniles (n=4) reared on particulate aquaculture waste
633 (none) or with aquaculture waste amended with various carbon sources (i.e. glucose, starch or
634 cellulose) for four months.

635




Fig 1.





Fig 2.



643

644 Fig 3.

Table 1. Description of experimental treatments including daily additions of particulate aquaculture waste (2.41 g wet weight) and the different carbon sources standardised to 200 mmol C m⁻² d⁻¹. *First-order decomposition rate constant is the percentage of a given compound that degrades each day; **Half-life is the time it takes to reach 50% of the complete degradation of a given substrate concentration.

Carbon source	Solubility	*First-order decomposition rate (d ⁻¹)	**Half-life (d)	Carbon content (%)	Dry weight (g)	C:N
None	N/A	N/A	N/A	N/A	N/A	5:1
D-glucose	Soluble	1.1500	0.6	39.99	0.75	20:1
Starch	50 g L ⁻¹ at 90 °C	0.8000	5.0	44.44	0.68	20:1
Cellulose, microcrystalline	Insoluble (20 °C)	0.0495	14.0	44.44	0.68	20:1

Table 2. Mean (\pm standard error) of water and sediment quality parameters recorded during the four-month experiment. C:N = carbon to nitrogen ratio.

	None	Glucose	Starch	Cellulose	Overall
<i>Water quality</i>					
Light (lux)	139.25 \pm 5.22	141.25 \pm 5.22	141.25 \pm 5.22	136.35 \pm 5.41	139.53 \pm 2.59
Temperature (°C)	29.07 \pm 0.50	29.24 \pm 0.53	29.21 \pm 0.55	29.19 \pm 0.51	29.18 \pm 0.26
Dissolved oxygen (mg L ⁻¹)	7.28 \pm 0.17	7.19 \pm 0.12	7.47 \pm 0.14	7.11 \pm 0.14	7.26 \pm 0.07
Ammonia (mg L ⁻¹)	0.34 \pm 0.07	0.35 \pm 0.09	0.40 \pm 0.11	0.33 \pm 0.09	0.35 \pm 0.04
Nitrate (mg L ⁻¹)	3.98 \pm 0.41	4.09 \pm 0.43	3.91 \pm 0.40	3.68 \pm 0.38	3.92 \pm 0.20
<i>Sediment quality</i>					
Total carbohydrate ($\mu\text{g g}^{-1}$)	64.27 \pm 28.08	49.90 \pm 19.37	41.11 \pm 21.96	96.08 \pm 40.80	62.84 \pm 14.30
Chlorophyll <i>a</i> ($\mu\text{g g}^{-1}$)	0.86 \pm 0.56	0.85 \pm 0.37	1.29 \pm 0.54	2.67 \pm 1.19	1.42 \pm 0.37
Phaeopigment ($\mu\text{g g}^{-1}$)	1.08 \pm 0.77	0.97 \pm 0.49	3.04 \pm 1.10	4.76 \pm 2.15	2.46 \pm 0.66
Organic carbon (%)	0.12 \pm 0.02	0.12 \pm 0.02	0.11 \pm 0.02	0.20 \pm 0.05	0.14 \pm 0.01
Nitrogen (%)	0.01 \pm 0.00	0.01 \pm 0.00	0.02 \pm 0.00	0.02 \pm 0.00	0.02 \pm 0.00
C:N	8.97 \pm 0.44	8.43 \pm 0.43	7.41 \pm 0.23	9.84 \pm 0.51	8.66 \pm 0.23

Captions for supplementary tables:

Table S1: Supporting statistics for environmental and water quality parameter data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Table S2: Supporting statistics for sediment reduction-oxidation potential, total carbohydrate, chlorophyll *a* and phaeopigment data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Table S3: Supporting statistics for sediment organic carbon, nitrogen and carbon to nitrogen ratio data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Table S4: Supporting statistics for sea cucumber juvenile growth data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Supplementary:

**The effect of resource quality on the growth of *Holothuria scabra* during aquaculture
waste bioremediation**

Georgina Robinson^{1,2,#*}, Gary S. Caldwell¹, Clifford L.W. Jones², Selina M. Stead¹

¹School of Marine Science and Technology, Newcastle University, Newcastle, NE1 7RU, UK.

²Department of Ichthyology and Fisheries Science, Rhodes University, Grahamstown 6140, South Africa.

#Current address: Scottish Association for Marine Science, Scottish Marine Institute, PA37 1QA, Oban, UK

*Corresponding author. Tel +230 5982 4971; Email address Georgina.Robinson@sams.ac.uk (G. Robinson)

Supplementary information

Table S1: Supporting statistics for environmental and water quality parameter data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Source of variation	df	Light (lux)			Temperature (°C)			Dissolved oxygen (mg L ⁻¹)			pH		
		MS	F	p	MS	F	p	MS	F	P	MS	F	p
Carbon source (C)	3	8.75E+07	0.226	0.877	0.11	0.1	0.972	0.485	1.61	0.240	0.05	5.62	0.012*
Time (T)	4	4.02E+08	3.038	0.026*	93.44	190.7	0.000*	5.444	52.52	0.000*	0.575	67.4	0.000*
C x T	12	3.52E+07	0.265	0.992	0.06	0.1	1.000	0.081	0.78	0.663	0.024	2.79	0.006*

Table S2: Supporting statistics for sediment reduction-oxidation potential, total carbohydrate, chlorophyll *a* and phaeopigment data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Source of variation	df	Reduction-oxidation potential (mV)			Total carbohydrate (µg g ⁻¹)			Chlorophyll <i>a</i> (µg g ⁻¹)			Phaeopigment (µg g ⁻¹)	
		MS	F	p	MS	F	p	MS	F	p	MS	F
Carbon source (C)	3	3.62E+04	30.1	0.000*	1.16E+04	0.985	0.432	14.81	1.530	0.257	64.9	1.636
Time (T)	4	1.77E+05	251.9	0.000*	8.32E+04	5.373	0.001*	48.62	5.593	0.001*	136.3	5.244
C x T	12	3347	4.8	0.000*	3,323.00	0.214	0.997	7.51	0.864	0.587	21.8	0.840

Table S3: Supporting statistics for sediment organic carbon, nitrogen and carbon to nitrogen ratio data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Source of variation	df	Organic carbon (%)			Nitrogen (%)			C/N ratio		
		MS	F	p	MS	F	p	MS	F	p
Carbon source (C)	3	0.048	1.852	0.238	0.000	1.237	0.376	16.45	4.211	0.064
Time (T)	4	0.060	6.061	0.002*	0.001	5.877	0.002*	5.43	2.695	0.055
C x T	12	0.015	1.548	0.175	0.000	1.422	0.223	4.38	2.174	0.051

Table S4: Supporting statistics for sea cucumber growth data analysed by repeated measures ANOVA, $p < 0.05$ (* indicates significant differences).

Source of variation	Wet weight (g)				Biomass density (g m ⁻²)				Growth rate (g d ⁻¹)			
	df	MS	F	p	df	MS	F	p	df	MS	F	P
Carbon source (C)	3	1.16E+04	0.985	0.432	3	7.98E+04	4.1	0.031*	3	0.023	4.22	0.030*
Time (T)	4	8.32E+04	5.373	0.001*	4	1.50E+06	224.2	0.000*	3	0.126	44.53	0.000*
C x T	12	3,323.00	0.214	0.997	12	1.67E+04	2.5	0.013*	9	0.002	0.72	0.690