


RESEARCH ARTICLE

More miles on the clock: Neighbourhood stressors are associated with telomere length in a longitudinal study

Anne Ellaway^{1*}, Ruth Dundas¹, Tony Robertson², Paul G. Shiels³

1 MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, United Kingdom, **2** Centre for Public Health and Population Health Research, Faculty of Health Sciences & Sport, University of Stirling, Stirling, United Kingdom, **3** Institute of Cancer Sciences, University of Glasgow, Glasgow, United Kingdom

* anne.ellaway@glasgow.ac.uk



Abstract

Background

There is a substantial gap in health and longevity between more affluent and more deprived areas, and more knowledge of the determinants of this health divide is required. Experience of the local residential environment is important for health although few studies have examined this in relation to biological markers of age such as telomere length. We sought to examine if residents' perceptions of neighbourhood stressors over time were associated with telomere length in a community study.

Methodology/Principal findings

In a prospective cohort study of 2186 adults in the West of Scotland, we measured neighbourhood stressors at three time points over a 12-year period and telomere length at the end of the study. Using linear regression models, we found that a higher accumulation of neighbourhood stressors over time was associated with shorter telomere length, even after taking cohort, social class, health behaviours (smoking status, diet, physical activity), BMI and depression into account among females only (Beta = 0.007; 95%CI [0.001, 0.012]; $P < 0.014$).

Conclusions/Significance

Neighborhood environments are potentially modifiable, and future efforts directed towards improving deleterious local environments may be useful to lessen telomere attrition.

Introduction

A number of studies have shown that residents' perceptions of the local environment are associated with their health, independently of individual characteristics such as socioeconomic

OPEN ACCESS

Citation: Ellaway A, Dundas R, Robertson T, Shiels PG (2019) More miles on the clock: Neighbourhood stressors are associated with telomere length in a longitudinal study. PLoS ONE 14(3): e0214380. <https://doi.org/10.1371/journal.pone.0214380>

Editor: Adewale L. Oyeyemi, University of Maiduguri College of Medical Sciences, NIGERIA

Received: September 13, 2018

Accepted: March 12, 2019

Published: March 28, 2019

Copyright: © 2019 Ellaway et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data underlying this study cannot be shared publicly in order to protect participant privacy. Interested and qualified researchers can request the data through application to the MRC/CSO Social and Public Health Sciences Unit. For instructions and details on requesting access to the study data, please see: <http://2007study.sphsu.mrc.ac.uk/Information-on-data-sharing.html>. Please contact Professor Anne Ellaway for further information: Anne.Ellaway@glasgow.ac.uk.

Funding: The West of Scotland: Twenty-07 Study is funded by the UK Medical Research Council (MRC; MC_A540_53462). A.E. and R.D. are funded by the Medical Research Council (MC_UU_12017/10; MC_UU_12017/13) and the Chief Scientist's Office (SPHSU10; SPHSU13). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

status (SES), age and gender [1–5]. However, the majority of the findings to date are based on self-reported health outcomes, leaving open the possibility of 'plaintive-set' reporting bias [6]. Moreover, given the substantial gap in health and longevity between more affluent and more deprived areas [7, 8], more knowledge on the determining factors of poorer health in deprived areas, particularly in relation to biological mechanisms over the life course [9] are required to inform interventions [10]. It has been suggested that neighbourhood perceptions are important to examine as they can stimulate psychological and/or physiological stress responses that can affect mental and physical health [11]. One measure, which has received increased attention in relation to life stress and accelerated aging, is telomere shortening [12, 13].

Telomeres are specialised nucleoprotein complexes at the end of chromosomes which maintain genomic integrity by providing a cap for the chromosome, thus preventing chromosomal fusions [14]. Telomeres typically shorten during somatic cell division as a consequence of the 'end replication problem' [15, 16]. Telomere length has been posited as good marker for aging [17] and shorter telomeres have been observed by individual characteristics such as sex, among males compared to females [18] and lower SES in some populations [13, 19] but not others [20]. Shortened telomere length has been associated with exposure to chronic psychological stress [21], depression [22] and a range of non-communicable diseases including cancer [23] chronic kidney disease and diabetes [24]. As telomere length is modifiable [25] [23], a more nuanced understanding of the underlying mechanisms which facilitate accelerated telomere shortening can provide useful information to inform potential interventions that may mitigate the rate of telomeric attrition [10].

To date, only a small number of studies have examined whether exposure to neighbourhood conditions are associated with telomere length. In the USA, shorter telomeres were associated with perceived neighbourhood safety and social cohesion in the Multi-Ethnic Study of Atherosclerosis (MESA) [26]; perceived neighbourhood problems such as heavy traffic and excessive noise in a sample of African American females [27]; and in the Netherlands, perceived neighbourhood disorder (vandalism), fear of crime and noise were associated with telomere shortening [11]. As we have previously shown that social housing tenure is associated with shorter telomere length [28], and given that social housing tenure may expose individuals to poorer neighbourhood quality [29] it is key to explore whether neighbourhood quality is part of the pathway linking housing tenure to telomere attrition. Moreover, as most studies to date which have examined perceptions of the neighbourhood and telomere length are cross sectional, there have been calls to examine the effects of cumulative exposures to adverse neighbourhood conditions on health [30]. As others have discussed [10, 11, 26, 31], prolonged exposure to stressful conditions may lead to chronic activation of the hypothalamic–pituitary–adrenal (HPA) axis, which may influence the rate of telomere shortening, through changes in hallmark processes of ageing [32, 33]. Indirect mechanisms whereby chronic stress may invoke telomere shortening are related to the exposome [34] and include unhealthy behaviours as coping strategies such as the consumption of 'comfort' foods or smoking to alleviate stress. In this study, we have examined whether perceived neighbourhood stressors were associated with telomere length using three sweeps of data (hereby referred to as 'waves'), covering 13 years of the life course, from three cohorts in the longitudinal West of Scotland Twenty-07 Study. We therefore assessed two models [35] to examine the impact of perceived neighbourhood stressors on telomere length. These comprised: (i) an accumulation model which uses a cumulative measure of neighbourhood perceptions over time; (ii) a critical periods model which separately analyses the relationship between measures of perceived neighbourhood stressors at each wave and subsequent telomere length. As a number of studies have shown that neighbourhood conditions may impact differently upon the health of males and females [1, 36–39] we conduct our analyses separately by sex.

Methods

Data were from the West of Scotland Twenty-07 Study, a community-based, prospective cohort study, which has followed three cohorts of men and women recruited (one individual per household) in 1987 at the (approximate) ages of 15 ('1970s cohort'), 35 ('1950s cohort') and 55 years ('1930s cohort') in 1987 (wave 1) and the same individuals were followed up in a further four waves over the next 20 years until 2007/8. Baseline respondents have been shown to be representative of the general population of the sampled area [40]. At Wave 5, 83.6% of those eligible (i.e. still living and contactable) to take part participated in an interview. As expected in a general population longitudinal survey, men, those in the oldest cohort and people from manual occupations were more likely to have died. The study design is described in more detail elsewhere [41]. The Tayside Committee on Medical Research Ethics approved the study. Informed, written consent was obtained from all respondents at each wave of the study. Here, we use three waves of the study for which we have data relevant to our research question in this paper (wave 3, 4 & 5). Data, including blood samples at wave 5 (2007/8, $n = 2604$), were collected by trained nurses in the homes of the study participants when respondents were aged approximately 35 (1970s cohort), 55 (1950s cohort) and 75 years (1930s cohort).

Individual level variables

Household social class (an accepted measure of socioeconomic position [42]) was coded into six classifications I- VI (I Professional, II Managerial and Technical, III skilled non manual, IV skilled Manual, V partly skilled occupations, VI unskilled occupation) according to the Registrar General's 1980 classification [43] for head of household's current or previous occupation. Depression, which was added as a covariate in models to partly account for the potential of poor mental health influencing perceptions of the local environment, and as it has been shown to be related to telomere length [22], was measured by the 7 item depression subscale of the Hospital Anxiety and Depression Scale [HADS], higher scores on HADS indicate greater reported symptoms. [44]. As this instrument was not available for all respondents who participated at Wave 3, a single item ('feeling unhappy or depressed over the previous week') from the 12 item General Health Questionnaire (GHQ-12 [45]) was used instead for critical period (cross sectional). Health behaviours, as correlates of telomere length, were assessed at Wave 5, include diet (consumption of sweets and chocolate) [46], smoking status [47], physical activity ('In an average week, on how many days do you spend at least 20 continuous minutes doing vigorous physical exercise, enough to make you sweaty and out of breath?') [48] and Body Mass Index (BMI: $\text{weight [kg]/height[m]}^2$) [49].

Biospecimen collection procedure

At Wave 5, respondents' blood was drawn by venepuncture from the median cubital vein (major arm vein) at the end of the interview. Whole blood (4 ml) was collected in potassium ethylenediaminetetraacetic acid vacutainer tubes (Becton Dickinson, Franklin Lakes, NJ, USA). The tubes were stored at 4°C until DNA extraction. DNA was extracted from peripheral blood leukocytes using the Maxwell automated purification system, according to the manufacturer's instructions (Promega, Fitchburg, WI, USA). The DNA concentration and purity were quantified using the Nanodrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA).

Telomere length

As previously reported [20, 22, 28], telomere length was analysed at the Institute of Cancer Sciences, University of Glasgow. Telomere lengths in the DNA extracted from peripheral blood

leukocytes were determined by quantitative polymerase chain reaction, following the method of Cawthon [50]. Telomere length determination was performed blindly using a Roche Light Cycler LC480. Telomere length analyses were performed in triplicate for each sample, using a single-copy gene amplicon primer set (acidic ribosomal phosphoprotein, 36B4) and a telomere-specific amplicon primer set [51]. Quality control parameters used for the amplifications comprised a cut-off of 0.15 for the standard deviation (SD) of the threshold cycle (Ct) for sample replicates. At an SD above 0.15, the sample was reanalyzed. The average SD across plates was 0.05. A control DNA sample was used to generate a standard curve on every plate to enable absolute quantification analysis. Relative telomere length was estimated from Ct scores using the comparative Ct method after confirming that the telomere and control gene assays yielded similar amplification efficiencies. This method determines the ratio of telomere repeat-copy number to single-copy gene number (T/S) in experimental samples relative to a control sample DNA. This normalized T/S ratio was used as the estimate of relative telomere length (relative T/S). It should be noted that the same control DNA sample was used on every plate that was analyzed. The intra-assay variation was assessed by comparing the relative telomere estimates (T/S ratio) across assays for the positive controls, which were assayed on every assay plate. The average intra-assay coefficient of variance was 0.56% for telomere and 0.19% for 36B4 plates [13, 50, 52].

Perceptions of neighbourhood quality

A similar suite of questions on respondents' perceptions of their local neighbourhood was asked at three waves of the study: wave 3 (1995), wave 4 (2000) and wave 5 (2007). Respondents were asked 'Around where you live would you say that any of the following are a serious problem, a minor problem or not a problem' and provided with a list of six socio-environmental problems: vandalism, litter and rubbish, assaults and muggings, disturbances by children or youngsters, smells and fumes, burglaries and invited to reply using a three-point scale ('not a problem' score 1, 'minor problem' score 2, 'serious problem' score 3). A similar suite of questions has been used in other UK studies including the General Household Survey and the Hereford Cohort Study [53, 54]. At each wave, to provide a more differentiated score on the perceived severity of the problems [55, 56], scores were revised so that 'serious problem' was given a score of one and 'minor problem' and 'not a problem' were each scored as zero. Responses to each of the individual problems were then summed, with scores ranging from 0 to 6 at each wave, with the higher score indicating greater problems in the neighbourhood. To create a cumulative measure of neighbourhood perceptions over time, scores were summed across the three waves and scores for this variable ranged from 0 to 15. All neighbourhood perceptions scores were measured and analysed at the individual level.

Statistical analysis

Associations between T/S (relative telomere length) and two models ('accumulation', critical) of perceptions of the neighbourhood were analyzed, separately for males and females, using linear regression models, adjusted for cohort, household social class, the HADS measure of depression, diet [46], smoking status [47], physical [48] and Body Mass Index [49]. All analyses were performed with IBM SPSS 21.

Results

Of the 2604 study participants who took part at wave 5, 2310 (88.7%) consented to have telomere analysis performed; 2193 (94.9%) participants had sufficient blood for DNA to be extracted. Telomere length was successfully measured for 2186 (94.6%) (1970s cohort $n = 776$;

Table 1. Mean telomere length by sex, age cohort and social class.

	n	%	T/S ratio Mean (SD), p<	Accumulation of neighbourhood problems mean (SD)	Neighbourhood problems Wave 3 mean (SD)	Neighbourhood problems Wave 4 mean (SD)	Neighbourhood problems Wave 5 mean (SD)
sex			P< 0.021				
male	994	45.6	0.78 (0.21)	1.16 (2.07)	0.43 (0.96)	0.36 (0.91)	0.38 (0.94)
females	1192	54.4	0.80 (0.20)	1.31 (2.14)	0.49 (1.03)	0.38 (0.86)	0.44 (0.98)
cohort			P< 0.001			P<0.020	
1970s	776	35.4	0.86 (0.21)	1.20 (1.96)	0.48 (1.05)	0.33 (0.86)	0.40 (0.92)
1950s	866	39.7	0.78 (0.19)	1.20 (2.15)	0.44 (0.98)	0.35 (0.84)	0.42 (0.98)
1930s	544	25.0	0.70 (0.19)	1.37 (2.24)	0.48 (0.96)	0.46 (0.99)	0.43 (1.00)
Head of Household social class			P< 0.001	P< 0.001	P< 0.001	P< 0.001	P< 0.001
I Professional	266		0.81 (0.21)	0.71 (1.27)	0.27 (0.67)	0.23 (0.61)	0.22 (0.60)
II managerial and technical	896		0.81 (0.21)	0.99 (1.81)	0.40 (0.94)	0.30 (0.77)	0.30 (0.77)
III skilled non manual	457		0.78 (0.19)	1.35 (2.13)	0.48 (0.98)	0.41 (0.89)	0.46 (1.04)
III skilled manual	300		0.77 (0.20)	1.58 (2.61)	0.60 (1.19)	0.40 (1.02)	0.60 (1.17)
IV partly skilled	199		0.75 (0.21)	2.08 (2.67)	0.67 (1.23)	0.64 (1.23)	0.75 (1.30)
V unskilled occupation	67		0.73 (0.23)	2.18 (2.63)	0.78 (1.14)	0.70 (1.13)	0.70 (1.24)
Total	2186	100	0.79 (0.21)	1.24 (2.11)	0.46 (1.00)	0.37(0.89)	0.41(0.96)

<https://doi.org/10.1371/journal.pone.0214380.t001>

1950s cohort n = 866; 1930s cohort n = 544). There were various reasons for non-collection of blood among those who had consented, for example: blood not running freely enough to permit collection (n = 53), the respondent being pregnant at the time of the interview (n = 11), respondent feeling unwell (n = 17), sample only partially collected (n = 79).

Table 1 provides descriptive statistics for the Wave 5 sample broken down by sex, age cohort and social class. Just over half of the sample (54.4%, n = 1192) were female, 35.4% were aged around 35, 39.7% aged around 55 and 25% aged around 75. The mean T/S for the sample was 0.79 (SD 0.21), males had shorter telomeres than females, telomere length was longest among the youngest cohort and shorter among lower social class groups. The mean (SD) for neighbourhood stressors as measured over 3 waves (the accumulation measure) was 1.24 (SD 0.21); for wave 3 this was 0.46 (SD 1.00); at wave 4 it was 0.37 (SD 0.89) and 0.441 at wave 5 (0.96). Females had higher scores for most measures of neighbourhood stressors (except wave 4), but this was not statistically significant. The oldest cohort had higher scores in the main compared to the younger cohorts, although this was only statistically significant at wave 4. For the accumulation score and for all 3 cross sectional waves, significantly higher scores for neighbourhood problems were observed among lower social class groups (p<0.001).

In models adjusted for age cohort, social class, depression, smoking status, diet, physical activity and BMI (Table 2), neighbourhood stressors were significantly associated with telomere in the accumulation model, with poorer neighbourhood perceptions resulting in shorter telomere length for women only (Beta = 0.007; 95%CI [0.001, 0.012]; P<0.014). Looking at critical periods, (i.e. for each wave separately), neighbourhood stressors reported at Wave 5 were associated with telomere length again among women only (Beta = 0.015; 95%CI [0.003, 0.027]; p<0.016).

Discussion

In this study, perceived neighbourhood stressors are associated with biological aging as measured by telomere length, even after taking known correlates such as chronological age, SES/

Table 2. Associations between perceived neighbourhood stressors and telomere length.

Perceived neighbourhood problems	<i>n</i>	β	CI	<i>P</i> <
Accumulation				
<i>Neighbourhood problems summed across 3 waves</i>				
<i>males</i>	951	0.002	-0.005, 0.008	0.636
<i>females</i>	1143	0.007	0.001, 0.012	0.014
Critical periods				
Wave 3				
<i>males</i>	951	0.002	-0.012, 0.015	0.826
<i>females</i>	1143	0.010	-0.001, 0.021	0.083
Wave 4				
<i>males</i>	951	0.008	-0.006, 0.023	0.249
<i>females</i>				
Wave 5				
<i>males</i>	951	-0.002	-0.016, 0.012	0.741
<i>females</i>	1143	0.015	0.003, 0.027	0.016

<https://doi.org/10.1371/journal.pone.0214380.t002>

social class, mental health (depression), health behaviours (smoking, diet, physical activity) and BMI into account. Our study is novel and adds to the literature, as although there is a growing body of research on the influence of characteristics of the neighbourhood environment and aging [10, 11, 26, 27], most studies to date which have explored perceptions of the local neighbourhood with telomere length have been cross sectional and few have investigated sex differences in these. Our models have explored perceptions of the neighbourhood measured over a 13-year time period and shown that the accumulation of neighbourhood stressors over time, and contemporaneous conditions, impact on telomere length. We found few associations between neighbourhood perceptions measured at earlier waves being associated with telomere length when examined cross-sectionally.

We found a stronger relationship between neighbourhood perceptions and telomere length for women compared to men. Other studies have not explored sex differences in the relationship between perceptions of neighbourhood and telomere length [11, 26] or have focussed on only one sex [27]. The gendered experience and consequences of place for health has previously been reported. For example, in studies comparing men and women, perceived neighbourhood crime is more strongly associated with the likelihood of women being a smoker [38], and neighbourhood trust was more strongly related to self-rated health among women [39]. The reasons for this may include differential exposure (women may spend more time than men in their local neighbourhood); women may be more sensitised to the quality of their neighbourhood, perhaps due to their being more likely to be the main carers of children; it is also possible that men and women process stressful environments differently in ways which might lead to different physiological responses. For example, it has previously been noted that there are sex differences in activation of the HPA axis as a response to stressful conditions [57].

Our findings are robust with respect to a measure of mental health (depression) known to be associated with telomere length [58]. In addition, by including depression in our models, we have partly accounted for the possibility that more negative perceptions of the neighbourhood are merely reflecting depressive symptoms. Moreover, depression may result from chronic stress which impacts on body systems (such as the immune system) which can affect telomere length [21, 22]. We also controlled for other health behaviours known to be

associated with telomere length such as smoking status, diet, physical activity and a measure of obesity (BMI).

Our study has been able to examine the associations between exposures of neighbourhood stressors over time in a large well characterized population study comprising a wide range of ages. Although not identical, our measures of perceived neighbourhood problems are broadly similar to other studies in the USA [26] and the Netherlands [11] which have explored the influence of these perceived problems on telomere length. A limitation of our study is that telomere length was only measured at one time point, the final wave. However, as we have previously noted, given that assays for this are a relatively recent development and the difficulties of collecting blood samples in large epidemiological studies, this is still one of the largest studies to examine telomere length across different age and SES groups [22]. The telomere length data used in our analyses were generated historically and contemporary with the original Cawthon method, prior to an updated methodology from his lab [59]. Our findings that perceived neighbourhood problems are associated with telomere length may be subject to residual confounding in that there are likely to be unmeasured socioeconomic circumstances that affect these associations, particularly those measured across the life course [60, 61]. Health selection may also play a part in that individuals in poor health may be 'sorted' into social housing (and associated poor neighbourhood conditions) due to the UK priority points system [60, 62]. In addition, we do not have data on how long respondents spend in their neighbourhoods on a day-to-day basis and therefore the extent of their exposure to neighbourhood conditions. Finally, although we have found that chronic environmental stressors, such as neighbourhood problems accumulated over time, are related to shortened telomere length, we do not have a measure of physiological stress in our study and hence are unable to examine direct biological pathways between neighbourhood conditions and telomere length.

In conclusion, we have shown that chronic neighbourhood stressors are associated with shorter telomere length, particularly among women. Neighbourhood environments are potentially modifiable, and future efforts directed towards improving deleterious local environments may be useful to lessen telomere attrition.

Acknowledgments

We are grateful to participants and staff of all three studies.

Author Contributions

Conceptualization: Anne Ellaway, Paul G. Shiels.

Formal analysis: Anne Ellaway, Ruth Dundas, Paul G. Shiels.

Methodology: Anne Ellaway, Ruth Dundas, Paul G. Shiels.

Writing – original draft: Anne Ellaway.

Writing – review & editing: Ruth Dundas, Tony Robertson, Paul G. Shiels.

References

1. Ellaway A, Macintyre S. Are perceived neighbourhood problems associated with the likelihood of smoking? *Journal of Epidemiology and Community Health*. 2009; 63(1):78–80. <https://doi.org/10.1136/jech.2007.068767> WOS:000261761100014. PMID: 19088119
2. Ellaway A, Morris G, Curtice J, Robertson C, Allardice G, Robertson R. Associations between health and different types of environmental incivility: a Scotland-wide study. *Public Health*. 2009; 123(11):708–13. Epub 2009/11/04. S0033-3506(09)00276-5 [pii] <https://doi.org/10.1016/j.puhe.2009.09.019> PMID: 19883927.

3. Jones R, Heim D, Hunter S, Ellaway A. The relative influence of neighbourhood incivilities, cognitive social capital, club membership and individual characteristics on positive mental health. *Health Place*. 2014; 28:187–93. Epub 2014/06/10. S1353-8292(14)00064-1 [pii] <https://doi.org/10.1016/j.healthplace.2014.04.006> PMID: 24911971.
4. Kim D. Blues from the neighborhood? Neighborhood characteristics and depression. *Epidemiol Rev*. 2008; 30:101–17. Epub 2008/08/30. mxn009 [pii] <https://doi.org/10.1093/epirev/mxn009> PMID: 18753674.
5. Sooman A, Macintyre S. Health and perceptions of the local environment in socially contrasting neighbourhoods in Glasgow. *Health & Place*. 1995; 1(1):15–26.
6. Stansfeld S, Smith G, Marmot M. Association between physical and psychological morbidity in the Whitehall II Study. *Journal of Psychosomatic Research*. 1993;37.
7. Diez Roux A, Mair C. Neighborhoods and health. *Ann N Y Acad Sci*. 2010; 1186:125–45. <https://doi.org/10.1111/j.1749-6632.2009.05333.x> PMID: 20201871
8. Ellaway A, Benzeval M, Green M, Leyland A, Macintyre S. "Getting sicker quicker": Does living in a more deprived neighbourhood mean your health deteriorates faster? *Health & Place*. 2012; 18(2):132–7. <https://doi.org/10.1016/j.healthplace.2011.08.005> WOS:000301276800002. PMID: 21873103
9. Evans G, Chen E, Miller G, Seeman T. How Poverty Gets Under the Skin: A Life Course Perspective In: King R, Maholmes V, editors. *The Oxford Handbook of Poverty and Child Development* 2012.
10. Lynch SM, Mitra N, Ravichandran K, Mitchell J, Spangler E, Zhou W, et al. Telomere Length and Neighborhood Circumstances: Evaluating Biological Response to Unfavorable Exposures. *Cancer Epidemiol Biomarkers Prev*. 2017; 26(4):553–60. Epub 2017/04/05. 26/4/553 [pii] <https://doi.org/10.1158/1055-9965.EPI-16-0554> PMID: 28373169.
11. Park M, Verhoeven JE, Cuijpers P, Reynolds CF III, Penninx BW. Where You Live May Make You Old: The Association between Perceived Poor Neighborhood Quality and Leukocyte Telomere Length. *PLoS One*. 2015; 10(6):e0128460. Epub 2015/06/18. <https://doi.org/10.1371/journal.pone.0128460> PONE-D-14-55855 [pii]. PMID: 26083263; PubMed Central PMCID: PMC4471265.
12. McGuinness D, McGlynn LM, Johnson PC, MacIntyre A, Batty GD, Burns H, et al. Socio-economic status is associated with epigenetic differences in the pSoBid cohort. 2012; 41(1):151–60. Epub 2012/01/19. dyr215 [pii] <https://doi.org/10.1093/ije/dyr215> PMID: 22253320.
13. Shiels PG, McGlynn LM, MacIntyre A, Johnson PC, Batty GD, Burns H, et al. Accelerated telomere attrition is associated with relative household income, diet and inflammation in the pSoBid cohort. *PLoS One*. 2011; 6(7):e22521. Epub 2011/08/06. <https://doi.org/10.1371/journal.pone.0022521> PONE-D-11-07269 [pii]. PMID: 21818333; PubMed Central PMCID: PMC3144896.
14. Shiels PG, Davies RW. Ageing and death in neurones. In: Davies R, Morris B, editors. *The Molecular Biology of the Neurone*. Oxford: Oxford University Press; 2003. p. 435–64.
15. Olovnikov AM. A theory of marginotomy. The incomplete copying of template margin in enzymic synthesis of polynucleotides and biological significance of the phenomenon. *J Theor Biol*. 1973; 41(1):181–90. Epub 1973/09/14. 0022-5193(73)90198-7 [pii]. PMID: 4754905.
16. Watson J. Origin of concatemeric T7 DNA. *Nat New Biol*. 1972;239.
17. Der G, Batty GD, Benzeval M, Deary IJ, Green MJ, McGlynn L, et al. Is telomere length a biomarker for aging: cross-sectional evidence from the west of Scotland? *PLoS One*. 2012; 7(9):e45166. Epub 2012/10/03. <https://doi.org/10.1371/journal.pone.0045166> PONE-D-12-13258 [pii]. PMID: 23028820; PubMed Central PMCID: PMC3454419.
18. Gardner M, Bann D, Wiley L, Cooper R, Hardy R, Nitsch D, et al. Gender and telomere length: systematic review and meta-analysis. *Exp Gerontol*. 2014; 51:15–27. Epub 2013/12/25. S0531-5565(13)00360-4 [pii] <https://doi.org/10.1016/j.exger.2013.12.004> PMID: 24365661; PubMed Central PMCID: PMC4523138.
19. Cherkas LF, Aviv A, Valdes AM, Hunkin JL, Gardner JP, Surdulescu GL, et al. The effects of social status on biological aging as measured by white-blood-cell telomere length. *Aging Cell*. 2006; 5(5):361–5. Epub 2006/07/22. ACE222 [pii] <https://doi.org/10.1111/j.1474-9726.2006.00222.x> PMID: 16856882.
20. Robertson T, Batty GD, Der G, Fenton C, Shiels PG, Benzeval M. Is socioeconomic status associated with biological aging as measured by telomere length? *Epidemiol Rev*. 2013; 35:98–111. Epub 2012/12/22. mxs001 [pii] <https://doi.org/10.1093/epirev/mxs001> PMID: 23258416; PubMed Central PMCID: PMC3578449.
21. Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, et al. Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci U S A*. 2004; 101(49):17312–5. Epub 2004/12/03. 0407162101 [pii] <https://doi.org/10.1073/pnas.0407162101> PMID: 15574496; PubMed Central PMCID: PMC534658.

22. Phillips AC, Robertson T, Carroll D, Der G, Shiels PG, McGlynn L, et al. Do symptoms of depression predict telomere length? Evidence from the West of Scotland Twenty-07 study. *Psychosom Med*. 2013; 75(3):288–96. Epub 2013/03/21. PSY.0b013e318289e6b5 [pii] <https://doi.org/10.1097/PSY.0b013e318289e6b5> PMID: 23513237.
23. Ornish D, Lin J, Chan JM, Epel E, Kemp C, Weidner G, et al. Effect of comprehensive lifestyle changes on telomerase activity and telomere length in men with biopsy-proven low-risk prostate cancer: 5-year follow-up of a descriptive pilot study. *Lancet Oncol*. 2013; 14(11):1112–20. [https://doi.org/10.1016/S1470-2045\(13\)70366-8](https://doi.org/10.1016/S1470-2045(13)70366-8) PMID: 24051140.
24. Kooman JP, Kotanko P, Schols AM, Shiels PG, Stenvinkel P. Chronic kidney disease and premature ageing. *Nature Reviews Nephrology*. 2014; 10(12):732–42. <https://doi.org/10.1038/nrneph.2014.185> PMID: 25287433
25. Crous-Bou M, Fung TT, Prescott J, Julin B, Du M, Sun Q, et al. Mediterranean diet and telomere length in Nurses' Health Study: population based cohort study. *Bmj-Brit Med J*. 2014; 349. ARTN g6674 <https://doi.org/10.1136/bmj.g6674> WOS:000346116300002. PMID: 25467028
26. Needham BL, Carroll JE, Diez Roux AV, Fitzpatrick AL, Moore K, Seeman TE. Neighborhood characteristics and leukocyte telomere length: the Multi-Ethnic Study of Atherosclerosis. *Health Place*. 2014; 28:167–72. Epub 2014/05/27. S1353-8292(14)00067-7 [pii] <https://doi.org/10.1016/j.healthplace.2014.04.009> PMID: 24859373; PubMed Central PMCID: PMC4096814.
27. Gebreab SY, Riestra P, Gaye A, Khan RJ, Xu R, Davis AR, et al. Perceived neighborhood problems are associated with shorter telomere length in African American women. *Psychoneuroendocrinology*. 2016; 69:90–7. Epub 2016/04/14. S0306-4530(16)30079-8 [pii] <https://doi.org/10.1016/j.psyneuen.2016.03.018> PMID: 27070760; PubMed Central PMCID: PMC5051547.
28. Robertson T, Batty GD, Der G, Green MJ, McGlynn LM, McIntyre A, et al. Is telomere length socially patterned? Evidence from the West of Scotland Twenty-07 Study. *PLoS One*. 2012; 7(7):e41805. <https://doi.org/10.1371/journal.pone.0041805> PMID: 22844525; PubMed Central PMCID: PMC3402400.
29. Macintyre S, Ellaway A, Hiscock R, Kearns A, Der G, McKay L. What features of the home and the area might help to explain observed relationships between housing tenure and health? Evidence from the west of Scotland. *Health Place*. 2003; 9(3):207–18. Epub 2003/06/18. S1353829202000400 [pii]. PMID: 12810328.
30. Clarke P, Morenoff J, Debbink M, Golberstein E, Elliott MR, Lantz PM. Cumulative exposure to neighborhood context: consequences for health transitions over the adult life course. *Res Aging*. 2014; 36(1):115–42. <https://doi.org/10.1177/0164027512470702> PMID: 24465068; PubMed Central PMCID: PMC3900407.
31. Aneshensel C. Neighborhood as a social context of the stress process. In: Avison W, Aneshensel C, Schieman S, Wheaton B, editors. *Advances in the conceptualization of the stress process Essays in honor of Leonard L Pearlman*. New York: Springer; 2010. p. 35–52.
32. Lopez-Isaac E, Bossini-Castillo L, Simeon CP, Egurbide MV, Alegre-Sancho JJ, Callejas JL, et al. A genome-wide association study follow-up suggests a possible role for PPARγ in systemic sclerosis susceptibility. *Arthritis Res Ther*. 2014; 16(1):R6. <https://doi.org/10.1186/ar4432> PMID: 24401602; PubMed Central PMCID: PMC3978735.
33. Shiels PG, McGuinness D, Eriksson M, Kooman JP, Stenvinkel P. The role of epigenetics in renal ageing. *Nat Rev Nephrol*. 2017; 13(8):471–82. <https://doi.org/10.1038/nrneph.2017.78> PMID: 28626222.
34. Shiels PG, Stenvinkel P, Kooman JP, McGuinness D. Circulating markers of ageing and allostatic load: A slow train coming. *Pract Lab Med*. 2017; 7:49–54. <https://doi.org/10.1016/j.plabm.2016.04.002> PMID: 28856219; PubMed Central PMCID: PMC39574864.
35. Mishra G, Nitsch D, Black S, De Stavola B, Kuh D, Hardy R. A structured approach to modelling the effects of binary exposure variables over the life course. *Int J Epidemiol*. 2009; 38(2):528–37. Epub 2008/11/26. dyn229 [pii] <https://doi.org/10.1093/ije/dyn229> PMID: 19028777; PubMed Central PMCID: PMC2663717.
36. Ellaway A, Emslie C. Connecting gender, space and place: are there gender differences in the relationships between the social environment and health-related behaviours?. *Neighbourhood Structure and Health Promotion* Springer; 2013.
37. Ellaway A, Macintyre S. Women in their place. Gender perceptions of neighbourhoods and health in the West of Scotland. In: Dyack I, vis Lewis N. and McLafferty S. (eds.) editor. *Geographies of Women's Health*. London: Routledge; 2001.
38. Shareck M, Ellaway A. Neighbourhood crime and smoking: the role of objective and perceived crime measures. *Bmc Public Health*. 2011; 11. <https://doi.org/10.1186/1471-2458-11-930> WOS:000300286200003. PMID: 22168151

39. Stafford M, Cummins S, Macintyre S, Ellaway A, Marmot M. Gender differences in the associations between health and neighbourhood environment. *Soc Sci Med*. 2005; 60(8):1681–92. Epub 2005/02/03. S0277-9536(04)00418-6 [pii] <https://doi.org/10.1016/j.socscimed.2004.08.028> PMID: 15686801.
40. Der G. A comparison of the West of Scotland Twenty-07 study sample and the 1991 census SARS. Glasgow: MRC Social and Public Health Sciences Unit, 1998.
41. Benzeval M, Der G, Ellaway A, Hunt K, Sweeting H, West P, et al. Cohort profile: west of Scotland twenty-07 study: health in the community. *Int J Epidemiol*. 2009; 38(5):1215–23. Epub 2008/10/22. dyn213 [pii] <https://doi.org/10.1093/ije/dyn213> PMID: 18930962; PubMed Central PMCID: PMC2935558.
42. Connelly R, Gayle V, Lambert P. A Review of occupation-based social classifications for social survey research. *Methodological Innovations*. 2016; 9:1–14.
43. Office of Population Censuses and Surveys. OPCS standard occupational classification. London: HMSO, 1991.
44. Zigmond A, Snaith R. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand*. 1983; 67:361–70. PMID: 6880820
45. Goldberg D, Williams P. A user's guide to the General Health Questionnaire. Basingstoke NFER-Nelson; 1988.
46. Nettleton JA, Diez-Roux A, Jenny NS, Fitzpatrick AL, Jacobs DR Jr. Dietary patterns, food groups, and telomere length in the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr*. 2008; 88(5):1405–12. Epub 2008/11/11. 88/5/1405 [pii]. <https://doi.org/10.3945/ajcn.2008.26429> PMID: 18996878; PubMed Central PMCID: PMC3037593.
47. Astuti Y, Wardhana A, Watkins J, Wulaningsih W. Cigarette smoking and telomere length: A systematic review of 84 studies and meta-analysis. *Environ Res*. 2017; 158:480–9. Epub 2017/07/14. S0013-9351(17)30433-4 [pii] <https://doi.org/10.1016/j.envres.2017.06.038> PMID: 28704792; PubMed Central PMCID: PMC5562268.
48. Arsenis NC, You T, Ogawa EF, Tinsley GM, Zuo L. Physical activity and telomere length: Impact of aging and potential mechanisms of action. *Oncotarget*. 2017; 8(27):45008–19. Epub 2017/04/15. 16726 [pii] <https://doi.org/10.18632/oncotarget.16726> PMID: 28410238; PubMed Central PMCID: PMC5546536.
49. Wulaningsih W, Kuh D, Wong A, Hardy R. Adiposity, Telomere Length, and Telomere Attrition in Midlife: the 1946 British Birth Cohort. *J Gerontol A Biol Sci Med Sci*. 2018; 73(7):966–72. <https://doi.org/10.1093/gerona/glx151> PMID: 28957990; PubMed Central PMCID: PMC6001881.
50. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res*. 2002; 30(10). WOS:000175591600024.
51. Koppelstaetter C, Schratzberger G, Perco P, Hofer J, Mark W, Ollinger R, et al. Markers of cellular senescence in zero hour biopsies predict outcome in renal transplantation. *Aging Cell*. 2008; 7(4):491–7. <https://doi.org/10.1111/j.1474-9726.2008.00398.x> PMID: 18462273.
52. Carrero JJ, Stenvinkel P, Fellstrom B, Qureshi AR, Lamb K, Heimbürger O, et al. Telomere attrition is associated with inflammation, low fetuin-A levels and high mortality in prevalent haemodialysis patients. *J Intern Med*. 2008; 263(3):302–12. Epub 2007/12/12. JIM1890 [pii] <https://doi.org/10.1111/j.1365-2796.2007.01890.x> PMID: 18070000.
53. Gale CR, Dennison EM, Cooper C, Sayer AA. Neighbourhood environment and positive mental health in older people: the Hertfordshire Cohort Study. *Health Place*. 2011; 17(4):867–74. <https://doi.org/10.1016/j.healthplace.2011.05.003> PMID: 21605999.
54. Timmermans EJ, van der Pas S, Cooper C, Schaap LA, Edwards MH, Deeg DJ, et al. The neighbourhood environment and use of neighbourhood resources in older adults with and without lower limb osteoarthritis: results from the Hertfordshire Cohort Study. *Clin Rheumatol*. 2016; 35(11):2797–805. <https://doi.org/10.1007/s10067-016-3388-5> PMID: 27567628; PubMed Central PMCID: PMC5063902.
55. Ellaway A, Dundas R, Olsen JR, Shiels PG. Perceived Neighbourhood Problems over Time and Associations with Adiposity. *Int J Environ Res Public Health*. 2018; 15(9). <https://doi.org/10.3390/ijerph15091854> PMID: 30154306; PubMed Central PMCID: PMC6164418.
56. Olsen JR, Dundas R, Ellaway A. Are Changes in Neighbourhood Perceptions Associated with Changes in Self-Rated Mental Health in Adults? A 13-Year Repeat Cross-Sectional Study, UK. *Int J Environ Res Public Health*. 2017; 14(12). <https://doi.org/10.3390/ijerph14121473> PMID: 29186035; PubMed Central PMCID: PMC5750892.
57. Pasquali R. The hypothalamic-pituitary-adrenal axis and sex hormones in chronic stress and obesity: pathophysiological and clinical aspects. *Ann N Y Acad Sci*. 2012; 1264:20–35. <https://doi.org/10.1111/j.1749-6632.2012.06569.x> PMID: 22612409; PubMed Central PMCID: PMC3464358.

58. Ridout KK, Ridout SJ, Price LH, Sen S, Tyrka AR. Depression and telomere length: A meta-analysis. *J Affect Disord.* 2016; 191:237–47. Epub 2015/12/22. S0165-0327(15)31078-8 [pii] <https://doi.org/10.1016/j.jad.2015.11.052> PMID: 26688493; PubMed Central PMCID: PMC4760624.
59. Cawthon R. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res.* 2009; 37(3).
60. Ellaway A, Macdonald L, Kearns A. Are housing tenure and car access still associated with health? A repeat cross-sectional study of UK adults over a 13-year period. *BMJ Open.* 2016; 6(11):e012268. <https://doi.org/10.1136/bmjopen-2016-012268> PMID: 27807086; PubMed Central PMCID: PMCPMC5128997.
61. Galobardes B, Shaw M, Lawlor DA, Lynch JW, Davey Smith G. Indicators of socioeconomic position (part 1). *J Epidemiol Community Health.* 2006; 60(1):7–12. Epub 2005/12/20. 60/1/7 [pii] <https://doi.org/10.1136/jech.2004.023531> PMID: 16361448; PubMed Central PMCID: PMC2465546.
62. Smith S. Explaining the Link between Housing and Health: All or Nothing? *Housing, Theory and Society.* 2012; 29(1):40–6.