

Accepted Manuscript

Survival and immune response of white shrimp *Litopenaeus vannamei* following single and concurrent infections with WSSV and *Vibrio parahaemolyticus*

Huanying Pang, Gang Wang, Shihui Zhou, Junlin Wang, Jichen Zhao, Rowena Hoare, Sean J. Monaghan, Ziling Wang, Chengbo Sun



PII: S1050-4648(19)30686-2

DOI: <https://doi.org/10.1016/j.fsi.2019.06.039>

Reference: YFSIM 6235

To appear in: *Fish and Shellfish Immunology*

Received Date: 22 February 2019

Revised Date: 12 June 2019

Accepted Date: 19 June 2019

Accepted refereed manuscript of: : Pang H, Wang G, Zhou S, Wang J, Zhao J, Hoare R, Monaghan SJ, Wang Z, Sun C (2019) Survival and immune response of white shrimp *Litopenaeus vannamei* following single and concurrent infections with WSSV and *Vibrio parahaemolyticus*, *Fish and Shellfish Immunology*, 92, pp. 712-718, doi: <https://doi.org/10.1016/j.fsi.2019.06.039>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

**Survival and immune response of white shrimp *Litopenaeus*
vannamei following single and concurrent infections with WSSV
and *Vibrio parahaemolyticus***

Huanying Pang^{abef1}, Gang Wang^{a1}, Shihui Zhou^{ab1}, Junlin Wang^a, Jichen Zhao^a,

Rowena Hoare^d, Sean J. Monaghan^d, Ziling Wang^a, Chengbo Sun^{ac□*}

^aFisheries College, Guangdong Ocean University, Zhanjiang 524025, China;

^bGuangdong Provincial Key Laboratory of Pathogenic Biology and Epidemiology for
Aquatic Economic Animals, Zhanjiang 524025, China; Guangdong Key Laboratory
of Control for Diseases of Aquatic Economic Animals, Zhanjiang 524025, China

^cTropical Invertebrates Aquaculture Research Center of Guangdong Colleges and
Universities, Zhanjiang 524025, China.

^dInstitute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK

^eKey Laboratory of Experimental Marine Biology, Institute of Oceanology, Chinese
Academy of Sciences, Qingdao 266071, China; Laboratory for Marine Biology and
Biotechnology, Qingdao National Laboratory for Marine Science and Technology,
Qingdao 266071, China

^fGuangdong provincial engineering research center for aquatic animal health
assessment, Shenzhen 518120, China

* Corresponding author: Chengbo Sun, E-mail: scb248@126.com

Address: College of Fishery, Guangdong Ocean University, Zhanjiang 524025, China

1 These authors contributed to the work equally and should be regarded as co-first
authors.

ABSTRACT

The survival and immune responses of *Litopenaeus vannamei* were evaluated during white spot syndrome virus (WSSV) or *Vibrio parahaemolyticus* single and concurrent infections. The mortality, WSSV load, activities of 4 immune enzymes: acid phosphatase (ACP), alkaline phosphatase (AKP), peroxidase (POD) and superoxide dismutase (SOD), and the transcription of Evolutionarily Conserved Signaling Intermediate in Toll pathways of *L.vannamei* (LvECSIT) were quantified at 0, 3, 6, 12, 24, 48, 72 and 96 h post-infection (pi). The results showed: (i) the cumulative mortality of the co-infection group (WSSV and *V. Parahaemolyticus* 83 %) was significantly lower than the WSSV infection group (97%) ($P < 0.05$) at 96 hpi; (ii) copies of WSSV in the co-infection group were significantly lower than that of the single infection group from 24 to 96 hpi ($P < 0.05$); (iii) ACP, AKP, POD and SOD activity in the gills of the co-infection group was higher than that of the WSSV group at 12, 48 and 96 hpi ($P < 0.05$). The expression of LvECSIT mRNA in the co-infection group was significantly higher than in the WSSV infection group from 12 to 72 hpi ($P < 0.05$). The results indicate that proliferation of WSSV is inhibited by *V.parahaemolyticus* infection. In addition, infection with WSSV alone causes a significant reduction in some immune responses of shrimp than co-infection with WSSV and *V.parahaemolyticus* occurs at 26 °C. Third, LvECSIT, an essential member of TLR signaling pathway might play a crucial role in shrimp defense against WSSV – *Vibrio* co- infection.

Keywords: *Litopenaeus vannamei*, Immune response, White spot syndrome virus (WSSV), *Vibrio parahaemolyticus*, Co-infection

1. Introduction

Shrimp aquaculture has developed very fast in China over the last two decades, but the production of shrimp has been seriously affected by white spot syndrome virus (WSSV) and *Vibrio* spp. [1]. WSSV - *Vibrio* co-infection is the normal manner of shrimp disease breakouts and shrimp infected with the virus are more susceptible to *Vibrio* spp.[2]. It has been reported previously that *Vibrio alginolyticus* was isolated from shrimp during a breakout of white spot syndrome virus [3]. Another study showed that during a WSSV and *Vibrio anguillarum* co-infection test in shrimp, WSSV increased more rapidly under co-infection conditions than in the single infection[4]. Similarly, the transcription of immune-related genes was suppressed in the co-infection groups, and the shrimp would suffer higher mortality in multiple infections [5]. Unlike the above observations, an outbreak of WSSV was postponed after co-infection with WSSV and *Vibrio harveyi* in *Penaeus vannamei* [6]. These studies about the WSSV - *Vibrio* co-infections in shrimp seem to be conflicting and the pathogenesis involved is unclear.

Although the defense mechanism of shrimp to WSSV - *Vibrio* co-infections remains unknown, it has been reported that bacterial infection could reduce the copies of virus in some arthropods [7-8]. *Drosophila melanogaster* infected with *Wolbachia* appeared to inhibit the proliferation of Drosophila C virus[7]. Furthermore, *Wolbachia* induces reactive oxygen species (ROS)-dependent activation of the Toll/Toll-like receptor (TLR)-mediated signaling pathway to control dengue virus in the mosquito *Aedes aegypti*. Some Toll pathway-related genes (Spn27A, SPZ1, CECD, and DEFC)

were up-regulated in *Aedes aegypti* after co-infection with *Wolbachia* and dengue virus [8]. Such virus suppression mechanisms may exist in shrimp, which warrants further exploration.

In shrimp, the innate immune system is the first line of defense against pathogenic infections [9]. When pathogens invade shrimp, they stimulate a series of immune responses including lymphatic hemocyte agglutination, melanisation, hemocyte phagocytosis, formation of cysts [10-12] and humoral immune factors (a variety of enzymes have been identified). It was reported that ACP, AKP, POD and SOD were susceptible to WSSV and *Vibrio* infections, and they could be used as indicators of immune response to these pathogens [13-15].

Under the stimulus of pathogens, various humoral and cellular immune responses of shrimp are activated through signaling pathways, among which Toll/Toll-like receptor (TLR)-mediated signaling pathway are the best known and can be activated by pathogenic related molecular patterns (PAMPs) [16]. After PAMP recognition, TLRs can either directly or indirectly trigger downstream signaling cascades, resulting in the regulation of cytokine gene expression [17]. TRAF6 is an important downstream signal ligand of Toll-1 receptor protein and ECSIT is the first gene that has been approved to interact with TRAF 6 [18]. As an important adaptor protein of TLR, ECSIT have been demonstrated to be an immune-response gene since its transcript expression level is up-regulated after *Vibrio anguillarum* [19] or WSSV infection [20].

White spot syndrome virus (WSSV) is one of the most detrimental pathogens

affecting shrimp [21]. It is a baculovirus with double stranded DNA [22], and the mortality rate of WSSV-infected shrimp can reach 100% in 7-10 days. Recently, researchers found another serious shrimp disease (acute hepatopancreatic necrosis disease AHPNS/early mortality syndrome EMS), which is characterised by empty stomach, severe atrophy of hepatopancreas and soft carapace. *Vibrio parahaemolyticus* is one of the causative agents of AHPNS / EMS, and it has caused big economic losses in the shrimp industry in China [23-25]. Nowadays, there is limited information available on molecular immune responses in shrimp under WSSV or *V. parahaemolyticus* single and concurrent infections.

In an attempt to provide a theoretical basis for the control of WSSV in *L. vannamei*, a number of parameters (mortality, WSSV load, the activities of the several immune enzymes, transcription of LvECSIT) were investigated following single infections and co-infection with WSSV and *V. parahaemolyticus*.

2. Materials and methods

2.1. Experimental animals and conditions

L. vannamei (size 7.66 ± 0.82 cm) were obtained from the East Sea Island Marine Biological Research Center in Guangdong Ocean University. Before the experiment, 20 shrimp were randomly selected to ensure that they were free of WSSV and *V. parahaemolyticus*, according to Sun *et al.* [14]. They were fed with artificial pellet diets twice a day and were kept at 26°C and salinity at 25 ‰. Filtered seawater was sterilized with 1.5 ppm trichloroisocyanuric acid and the residual chlorine was

detected to ensure that it was safe for shrimp. About 1/3 of the water was replaced and un-eaten pellet diet was removed by siphoning daily.

2.2 Preparation of virus and *V. parahaemolyticus* suspension for injection

WSSV extracts were prepared from crude extracts of disease shrimp and stored at - 80 °C. Healthy shrimp were injected intramuscularly with 3.3×10^2 copies μL^{-1} virus (in PBS) and mortalities occurred at 48 h post-injection (pi). Following removal of the exoskeletons, WSSV infected shrimp were homogenized in cold PBS (KH_2PO_4 0.27g, Na_2HPO_4 0.01g, NaCl 8g, KCl 0.2g, diluted with water to 1 L and adjust pH to 7.4) (1 mL g^{-1}). After centrifugation at 12,000 g for 10 min, the crude viral supernatant was filtered using a membrane filter (220 nm).

V. parahaemolyticus was obtained from the Economic Aquatic Animal Disease Control Laboratory of the Guangdong Ocean University [26]. *V. parahaemolyticus* was cultured in trypticase soy broth (TSB, Huankai Co Ltd., Guangzhou, China) at 28 °C for 18 h. The culture medium was centrifuged in an 8 mL tube at 4000 g for 15 min. The supernatant was removed and *V. parahaemolyticus* was re-suspended in PBS to 1.22×10^6 CFU mL^{-1} .

2.3 Experimental design

The laboratory challenge test contained 4 treatments in triplicate (n=40 for each sample group, n=10 for mortality group). For *V. parahaemolyticus* treatment, shrimp were intramuscularly injected with 50 μL of *V. parahaemolyticus* (1.22×10^6 CFU

mL⁻¹). For WSSV treatment, shrimp were intramuscularly injected with 50 µL of WSSV viral suspension (3.3×10^2 copies µL⁻¹). For co-infection treatment, shrimp were intramuscularly injected with 50 µL of cocktail suspensions containing *V. prahaemolyticus* (1.22×10^6 CFU mL⁻¹) and WSSV (3.3×10^2 copies µL⁻¹). The PBS treatment was injected with 50 µL of PBS. Tissues (muscle, gills) of one shrimp per group were sampled individually at PBS 0 h post-infection (pi), and at each time point (3, 6, 12, 24, 48, 72 and 96 hpi) from each group to measure virus load, immune-related enzymes, and immune-related gene LvECSIT expression analysis (Table 1-2). The experiments were repeated three times.

2.4 Analysis of virus load

The muscle of the first abdominal segment (about 0.05 g) was dissected and added to 45 µL 50 mM NaOH and homogenized on ice, mixed and then boiled in water bath for 10 min. Then, 5 µL 1M Tris solution was added, mixed and centrifuged at 12,000 g for 10 min [14]. The supernatant was used as WSSV template for quantitative PCR. The qPCR was carried out in 15 µL volume, and the primer sequences are shown in Table 3. The standard curve was made according to the method of Xin *et al.* [27].

2.5 Determination of activities of immune-related enzymes in the gills

The gills (0.2g) were cut off from the samples stored in liquid nitrogen and homogenized on ice after adding 1.8 mL PBS. The samples were centrifuged at 3000g

for 10 min at 4 °C, the precipitate was removed and the supernatant was used for acid phosphatase (ACP), alkaline phosphatase (AKP), peroxidase (POD) and superoxide dismutase (SOD) immune enzyme analysis. Enzymatic activities for ACP, AKP, POD, SOD were determined using kits purchased from Jiancheng Bioengineering Institute (NJCBio, Nanjing, China), according to the methods described by Sun *et al.* and Liu *et al.* [14,28]. ACP and AKP activities are expressed in King unit (mg protein)⁻¹. POD and SOD activities are expressed in U (mg protein)⁻¹. Each enzymatic assay was performed in triplicate.

2.6 Immune-related gene LvECSIT expression analysis by real-time PCR

Gills from one shrimps were sampled [20] at PBS 0 h post-infection (pi) and at each time point (3, 6, 12, 24, 48, 72 and 96 hpi) from each group. The transcriptional level of LvECSIT was detected with real-time PCR. Primers for LvECSIT (Genbank accession No. is XM_027378031) were shown in Table 3. β -actin was used as internal reference. RNA extraction, cDNA synthesis, real-time PCR for analysis of immune gene expression were as described by Li et al. [29].

2.7 Statistical analysis

Statistical analysis was carried out using the software SPSS 21. Results were analyzed using One-way ANOVA and Duncan's multiple comparisons of the means. Differences were considered significant when $P < 0.05$.

3. Results

3.1 Effect of WSSV and *V. parahaemolyticus* infection on shrimp survival

Shrimp in each challenge group started to die at 12 hpi. The cumulative mortality reached peak at 96 hpi, and the mortality of WSSV group (97 %) was significantly higher than co-infection group (83 %) and *V. parahaemolyticus* group (34 %) ($P < 0.05$) (Fig.1).

3.2 Effects of WSSV and *V. parahaemolyticus* infection on the proliferation of WSSV in *L. vannamei*

In the experiment, we collected the muscle of shrimp to detect the copies of WSSV by real time PCR. The results illustrated that WSSV could be detected in muscle within 3 h, and the maximum viral load in the WSSV infection group was 6.71×10^5 copies μL^{-1} at 72 hpi, significantly higher than that in co-infection group (1.80×10^4 copies μL^{-1}). The viral load in the WSSV infection group was approximately 10 times more than that in co-infection group at 24, 48, 72 and 96 hpi (Fig.2).

3.3 Effects of WSSV and *V. parahaemolyticus* infection on shrimp gill immune enzyme activity

The ACP activity in the gills of shrimp infected with *V. parahaemolyticus* alone and the co-infection groups showed an initial rise and subsequent fall, and reached maximum activity at 24 and 6 hpi respectively. In the *V. parahaemolyticus* group and

co-infection group, the maximum ACP activity was significantly higher than the PBS group and WSSV group at 6, 12, 24, 48, 72 and 96 hpi ($P < 0.05$). By the end of the experiment, the ACP activity of WSSV group remained at a low level, and was consistently lower than both the *V. parahaemolyticus* and the co-infection groups. Comparison of the degree of variation of each treatment group showed the following trend: PBS group (0.14) < WSSV group (0.33) < *V. parahaemolyticus* group (0.45) < co-infection group (0.58) (Fig.3A).

In the *V. parahaemolyticus* group and co-infection group, the AKP activity decreased after the initial rise, and was higher than the WSSV group and PBS group at all time points, and the maximum AKP activity was recorded at 6 h and 24 hpi respectively. The AKP activity of WSSV group was significantly lower than the co-infection group from 6-96 hpi. The AKP activity of *V. parahaemolyticus* group varied over the course of the experiment whereas the AKP activity of the PBS group was stable. Degree of variation: PBS group (0.18) < WSSV group (0.21) < co-infection group (0.29) < *V. parahaemolyticus* group (0.45) (Fig.3B).

The POD activity of the PBS group remained higher than 3 challenge groups until the end of experiment, and the difference was significant at 48 hpi ($P < 0.05$). For the *V. parahaemolyticus* group, co-infection group and WSSV group, the minimum POD activity occurred at 3, 6 and 24 hpi respectively. The POD activity of the co-infection group was higher than the WSSV group at 6, 12, 48 and 96 hpi, and was significantly higher at 6 hpi. Degree of variation: PBS group (0.05) < *V. parahaemolyticus* group (0.11) < co-infection group (0.14) < WSSV group (0.15)

(Fig.3C).

SOD activity of the WSSV and co-infection groups showed the lowest value at 96 h pi, which was significantly lower than PBS group ($P < 0.05$). The SOD activity of the co-infection group was significantly higher than the WSSV group at 48 hpi ($P < 0.05$). The SOD activity of *V. parahaemolyticus* group was significantly higher than WSSV group at 3, 6, 48 and 96 hpi ($P < 0.05$). SOD activity in each group variation coefficient: PBS group (0.11) $<$ *V. parahaemolyticus* group (0.18) $<$ co-infection group (0.24) $<$ WSSV group (0.32) (Fig.3D).

3.4 Effects of WSSV, *V. parahaemolyticus*, and WSSV and *V. parahaemolyticus* co-infection on LvECSIT expression in shrimp

In the challenge test, the expression of LvECSIT was detected in gill at 0, 3, 6, 12, 24, 48, 72 and 96 hpi. The transcription levels of LvECSIT in the PBS group up-regulated from 6 to 48 hpi. WSSV infection group showed a degree of fluctuation and reached maximum expression at 48h. Furthermore, LvECSIT expression up-regulated significantly in WSSV infection group more than co-infection group at 3hpi, and was significantly more up-regulated than *V. parahaemolyticus* group at 6 hpi. The LvECSIT expression was significantly up-regulated in *V. parahaemolyticus* group or co-infection group when compared with the WSSV infection group from 12 to 72 hpi ($P < 0.01$). There was no significant difference between the *V. parahaemolyticus* group and co-infection group from 12 to 48 hpi. Each treatment group showed minimum LvECSIT expression at 96 hpi and was all significantly

lower than PBS group ($P < 0.05$) (Fig. 4).

4. Discussion

In complex aquaculture environments, the outbreak of shrimp disease is accompanied with sharply defined changes of physical factors or secondary infection and co-infection by pathogens [30-32]. Nonetheless, the conclusions about *Vibrio* spp. and WSSV co-infection in shrimp have been conflicting. Previous studies have shown that mortality in co-infections (39%) was significantly higher than in single WSSV infections (25%) and single infections with *Vibrio anguillarum* (25%) [5]. However, other studies have revealed that the outbreak of WSSV was postponed after *P. vannamei* co-infection with WSSV and *V. harveyi* [6]. In this study, the mortality of WSSV group (97 %) was significantly higher than the co-infection group (83%) and *V. parahaemolyticus* group (34 %) ($P < 0.05$), which conflicted with the reported in *L. vannamei* after co-infection with WSSV and *V. anguillarum* [5], but was similar to previous findings in *P. vannamei* after co-infection with WSSV and *V. harveyi* [6]. The synergistic effect between WSSV and *Vibrio* may be influenced by the species of the *Vibrio* bacteria [6].

In this experiment, the WSSV copy number measured in the co-infection group was always lower than in the WSSV group. It might be the key factor of lower mortality in the co-infection group. The proliferation of WSSV result also demonstrated that the WSSV replication was controlled under co-infection conditions. It is possible that WSSV must make use of the metabolites in the host cell to assemble

nucleotides and proteins of the virus [33] after infection of the shrimp, but the metabolites were used by *V. parahaemolyticus* or the metabolism of shrimp was slowed down by *V. parahaemolyticus*. This suggests that virus couldn't replicate without the metabolites, hence the WSSV proliferation was inhibited.

ACP is a typical lysosomal enzyme and plays a key role in eliminating and hydrolyzing microbes [34]. In *Chlamys farreri* [35], the ACP activity was significantly increased at the early stage of *Vibrio anguillarum* challenge. In this experiment, the ACP activity is most sensitive to *V. parahaemolyticus* infection from 3 h after infection and reached the peak at 6 hpi. However, the ACP activity of the WSSV infected group declined at 3 hpi then increased and reached the peak at 12 hpi. The result was consistent with ACP activity in *Penaeus monodon* with WSSV in latent period on reinfection [36], but the time of appearance of the peak varied. The difference in the appearance of the peak might be associated with the dose of infection and environment. Furthermore, ACP activity in the virus infected group was always significantly lower than that of the co-injection group throughout the experimental period. In other words, the *V. parahaemolyticus* infection has, to some extent, affected ACP vitality of the shrimp. The ACP activity of the co-injection group from 3 to 96 hpi was always higher than the WSSV group. The ACP activity of the co-injection group from 6 to 24 hpi was significantly higher than that of the *V. parahaemolyticus* injected group which suggests that co-infection stimulates the immune response in *L. vannamei*. In the co-infection group, the ACP activity declined from 48 hpi, but remained significantly higher than the WSSV group. The co-infection may cause

disturbance of cell metabolism and immune function, which is consistent with the previous report in *Penaeus (Marsupenaeus) japonicus*[37].

AKP is a regulatory enzyme associated with the metabolism and can be seen as an important index in the assessment of the immune status of shrimp [38]. After an initial rise at 3 hpi, the AKP activity of WSSV-injected group decreased significantly at 6 hpi in this experiment which was similar to previous reports [39]. We observed that AKP activity in the gills of the shrimp is more sensitive to *V. parahaemolyticus* infection than WSSV infection; the AKP activity of the co-injection group varied in a similar manner.

Reactive oxygenspecies (ROS), including superoxide anion (O_2^-),hydroxyl radical (OH) and hydrogen peroxide (H_2O_2) are an important part of the innate immune defense system that is produced to help eliminate invading microbes[40].Antioxidant enzymes such as peroxidases (POD) and superoxide dismutase (SOD) either convert O_2 to H_2O_2 (SOD), convert H_2O_2 to water and oxygen bycatalase (CAT), or use H_2O_2 to oxidize substrates by various peroxidases [41].POD activity can serve as an immune index to evaluate the immune status of crustacean[42].After infection with WSSV, the POD activity of *Cherax quadricarinatus* was shown to decrease significantly [43].In this study, the POD activity in gill decreased initially in all 3 challenge groups at 3 hpi. The minimum activity of the WSSV-injected groups was recorded at 6 hpi and was significantly lower than other groups. The POD activity in the co-infection group was significantly higher than WSSV group at 6 hpi, which may have contributed to enhancing the

ability of the co-infection group to resist the infection of WSSV at 6 hpi.

SOD is an enzyme that catalyses the rapid two-step dismutation of the toxic superoxide anion to molecular oxygen and hydrogen peroxide through the alternate reduction and oxidation of the active-site metal ion [44]. A previous study indicated that a significant decrease in SOD activity occurred earlier at 3 hpi in white shrimp *L. vannamei* that received *V. alginolyticus* injection, followed by recovery after 96 hpi [45]. In this study, the SOD activity of the *V. parahaemolyticus* -injected group significantly increased at 3 hpi which conflicted with the previous report [45]. A significant decrease in SOD activity occurred in WSSV -injected group at 6 hpi. It was consistent with reports in the shrimp *Penaeus monodon* [46] and *L. vannamei* [47], which showed a decrease of SOD activity after WSSV infection. According to the study in *Fenneropenaeus indicus* [48], the lower activities of SOD may have been due to inactivation of SOD by the oxidative stress generated singlet oxygen. In the present study, the SOD activity of co-infection group and *V. parahaemolyticus* group was significantly higher than that in WSSV group at 48 hpi, which suggests that the shrimp in the co-infection and *V. parahaemolyticus* group could clear the oxyradical more efficiently compared to WSSV group, and avoid the oxidative damage induced by pathogens. Previous studies have shown an increase in activity of antioxidant enzymes in shrimp during bacterial infections, with a decrease observed during viral infection with WSSV [41].

As far as we know, viral suppression mechanisms exist in arthropods [7]. Studies had revealed that the proliferation of West Nile and chikungunya virus were

332 suppressed in individuals after infection with *Wolbachia* [50, 51]. In mosquito during
333 co-infection with *Wolbachia* and dengue virus, the TLR signaling pathway was
334 activated by ROS and expressed more immune factors than in the mosquito group
335 infected with virus only [8]. ECSIT is a multifunctional adaptor protein of TLR
336 signaling pathway, and represented a constitutive expression pattern in some tissues
337 [51]. In shrimp, MjECSIT was previously shown to be expressed in hemocyte, gill,
338 hepatopancreatic, stomach, heart, intestinal, testicular, and ovarian tissues, and the
339 expression level in gill was higher than in hemocyte [19]. The mRNA transcript of
340 LvEcsit in gill was also higher than in hemocyte (Data will be showed in another
341 paper), which are considered with the result in MjECSIT [19]. So gill was chosen for
342 the sample tissue in this study. TLR pathway is of major importance during innate
343 immunity. Most genes in TLR pathway are reported to up-regulated in the stress of
344 pathogen. ECSIT, an essential member of this pathway, was found to be significantly
345 up-regulated after *Vibrio anguillarum* challenge in *Crassostrea gigas* [52], and by
346 challenge with microorganisms (*Vibrio alginolyticus*, *Staphylococcus haemolyticus*
347 and *Saccharomyces cerevisiae*) in the Hong Kong oyster *Crassostrea hongkongensis*
348 (ChECSIT) [17]. In this study, the expression of LvECSIT was up-regulated by
349 infection with *V. parahaemolyticus* (Fig.4). The transcription level of LvECSIT in the
350 co-infection group was higher than WSSV group from 12h to 72hpi (Fig.4), which
351 was consistent with the expression pattern of Toll pathway-related genes in *Aedes*
352 *aegypti* [8]. Furthermore, the transcription levels of LvECSIT in the PBS group
353 up-regulated from 6 to 48 hpi, was consistent with MjECSIT at 6 hpi [19], and

ChECSIT at 3 and 12 hpi [17]. The difference in the kinetics of expression between these studies could be associated with the animal, dose of infection and environment. However, further study is required to elucidate the potential mechanism in shrimp. In summary, this study demonstrated that 1) shrimp in co-infection groups suffered lower mortality than groups with single infection by WSSV only; 2) the amount of WSSV in co-infection group was always lower than that of WSSV single infection group over the course of the trial; 3) ACP and AKP activity in gills of shrimp co-infected with *V. parahaemolyticus* and WSSV was significantly higher than that of WSSV single infection group from 6 to 72 hpi; ACP and AKP enzyme activity can be used as indicators of immune response to these pathogens; POD and SOD activity may not be the best indicators of immune response to WSSV - *Vibrio* infections. 4) the transcription level of LvECSIT was up-regulated in *V. parahaemolyticus* infected and multiple infection groups. This study provided information for understanding the effect of WSSV - *Vibrio* infections on survival and immune responses in shrimp. Further study is needed to develop prevention and management strategies to reduce losses caused by multiple pathogens in aquaculture.

Acknowledgments

This work was funded by the National Key R & Development Program of China (2018YFD0900501), National Natural Science Foundation of China (No. 31402344), Natural Science Foundation of Guangdong Province (No. 2017A030313174).

Reference

- [1] R.-C.o. Santiago, White spot syndrome virus (WSSV) infection in shrimp (*Litopenaeus vannamei*) exposed to low and high salinity, Archives of virology 9(159) (2014) 2213-2222.
- [2] R.E. Anaya-Rosas, M.E. Rivas-Vega, A. Miranda-Baeza, P. Pina-Valdez, M. Nieves-Soto, Effects of a co-culture of marine algae and shrimp (*Litopenaeus vannamei*) on the growth, survival and immune response of shrimp infected with *Vibrio parahaemolyticus* and white spot virus (WSSV), Fish & shellfish immunology 87 (2019) 136-143.
- [3] S. Manivannan, S.K. Otta, I. Karunasagar, I. Karunasagar, Multiple viral infection in *Penaeus monodon* shrimp postlarvae in an Indian hatchery, Diseases of aquatic organisms 48(3) (2002) 233-236.
- [4] I.K. Jang, G. Qiao, S.K. Kim, Effect of multiple infections with white spot syndrome virus and *Vibrio anguillarum* on Pacific white shrimp *Litopenaeus vannamei* (L.): mortality and viral replication, Journal of fish diseases 37(10) (2014) 911-920.
- [5] G. Qiao, D.H. Xu, Z. Wang, I.K. Jang, Z. Qi, M. Zhang, S.K. Kim, Comparison of immune response of Pacific white shrimp, *Litopenaeus vannamei*, after multiple and single infections with WSSV and *Vibrio anguillarum*, Fish & shellfish immunology 44(1) (2015) 257-264.
- [6] H.P. Le, M. Corteel, N.C. Thanh, H. Nauwynck, M. Pensaert, V. Alday-Sanz, W.V.D. Broeck, P. Sorgeloos, P. Bossier, Effect of dose and challenge routes of

- 398 *Vibrio* spp. on co-infection with white spot syndrome virus in *Penaeus vannamei*,
399 *Aquaculture* 290(1) (2009) 61-68.
- 400 [7] L. Teixeira, A. Ferreira, M. Ashburner, The bacterial symbiont *Wolbachia* induces
401 resistance to RNA viral infections in *Drosophila melanogaster*, *PLoS biology* 6(12)
402 (2008) e2.
- 403 [8] X. Pan, G. Zhou, J. Wu, G. Bian, P. Lu, A.S. Raikhel, Z. Xi, *Wolbachia* induces
404 reactive oxygen species (ROS)-dependent activation of the Toll pathway to control
405 dengue virus in the mosquito *Aedes aegypti*, *Proceedings of the National Academy of*
406 *Sciences of the United States of America* 109(1) (2012) E23-31.
- 407 [9] E. S.Loker , C. M. Adema, , Si-Ming Zhang, T. B.Kepler, Invertebrate immune
408 systems--not homogeneous, not simple, not well understood. *Immunological Reviews*
409 198(1)(2010) 10-24.
- 410 [10] P. Jiravanichpaisal, B.L. Lee, K. Soderhall, Cell-mediated immunity in
411 arthropods: hematopoiesis, coagulation, melanization and opsonization,
412 *Immunobiology* 211(4) (2006) 213-236.
- 413 [11] L. Cerenius, K. Söderhäll, The prophenoloxidase-activating system in
414 invertebrates, *Immunol. Rev.* 198 (1) (2004) 116-126.
- 415 [12] S.Y. Lee, K. Soderhall, Early events in crustacean innate immunity, *Fish*
416 *& shellfish immunology* 12(5) (2002) 421-437.
- 417 [13] Z. Man, S. Wei-Yan, W. Jun, Effects of WSSV on Enzymes Activities Related
418 with Immunity of *Marsupenaeus japonicus* Postlarvae, *Journal of Xiamen University*,
419 50(1) (2011)117-122.

- 420 [14] C.B. Sun, G. Wang, S.F. Chan, Effects of artificial infection of *Litopenaeus*
421 *vannamei* by *Micrococcus lysodeikticus* and WSSV on the activity of immunity
422 related enzymes, *Fish & shellfish immunology* 46(2) (2015) 778-786.
- 423 [15] Q. Zhai, J. Li, Y. Feng, Q. Ge , Evaluation of combination effects of Astragalus
424 polysaccharides and florfenicol against acute hepatopancreatic necrosis
425 disease-causing strain of *Vibrio parahaemolyticus* in *Litopenaeus vannamei*, *Fish &*
426 *Shellfish Immunology*, 86(5) (2019)74-383.
- 427 [16] S.Valanne, J. Wang, M. Rämet, The *Drosophila* toll signaling pathway,
428 *The Journal of Immunology*186(2) (2011) 649-656.
- 429 [17] F. Qu, Z. Xiang, F. Wang, Y. Zhang, J. Li, Y. Zhang, S. Xiao, Z. Yu,
430 Identification and function of an evolutionarily conserved signaling intermediate in
431 Toll pathways (ECSIT) from *Crassostrea hongkongensis*, *Developmental and*
432 *comparative immunology* 53(1) (2015) 244-252.
- 433 [18] P.H. Wang, D.H. Wan, Z.H. Gu, X.X. Deng, S.P. Weng, X.Q. Yu, J.G. He,
434 *Litopenaeus vannamei* tumor necrosis factor receptor-associated factor 6 (TRAF6)
435 responds to *Vibrio alginolyticus* and white spot syndrome virus (WSSV) infection and
436 activates antimicrobial peptide genes, *Developmental & Comparative Immunology*
437 35(1) (2011) 105-114.
- 438 [19] D. Ding, X.W. Chen, L.H. Kang, H.S. Jiang, C.J. Kang, Role of evolutionarily
439 conserved signaling intermediate in Toll pathways (ECSIT) in the antibacterial
440 immunity of *Marsupenaeus japonicus*, *Developmental and comparative immunology*
441 46(2) (2014) 246-254.

- 442 [20] G.Wang, Cloning of immunogene and immune responses after co-infection in
443 *Litopenaeus vannamei*, Guangdong Ocean University, 2015.
- 444 [21] T.W. Flegel, K. Sritunyalucksana, Shrimp molecular responses to viral pathogens,
445 Marine biotechnology (New York, N.Y.) 13(4) (2011) 587-607.
- 446 [22] D. Yao, L. Ruan, X. Xu, H. Shi, Identification of a c-Jun homolog from
447 *Litopenaeus vannamei* as a downstream substrate of JNK in response to WSSV
448 infection, Developmental and comparative immunology 49(2) (2015) 282-289.
- 449 [23] L. Tran, L. Nunan, R.M. Redman, L.L. Mohny, C.R. Pantoja, K. Fitzsimmons,
450 D.V. Lightner, Determination of the infectious nature of the agent of acute
451 hepatopancreatic necrosis syndrome affecting penaeid shrimp, Diseases of aquatic
452 organisms 105(1) (2013) 45-55.
- 453 [24] C.O. Lomelí-Ortega, S.F. Martínez-Díaz, Phage therapy against *Vibrio*
454 *parahaemolyticus* infection in the whiteleg shrimp (*Litopenaeus vannamei*) larvae,
455 Aquaculture 434 (2014) 208-211.
- 456 [25] W.J. Jin, J.E. Han, S.S. Giri, K.F.J. Tang, X. Zhou, L.F. Aranguren, H.J. Kim, S.
457 Yun, C. Cheng, G.K. Sang, Phage Application for the Protection from Acute
458 Hepatopancreatic Necrosis Disease (AHPND) in *Penaeus vannamei*, Indian Journal of
459 Microbiology 58(1) (2018) 114-117.
- 460 [26] H. Pang, L. Chen, R. Hoare, Y. Huang, ZaoheWu, J. Jian, Identification of DLD,
461 by immunoproteomic analysis and evaluation as a potential vaccine antigen against
462 three *Vibrio* species in *Epinephelus coioides*, Vaccine 34(9) (2016) 1225-1231.
- 463 [27] X.X. You, Y.Q. Su, Y. Mao, M. Liu, J. Wang, M. Zhang, C. Wu, Effect of high

- 464 water temperature on mortality, immune response and viral replication of
 465 WSSV-infected *Marsupenaeus japonicus* juveniles and adults, *Aquaculture* 305(1)
 466 (2010) 133-137.
- 467 [28] X.L. Liu, Q.Y. Xi, L. Yang, H.Y. Li, Q.Y. Jiang, G. Shu, S.B. Wang, P. Gao, X.T.
 468 Zhu, Y.L. Zhang, The effect of dietary Panax ginseng polysaccharide extract on the
 469 immune responses in white shrimp, *Litopenaeus vannamei*, *Fish & shellfish*
 470 *immunology* 30(2) (2011) 495-500.
- 471 [29] W. Li, Z. Yao, L. Sun, W. Hu, J. Cao, W. Lin, X. Lin, Proteomics Analysis
 472 Reveals a Potential Antibiotic Cocktail Therapy Strategy for *Aeromonas hydrophila*
 473 Infection in Biofilm, *Journal of proteome research* 15(6) (2016) 1810-1820.
- 474 [30] W. Cheng, F.M. Juang, J.C. Chen, The immune response of Taiwan abalone
 475 *Haliotis diversicolor supertexta* and its susceptibility to *Vibrio parahaemolyticus* at
 476 different salinity levels, *Fish & shellfish immunology* 16(3) (2004) 295-306.
- 477 [31] S.B. Prayitno, J.W. Latchford, Experimental infections of crustaceans with
 478 luminous bacteria related to *Photobacterium* and *Vibrio*. Effect of salinity and pH on
 479 infectiosity, *Aquaculture* 132(1) (1995) 105-112.
- 480 [32] SUNG, HungHung, HSU, ShiFang, CHEN, ChihKun, TING, YunYuan, CHAO,
 481 WeiLiang, Relationships between disease outbreak in cultured tiger shrimp (*Penaeus*
 482 *monodon*) and the composition of *Vibrio communities* in pond water and shrimp
 483 hepatopancreas during cultivation, *Aquaculture* 192(2) (2001) 101-110.
- 484 [33] T. E. Xie, Z. H. Hu, General Virology. Science publishing Company (2002)
 485 73-157.

- 486 [34] X.L. Yin, Z.J. Li, K. Yang, H.Z. Lin, Z.X. Guo, Effect of guava leaves on growth
487 and the non-specific immune response of *Penaeus monodon*, Fish & shellfish
488 immunology 40(1) (2014) 190-196.
- 489 [35] X. Wang, L. Wang, H. Zhang, Q. Ji, L. Song, L. Qiu, Z. Zhou, M. Wang, L.
490 Wang, Immune response and energy metabolism of *Chlamys farreri* under *Vibrio*
491 *anguillarum* challenge and high temperature exposure, Fish & shellfish immunology
492 33(4) (2012) 1016-1026.
- 493 [36] T. Zhang, J.H. Huang, W.G. Wen, Q.B. Yang, Z.X. Guo, Immune response of
494 enzymes activities in *Penaeus monodon* serum with WSSV in latent period on
495 reinfection, South China Fish. Sci. 9 (1) (2013) 35-42.
- 496 [37] D. R. Hewitt, P. F. Duncan, Effect of high water temperature on the survival,
497 moulting and food consumption of *Penaeus (Marsupenaeus) japonicus* (Bate, 1888),
498 Aquaculture Research 32(4) (2001)305-313.
- 499 [38]P.J. Sarlin, R. Philip, Efficacy of marine yeasts and baker's yeast as
500 immunostimulants in *Fenneropenaeus indicus*: A comparative study, Aquaculture
501 321(3) (2011)173-178.
- 502 [39] J. Du, H. Zhu, P. Liu, J. Chen, Y. Xiu, W. Yao, T. Wu, Q. Ren, Q. Meng, W. Gu,
503 W. Wang, Immune responses and gene expression in hepatopancreas from
504 *Macrobrachium rosenbergii* challenged by a novel pathogen spiroplasma MR-1008,
505 Fish & shellfish immunology 34(1) (2013) 315-323.
- 506 [40] T. Anchalee, S. Kunlaya, S. Premruethai, T. Sureerat, Discovery of immune
507 molecules and their crucial functions in shrimp immunity, Fish & shellfish

- immunology 34(4) (2013) 954-967.
- [41] M. Muñoz, R. Cedeño, J. Rodríguez, W.P.W.V.D. Knaap, E. Mialhe, E. Bachère, Measurement of reactive oxygen intermediate production in haemocytes of the penaeid shrimp, *Penaeus vannamei*, Aquaculture 191(1) (2000) 89-107.
- [42] Y. Liu, X.L. Jiang, L.U. Qing, H.S. Guan, Effects of mannuronate polysaccharide on enzymes of *Penaeus chinensis* related with immune and hemolysis, Journal of Fisheries of China 24(6) (2000). 549-553
- [43] D.L. Wang, D. Zuo, L.M. Wang, T. Sun, Q. Wang, Y.L. Zhan, Effects of white spot syndrome virus infection on immuno-enzyme activities and ultrastructure in gills of *Cherax quadricarinatus*, Fish Shellfish Immunol. 32 (5) (2012) 645-650.
- [44] C.C. Li, S.T. Yeh, J.C. Chen, The immune response of white shrimp *Litopenaeus vannamei* following *Vibrio alginolyticus* injection, Fish & shellfish immunology 25(6) (2008) 853-860.
- [45] C.C. Li, S.T. Yeh, J.C. Chen, The immune response of white shrimp *Litopenaeus vannamei* following *Vibrio alginolyticus* injection, Fish & shellfish immunology 25(6) (2008) 853-860.
- [46] G. Balasubramanian, M. Sarathi, C. Venkatesan, J. Thomas, A.S. Hameed, Studies on the immunomodulatory effect of extract of *Cyanodon dactylon* in shrimp, *Penaeus monodon*, and its efficacy to protect the shrimp from white spot syndrome virus (WSSV), Fish & shellfish immunology 25(6) (2008) 820-828.
- [47] Y.C. Lin, S.T. Yeh, C.C. Li, L.L. Chen, A.C. Cheng, J.C. Chen, An immersion of *Gracilaria tenuistipitata* extract improves the immunity and survival of white shrimp

Litopenaeus vannamei challenged with white spot syndrome virus, Fish & shellfish immunology 31(6) (2011) 1239-1246.

[48] K. Mohankumar, P. Ramasamy, White spot syndrome virus infection decreases the activity of antioxidant enzymes in *Fenneropenaeus indicus*, Virus research 115(1) (2006) 69-75.

[49] R.L. Glaser, M.A. Meola, The native *Wolbachia* endosymbionts of *Drosophila melanogaster* and *Culex quinquefasciatus* increase host resistance to West Nile virus infection, PLoS one 5(8) (2010) e11977.

[50] A.F. van den Hurk, S. Hall-Mendelin, A.T. Pyke, F.D. Frentiu, K. McElroy, A. Day, S. Higgs, S.L. O'Neill, Impact of *Wolbachia* on infection with chikungunya and yellow fever viruses in the mosquito vector *Aedes aegypti*, PLoS neglected tropical diseases 6(11) (2012) e1892.

[51] Kopp, E., Medzhitov, R., Carothers, J., Xiao, C., Douglas, I., Janeway, C.A., Ghosh, S., ECSIT is an evolutionarily conserved intermediate in the Toll/IL-1 signaltransduction pathway. Genes Dev 13(16) (1999) 2059-2071.

[52] L. Zhang, L. Li, G. Zhang. The first identification of molluscan Ecsit in the Pacific oyster, *Crassostrea gigas*, and its expression against bacterial challenge [J]. Aquaculture Research 43(8) (2012) 1-10.

553 Table1 Design of experiment for virus load, enzymes, and gene expression analysis.

Treatments	WSSV copies μL^{-1}	<i>V.pra</i> CFU mL^{-1}	No. of shrimp	Sampling Number of shrimp at hours post-injection (hpi)							
				0	3	6	12	24	48	72	96
1 PBS	-	-	40×3	1×3	1×3	1×3	1×3	1×3	1×3	1×3	1×3
2 <i>V. pra</i>	-	1.22×10^6	40×3	0	1×3	1×3	1×3	1×3	1×3	1×3	1×3
3 WSSV	3.3×10^2	-	40×3	0	1×3	1×3	1×3	1×3	1×3	1×3	1×3
4 Co-infection	3.3×10^2	1.22×10^6	40×3	0	1×3	1×3	1×3	1×3	1×3	1×3	1×3

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

Table2 Design of experiment for Lethality

Treatments	WSSV copies μL^{-1}	<i>V.pra</i> CFU mL^{-1}	No. of shrimp
1 PBS	-	-	10×3
2 <i>V. pra</i>	-	1.22×10^6	10×3
3 WSSV	3.3×10^2	-	10×3
4Co-infection	3.3×10^2	1.22×10^6	10×3

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

Table 3 Sequences of primers used in this study.

Primer name	Primer sequence(5'-3')	references
WSSV-F	AAACCTCCGCATTCCTGTGA	[28]
WSSV-R	TCCGCATCTTCTTCCTTCAT	
LvECSIT-F	ATGATTCTTATGAACGCTT	This study
LvECSIT-R	AATTTGGGCATCCAGTAC	
β -actin-F	GAAGTAGCCGCCCTGGTTGT	This study
β -actin-R	GGATACCTCGCTTGCTCTGG	

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637


638

Figure 1. Cumulative mortality in shrimp, *L. vannamei* infected by intramuscular injection with *V. parahaemolyticus* only, by white spot syndrome virus (WSSV) only, or concurrently infected with *V. parahaemolyticus* and WSSV (Co-infection) at different time intervals pi (0, 3, 6, 12, 24, 48, 72, and 96 hours). Injection with PBS served as negative control. Groups that don't share a letter are significantly different ($P < 0.05$).

Figure 2. Effect of single injection (*V. parahaemolyticus* or WSSV) and co-infection injection (*V. parahaemolyticus* and WSSV) on the amount of WSSV (copies μL^{-1}) estimated in *L. vannamei* muscle at different time intervals pi (3, 6, 12, 24, 48, 72, and 96 hours). Values are expressed as mean \pm SD. Groups that don't share a letter are significantly different ($P < 0.05$).

Figure 3. Effect of single injection (*V. parahaemolyticus* or WSSV) and co-infection injection (*V. parahaemolyticus* and WSSV) on the gill ACP(A), AKP(B), POD(C) and SOD(D) activity of *L. vannamei* at different time intervals pi (0, 3, 6, 12, 24, 48, 72, and 96 hours). Groups that don't share a letter are significantly different ($P < 0.05$).

Figure 4. Effect of single injection (*V. parahaemolyticus* or WSSV) and co-infection injection (WSSV and *V. parahaemolyticus*) on the mRNA expression of LvECSIT of *L. vannamei* at different time intervals pi (0, 3, 6, 12, 24, 48, 72, and 96 hours). Groups that don't share a letter are significantly different ($P < 0.05$).

**Fig.1**







Fig.2




722



723

724



725

726

727

728

729

730

731

732

Fig.3




Fig.4

Shrimp in co-infection groups suffered lower mortality than WSSV group.

The amount of WSSV in co-infection group was lower than in WSSV group.

ACP and AKP enzyme activity can be used as indicators to co-infection.

The transcription level of LvECSIT was up-regulated in co-infection groups.