

Regular Article

The Enduring Effects of Parental Alcohol, Tobacco, and Drug Use on Child Well-being: A Multilevel Meta-Analysis

Sofie Kuppens^{1,2}, Simon C. Moore^{3,4}, Vanessa Gross³, Emily Lowthian⁵ and Andy P. Siddaway⁶

¹Department of Public Health and Primary Care, KU Leuven, Centre for Environment and Health, Leuven, Belgium; ²Karel de Grote University of Applied Sciences and Arts, Antwerp, Belgium; ³School of Dentistry, Cardiff University, Cardiff, Wales, UK; ⁴Crime and Security Research Institute Friary House, Cardiff, Wales, UK; ⁵DECIPHer, Cardiff School of Social Sciences, Cardiff University, Cardiff, Wales, UK and ⁶Stirling Management School, University of Stirling, Stirling, Stirlingshire, Scotland, UK

Abstract

The effects of psychoactive substance abuse are not limited to the user, but extend to the entire family system, with children of substance abusers being particularly at risk. This meta-analysis attempted to quantify the longitudinal relationship between parental alcohol, tobacco, and drug use and child well-being, investigating variation across a range of substance and well-being indices and other potential moderators. We performed a literature search of peer-reviewed, English language, longitudinal observational studies that reported outcomes for children aged 0 to 18 years. In total, 56 studies, yielding 220 dependent effect sizes, met inclusion criteria. A multilevel random-effects model revealed a statistically significant, small detriment to child well-being for parental substance abuse over time ($r = .15$). Moderator analyses demonstrated that the effect was more pronounced for parental drug use ($r = .25$), compared with alcohol use ($r = .13$), tobacco use ($r = .13$), and alcohol use disorder ($r = .14$). Results highlight a need for future studies that better capture the effect of parental psychoactive substance abuse on the full breadth of childhood well-being outcomes and to integrate substance abuse into models that specify the precise conditions under which parental behavior determines child well-being.

Registration: PROSPERO CRD42017076088

Keywords: alcohol, children, meta-analysis, parent, substance abuse, tobacco, well-being

(Received 14 June 2018; revised 4 February 2019; accepted 21 March 2019)

The effects of psychoactive substance abuse are not limited to the user but extend to the entire family system and society at large. In line with the growing body of research on the detrimental influence of adverse childhood experiences, children of users are particularly at risk, with effects on their health, well-being, and their own use of tobacco, alcohol, or drugs (e.g., National Center on Addiction and Substance Abuse, 2005). Millions of children are likely to be affected by parental psychoactive substance abuse, as estimates suggest that 12.3% of US children aged 17 years or younger reside in a home with at least one parent with a substance abuse disorder (Lipari & Van Horn, 2017). In Europe, the estimates for children under the age of 20 years living with alcohol abusing parents varies greatly and ranges from 5.7% in Finland, 10.5% in Denmark, 15.4% in Germany, and 17% to 23% in Poland (European Monitoring Centre for Drugs and Drug Addiction, 2008).

A large body of evidence spanning several decades has documented that the children of parents who abuse alcohol, tobacco, and drugs (henceforth collectively referred to as parental

substance abuse) are more likely to develop a variety of emotional, behavioral, physical, cognitive, academic, and social problems in the short and long run (e.g., Barnard & McKeganey, 2004; Straussner & Fewell, 2011). For example, passive tobacco exposure has been linked to somatic health problems in children and adolescents, together with an increased risk of children's own tobacco use initiation and dependence (Hussong et al., 2008). In addition, parental substance abuse has been linked to family breakdown, which is a key risk factor for children's poor mental health (Mallett, Rosenthal, & Keys, 2005; Størksen, Røysamb, Moum, & Tambs, 2005). It has also been linked to a reduction in the quality of the parent-child relationship and maladaptive relationship models that can be detrimental to the development of later peer relationships (Fearon, Bakermans, Kranenburg, Van Ijzendoorn, Lapsley, & Roisman, 2010; Hoeve et al., 2012).

Parental substance abuse has also been associated with a reduction in the extent that parents monitor their children, which may undermine parents' ability to provide a safe and nurturing home environment (Barnard & McKeganey, 2004). Instability with respect to employment, family structure, housing, childcare, and household finances has also been shown to co-occur with parental substance abuse, with consequences that extend beyond the family environment to influence children's social functioning (Berger, Paxson, & Waldfogel, 2009; De Goede, Branje, Delsing, & Meeus, 2009; Giesbrecht, Cukier, &

Author for Correspondence: Simon C. Moore, School of Dentistry, Heath Park, Cardiff University, Cardiff, CF14 4XY, UK; Email: mooresc2@cardiff.ac.uk

Cite this article: Kuppens S, Moore SC, Gross V, Lowthian E, Siddaway AP (2019). The Enduring Effects of Parental Alcohol, Tobacco, and Drug Use on Child Well-being: A Multilevel Meta-Analysis. *Development and Psychopathology* 1–14. <https://doi.org/10.1017/S0954579419000749>

Steeves, 2010; Lander, Howsare, & Byrne, 2013; Martin, Razza, & Brooks-Gunn, 2012; Öberg, Jaakkola, Woodward, Peruga, & Prüss-Ustün, 2011; Parsons, Adler, & Kaczala, 1982).

There appears to be a general consensus amongst researchers, clinicians, and policy makers that parental substance abuse negatively affects child well-being (Bountress & Chassin, 2015; Hussong, Huang, Curran, Chassin, & Zucker, 2010; McGrath, Watson, & Chassin, 1999; Peterson *et al.*, 2006; Puttler, Zucker, Fitzgerald, & Bingham, 1998; Rossow, Keating, Felix, & McCambridge, 2016). However, the extent and nature of this relationship is currently unknown, because findings in the literature vary considerably in magnitude and studies have typically focused on a subset of child well-being outcomes or a specific parental substance abuse type. Furthermore, there is considerable variation in the way that parental substance abuse and child well-being outcomes have each been operationalized. A systematic synthesis of the available evidence using an overarching framework is needed to quantify the extent to which parental substance abuse predicts detrimental child well-being outcomes over time in order to draw more general conclusions and to determine the degree of heterogeneity in this relationship while also identifying factors that could explain any inconsistencies. In doing so, it would also be possible to identify gaps in the research literature and key directions for future research.

The present meta-analysis

To address these important knowledge gaps, we conducted a systematic review and meta-analysis of the longitudinal relationship between parental psychoactive substance abuse and child well-being that included multiple substance abuse types and well-being domains. We focused on longitudinal studies, as this provides insight into the directionality of any effect and the long-term characteristics of any association. We used a state-of-the-art multilevel meta-analytic approach that allowed us to model the dependency among effect sizes, which is common in primary studies, in part because they include multiple outcome measures and multiple informants, or they report on multiple family members. Given the broad and comprehensive nature of this meta-analysis, heterogeneity within the results was expected. Therefore, the following study and sample characteristics that could potentially moderate the strength of the relationship between substance abuse and well-being were examined.

Parental substance abuse refers to the consumption of psychoactive substances, including licit and illicit substances, of which alcohol (Rossow *et al.*, 2016) and tobacco (Saulyte, Regueira, Montes-Martínez, Khudyakov, & Takkouche, 2014) are the most frequently used. Given differences in the legal and social status, addictive potential, and cognitive effects, the effect on family members has been proposed to vary according to the particular substance consumed (Straussner, 1994; Straussner & Fewell, 2011). Apart from the different types, the use of such substances have commonly been considered to range on a continuum from recreational use to chronic dependence (Straussner, 2004). Substance dependence may vary in severity from mild to severe and refers to compulsive and continued use irrespective of any adverse consequences. We hypothesised that the association with child well-being would be less pronounced for the use of licit substances of a recreational nature.

Well-being is a nebulous term that is informed by personal, cultural, and other factors (Diener, Oishi, & Lucas, 2003), and there is no clear distinction between well-being and quality of

life or mental health problems (Siddaway, Taylor, & Wood, 2018; Siddaway, Wood, & Taylor, 2017). Well-being is typically considered a multidimensional construct (Gallagher, Lopez, & Preacher, 2009; Keyes, 2009; Ryff & Keyes, 1995) that includes both subjective and objective features (Newman, Tay, & Diener, 2014) and, for young people, can refer to domains as disparate as relationships, health, activities, finance, education, and skills (Bradshaw, 2016). Because existing research has largely focused on a specific well-being domain, a comprehensive examination of whether and how well-being subtype moderates the substance abuse–well-being relationship is currently lacking. We sought to address this gap by conceptualizing child well-being as a broad, multidimensional construct that involves physical, psychological, cognitive, social, and economic subdomains. This broad conceptualization is similar to the World Health Organization's (WHO, 2004, p. 10) definition of mental and physical health ("a state of complete physical, mental and social well-being").

Although parental substance abuse can be detrimental at any point in a child's life, it is feasible that its effects could vary according to the age of the child. For example, parental tobacco use in a child's first year of life has been associated with a greater risk of physical health symptoms (Mitchell *et al.*, 2012). For substance abuse, it has been argued that, compared with younger children, adolescents are at greater risk due to more prolonged exposure to parental substance abuse (Straussner & Fewell, 2011). We therefore examined whether effects differed according to the age of the child.

Although substance abuse by the mother and the father are both important, previous research has often found more pronounced associations with multiple adverse child outcomes for maternal substance abuse (Mitchell *et al.*, 2012; Straussner & Fewell, 2011). These gender differences may be due to children spending more time in the presence of their mothers than their fathers, as mothers traditionally take a more active role in child rearing. It is also possible that parental substance abuse effects may manifest differently in boys and girls. For example, associations between parental smoke exposure and child mental health outcomes have been found to be more apparent for boys (Bandiera, Richardson, Lee, He, & Merikangas, 2011). Given previous findings on gender differences, we examined whether the gender of parent and child moderates the substance abuse–well-being relationship.

The choice of informants in behavioral research has been a subject of a long-lasting debate. For many measures of child functioning and parenting behavior, inter-informant agreement is low-to-moderate because parent and child reports are often discrepant (De Los Reyes *et al.*, 2015; Kuppens, Grietens, Onghena, & Michiels, 2009b). Similarly, for licit and illicit substances, there is variation in prevalence by data collection mode (Beck, Guignard, & Legleye, 2014). Such inconsistent research findings may reflect method bias related to the specific informant or data collection mode (e.g., socially desirable responding, shared method variance; Kuppens, Grietens, Onghena, & Michiels, 2009a; Podsakoff, MacKenzie, Lee, & Podsakoff, 2003; Tourangeau & Yan, 2007). As the potential method effect remains unexplored for the parental substance abuse–child well-being relationship, we examined the moderating role of informant and data collection mode.

Finally, the interval adopted in longitudinal studies can vary considerably, which in turn may influence the strength of the relationship between substance abuse and well-being over time for

several reasons (Collins & Horn, 1991). The strength of the relationship could potentially decrease over time, as there is increasing opportunity for other cognitive, biological, and environmental variables to exert an effect. In addition, respondents may more easily remember their answers to a previous assessment when the time interval is shorter. We hypothesized that longitudinal relationships would be less pronounced as the time interval between parental substance abuse exposure and assessment of child well-being outcomes increased.

Method

This meta-analysis was conducted in accordance with guidelines for conducting a systematic review (Siddaway, Wood, & Hedges, 2019), the Preferred Reporting Items for Systematic Reviews and Meta-analyses (Moher, Liberati, Tetzlaff, Altman, & The PRISMA Group, 2009) and Meta-Analysis of Observational Studies in Epidemiology (Stroup et al., 2000) standards.

Literature search

Four electronic databases (PubMed, Medline, Embase, PsychInfo) were searched from inception to June 26, 2017. In addition, reference lists from eligible publications and relevant reviews were hand-searched. Comprehensive search strategies were developed by combining key and index terms covering the concepts of *parental substance abuse AND child well-being* (or related terms, see Supplement).

Inclusion and exclusion criteria

Peer-reviewed, English language, longitudinal observational studies (Morrison et al., 2012) were eligible for inclusion. Studies were required to include at least one association over time between a measure of parental substance abuse (measured at Time 1) and well-being (measured at Time 2) for children aged 18 years or younger. As previous research revealed polysubstance abuse in parental substance abuse (Hussong et al., 2008; Straussner & Fewell, 2011), we included recreational and disordered (including clinically defined dependence) alcohol, tobacco, and drug use. We included studies reporting comorbid substance use, as this is frequently observed in dependent users and excluding such groups might bias results towards being more conservative and less generalizable to the general population.

We operationalized child well-being broadly according to five distinct domains, in-keeping with a narrative literature review of how well-being can be conceptualized (Pollard & Lee, 2003), and in order to achieve a comprehensive overview of subjective and objective well-being of children, namely, (a) physical (including overall health and risks to health); (b) psychological (including emotional and mental states of mind); (c) cognitive (including capacity for learning and recall, academic achievement, and learning disability); (d) social (including social relationships and behaviors and anti-social behavior); and (e) economic (whether outside financial support is required) well-being. As we attempted to provide estimates of the association in the general population, studies were excluded if they were sampled from specific groups experiencing conditions that were likely to significantly affect associations between parental substance abuse and child well-being (e.g., detained adolescents, parents diagnosed with HIV). One author (V.G.) screened titles and abstracts, and S.M. rechecked extraction methods. Two authors (V.G. and S.M.)

screened full text articles for inclusion, $N = 381$, $\kappa = .93$, 95% CI [.92, .99]. References were exported and managed with Endnote X7.

Coding

To capture as much variation in substance abuse type, effect sizes were categorized into "Alcohol," "Alcohol Use Disorder," "Drugs," "Drug Use Disorder," "Tobacco," and a non-descript category, "Any substance." We used "Use Disorder" to identify parents whose use of substances was clinically significant and recorded using a clinically validated instrument. Child well-being was coded according to the aforementioned five domains, namely physical, psychological, social, cognitive, or economic well-being.

Several other study and outcome characteristics were extracted from each study: year; country from which the sample was drawn; sample size; retention rate; sample type (general population or specific subpopulation such as school children); parent and child gender, child age (0–5, 6–11, or 12–18 years); data collection mode for child and parent variables (interview, questionnaire); measurement instrument (standardized diagnostic, validated but nondiagnostic, or unstandardized); informant (parent, child, or teacher report); and follow-up duration (number of data collection points and the time in months between first and last assessment).

Data analysis

Effect size calculations

Pearson correlations (r) were computed to represent associations such that a positive correlation denoted that higher levels of parental substance abuse were associated with lower child well-being over time. Results reported in another metric were transformed to r whenever possible (Wilson, 2017). Study authors were contacted for additional information when an effect size or data to construct an effect size were not reported. Before pooling effect sizes, the correlations were transformed using Fisher's Z_r transformation (Rosenthal, 1991). Pooled Z_r expressions were transformed back to r expressions for reporting.

Meta-analytic integration

As most studies (71%) reported more than one relevant effect size, the independence assumption that underlies traditional meta-analysis was violated. Common approaches to addressing such dependency include choosing only one effect size from many or averaging across effect sizes within studies. Although such approaches allow one to adopt traditional meta-analytic techniques, they also result in a loss of information, rule out critical analyses of within-study moderators (e.g., well-being outcome differences), and may distort meta-analytic results (e.g., Cheung & Chan, 2004). We therefore employed a multilevel random-effects meta-analysis that permitted incorporating all of the relevant effect sizes from each study, while accounting for dependency among these effect sizes with SAS PROC HP MIXED (Littell, Milliken, Stroup, Wolfinger, & Schabenberger, 2006; Van den Noortgate, Lopez-Lopez, Marin-Martinez, & Sanchez-Meca, 2013).

The meta-analysis was implemented as a three-level model that models sampling variation for each effect size (Level 1), within-study variation (Level 2), and between-study variation (Level 3) (Van den Noortgate et al., 2013). We first computed an overall estimate of the longitudinal association between




Figure 1. Flow diagram of study identification and selection, including reasons for exclusion.

parental substance abuse and child well-being in a random-effects model. Between-study heterogeneity was reflected by the between-study variance (Level 3, representing differences between studies), but the three-level model also yielded an estimate of the within-study variance (Level 2, representing differences between effect sizes within studies). A likelihood ratio test was used to test the variation in effect sizes between and within studies, while the ratio of the variance at each level and the total variance was computed to reflect the amount of variance situated between and within studies. Subsequently, six substantive (study region, child age, child gender, parent gender, parental consumption type, child well-being type) and six methodological moderators were examined with grand-mean centering applied to continuous moderators. Separate mixed-effects models were fitted for each moderator variable to avoid inflating Type II error (Raudenbush & Bryk, 2002), and subgroup analyses were only conducted when data from at least four studies were available (Fu et al., 2011).

Risk of bias

In the absence of a standard risk of bias tool for longitudinal observational studies (see Siddaway et al., 2019), six criteria were adopted to reflect the methodological rigor of each study (Faragher, Cass, & Cooper, 2005; Higgins & Green, 2011; Wong, Cheung, & Hart, 2008): whether the study sample size provided adequate statistical power (calculated as at least $N = 193$ to provide a power of .80 in order to detect $r = .20$ with $\alpha = .05$), retention rate, representativeness of the sample, standardized assessment tool for parental substance abuse, standardized assessment tool for child well-being, and whether associations were adjusted for confounding. We tested whether effect sizes differed according to the separate criteria.

Three strategies were used to assess publication bias. First, a funnel plot was created to visually search for evidence of bias,

which would be apparent in an asymmetrical plot. Next, asymmetry was assessed using Egger's weighted regression test (Egger, Smith, Schneider, & Minder, 1997; Torgerson, 2006). Third, a sensitivity analysis was performed, which applies different a priori weight functions to correct the population effect size estimate for different types and severities of potential publication bias (Vevea & Woods, 2005). A mean study effect size was used in the publication bias analyses because independence is assumed in these methods and it is not yet possible to account for effect size dependency in these tests.

Results

Study characteristics

The meta-analysis included 56 studies and 220 effect sizes. A flow diagram of study identification and selection is presented in Figure 1. All 56 studies explicitly stated or used terminology implying that children and parents cohabited. Sample sizes varied from 83 to 49,5. Most studies drew samples from a general (34%) or school (27%) population; the remaining studies sampled from clinically dependent parents (22%) or used a specific sampling frame (e.g., 16% of families living in rural areas). Most studies ($n = 35$) were conducted in North America, followed by Europe ($n = 9$), Australasia ($n = 9$), Asia ($n = 2$), and South America ($n = 1$).

Drug use was variously defined and predominantly included cannabis use. Most of the effect sizes (78%) examined outcomes in young people aged 12 to 18 years and on average included 45% female children. For parental consumption, 37% and 27% of reported effect sizes referred to maternal and paternal behavior, respectively, and 36% reported on the consumption for both parents combined. Effect sizes were evenly split across alcohol (38%) and tobacco (32%) use; drug use and alcohol use disorder were assessed in 6% and 19% of the effect sizes, respectively. The

remaining effect sizes (5%) considered “any substance abuse.” No studies considered prescription medication. Information on parental substance abuse was mainly retrieved directly from parents (72%) or their children (25%). Most child well-being effect sizes concerned physical (74%) or psychological (19%) well-being, while social, cognitive, and economic well-being were assessed less frequently (7%). Well-being was mainly child (76%) or parent reported (15%). Time between parental substance abuse measurement and child well-being outcome averaged 60.26 months across all effect sizes ($SD = 47.90$, range = 3–190 months). Table 1 presents descriptive and mean effects sizes for the 56 included studies.

Parental substance abuse and child well-being

The three-level random-effects meta-analysis yielded a statistically significant, small association between parental substance abuse and child well-being ($r = .15$), indicating that parental substance abuse was associated with poorer child well-being levels over time. Effect sizes differed significantly both between, $\sigma_v^2 = .003$; $\chi^2(1) = 17.26$, $p < .001$, and within, $\sigma_u^2 = .009$; $\chi^2(1) = 14,804.78$, $p < .001$, studies; 21% of the total variance was attributable to differences between studies, and 64% of the total variance was due to differences within studies.

Moderator analyses

Table 2 presents the results for substantive and methodological moderators. Parental “drug use disorder” ($n = 2$), “any substance abuse” ($n = 2$), “social” ($n = 1$) and “economic” ($n = 0$) child well-being subgroups were excluded from moderator analyses due to an insufficient number of studies. Significant differences were found across parental substance abuse type, with pairwise comparisons revealing that the mean association was significantly stronger for parental drug use compared to alcohol use, $t(205) = 3.75$, $p < .001$, tobacco use, $t(205) = 3.66$, $p < .001$, and alcohol use disorder, $t(205) = 2.83$, $p = .005$. Other pairwise differences were not statistically different. Adding the parental substance abuse type moderator and the child well-being domain moderator explained 8.6% of the within-study variance. We did not find significant effects for the other substantive or methodological moderators. For substances other than tobacco or alcohol, cannabis was specifically identified in 71% of the drug-related category (k studies = 5; n (effect sizes) = 10). Sensitivity analysis yielded a significant effect of parental cannabis use on child well-being ($r = .23$, 95% CI [.14, .32], $p < .001$).

Risk of bias

Three-level mixed-effect models did not reveal a statistically significant effect on the parental substance abuse–child well-being association for the methodological moderators: power, $t(218) = 0.97$, $p = .332$; retention rate, $t(159) = 0.06$, $p = .952$; sample representativeness, $t(122) = -1.92$, $p = .058$; properties of the parental substance use assessment tool, $t(212) = 0.11$, $p = .913$; properties of the child well-being assessment tool, $F(2, 215) = 0.17$, $p = .844$; or adjustment for confounding, $t(218) = 0.96$, $p = .336$.

The funnel plot was funnel-shaped (Figure 2), and Egger’s method found no statistically significant asymmetry, $t(54) = 0.44$, $p = .664$, suggesting no indication of publication bias. The sensitivity analyses corroborated these findings. Analyses revealed that the adjusted ($r = .14$) and unadjusted ($r = .15$) population

effect size estimates were almost identical under different a priori weight functions. Overall, these analyses indicate that it is highly unlikely that publication bias influenced the meta-analytic results.

Discussion

We conducted a meta-analysis to examine the association between various indices of parental substance abuse and child well-being over time. A total of 56 studies yielding 220 dependent effect sizes met our inclusion criteria. We found a statistically significant, small longitudinal relationship between parental substance abuse and child well-being, with significant differences both between and within studies. The association between parental substance abuse and child well-being did not differ according to the length of the period between recording the parental substance abuse and the subsequent child well-being outcome, suggesting an enduring effect over time (Evans, Li, & Whipple, 2013). These results indicate that parental substance abuse is a risk factor for subsequent child well-being. The prevalence of substance use and parenthood points to the clinical significance of this risk factor. However, we caution readers to avoid interpreting parental substance abuse as a *causal* risk factor (see Kraemer et al., 1997 for a discussion of risk factor typology) and explicitly recognize that parental substance abuse and child well-being may be caused by the same third variable (e.g., financial or marital problems) or that both concepts can have a reciprocal relationship.

The longitudinal associations observed in the current meta-analysis were heterogeneous, with significant differences relating to the parental substance abuse type. Parental alcohol use (whether dependent or not) had similar secondary effects on children as parental tobacco use, suggesting that recreational alcohol use is as harmful for child well-being as secondary tobacco use and alcohol use disorder. Parental drug use yielded the strongest effect, which could (partially) be due to the illegal nature of these substances that could result in additional disruptive effects on families, such as fines, arrest, or custodial sentences. It should be noted that none of the studies considered prescription drug use, which is an important shortcoming given concerns over prescription opioids (Cicero, Ellis, Surratt, & Kurtz, 2014). Parental substance abuse had similar effects on the physical, psychological, and cognitive well-being of their children. The effects did not differ for fathers, mothers, sons, or daughters, nor according to the age of the child. Although small in magnitude, the effect of parental substance abuse on child well-being has a clear public health significance given the high prevalence of substance abuse (Cumming, 2014).

Our findings corroborate the conjecture that the effect of parental substance abuse on child well-being manifests through the lens of the family environment, with broad detrimental effects on physical, psychological, and cognitive well-being. The substantial unexplained variance that remained in our meta-analysis combined with an effect across substance abuse types could arise from unmeasured factors influencing child well-being that are common across all types of parental substance abuse. If targeting parental substance abuse is to be used in family-centred interventions to improve child outcomes (Calhoun, Conner, Miller, & Messina, 2015), then it will be necessary to further investigate how parental substance abuse affects child outcomes through the parent-child environment. More generally, working with the context of substance abuse, including the family environment in activities is increasingly seen as important in supporting recovery (Friedmann, Hendrickson, Gerstein, & Zhang, 2004;

Table 1. Descriptives and mean effect size for studies included in the meta-analysis ($N=56$).

Study	Country	Child age (years)	Gender child (% female)	Gender parents	Parental consumption	Child Well-being	Informant (parental consumption)	Informant (child well-being)	Method of assessment	Time lag (months)	Mean N	Mean r	Lower 95% CI	Upper 95% CI
Adalbjarnardottir & Rafnsson, 2002	Iceland	12–18	54%	B	T, A	Tobacco, Alcohol	Child	Self-report	Q	36	928	.10	.03	.17
Barman, et al., 2004	Finland	12–18	50%	MF	T	Tobacco	Self-report	Self-report	Q	36	4739	.20	.13	.27
Barnow, et al., 2007	USA	12–18	48%	B	A	Behavioral problems	Self-report	Parent	I	60	83	.06	–.15	.26
Bor, McGee, & Fagan, 2004	Australia	12–18	48%	M	T	Antisocial behavior	Self-report	Mother	Q	168	5278	.09	–.10	.28
Brook, et al., 2001	Colombia	12–18	49%	MF	T, D	Drug use	Self-report	Self-report	I	24	1374	.12	.02	.21
Bufferd et al., 2014	USA	6–11	46%	B	DD	Depression	Self-report	Self-report	I	36	456	.15	.00	.29
Burke, et al., 2001	Australia	12–18	53%	MF	A	Alcohol	Self-report	Self-report	Q		798	.26	.15	.37
Chaffin, et al., 1996	USA	missing	43%	B	A	Physical abuse, Neglect	Self-report	Parents	I	12	7103	.30	.17	.41
Colder, et al., 1997	USA	12–18	43%	MF	AD	Heavy alcohol use	Self-report	Self-report	Q	36	428	.17	.02	.31
Connolly, et al., 1992	USA	12–18	missing	M	A	Behavior problems	Self-report	Self-report, Teacher	I	48	652	.05	–.09	.19
Crandall, et al., 2006	USA	0–5	0%	M	A	Head injury	Self-report	Mother	I	12	3808	.21	.02	.38
Cranford, et al., 2010)	USA	12–18	31%	MF	AD	Alcohol	Self-report	Self-report	I	72	148	.21	.12	.29
Dwyer, et al., 1999	Tasmania	0–5	49%	MF	T	Sudden Infant Death Syndrome	Self-report	Parent	I	3	9788	.26	.19	.32
Eiden, et al., 2007	USA	0–5	48%	B	A	Externalizing behavior	Self-report	Mother, Father, Teacher	I	48	227	.03	–.06	.13
Ellickson, et al., 2001	USA	12–18	missing	B	A	Problem Alcohol	Self-report	Self-report	Q	60	4283	.04	–.16	.22
Fergusson, et al., 1993	New Zealand	12–18	missing	M	T	Conduct disorder, Attention deficit, Total disruptive behavior	Self-report	Mother, Teacher	Q	84	960	.16	.07	.23
Fergusson, et al., 1994	New Zealand	12–18	missing	B	D	Multiple problem behaviors	Self-report	Self-report	Q	48	942	.07	–.13	.26
Fergusson, et al., 1996	New Zealand	12–18	missing	B	T	Depression, Nicotine dependence	Self-report	Self-report	Q	48	942	.10	–.04	.24
Frank et al., 2014	USA	12–18	0%	B	D	Drug use	Other	Self-report	I	180	157	.34	.11	.54
Fuller et al., 2003	USA	6–11	53%	B	AD	Aggression	Self-report	Parent	Q	36	186	.15	–.09	.37
Griesler, et al., 2011	USA	12–18	63%	B	T	Nicotine dependence	Self-report	Self-report	I	24	814	.15	–.05	.34
Gritz et al., 2003	USA	12–18	missing	B	T	Tobacco	Self-report	Self-report	Q	12	659	.11	–.10	.30

Hanewinkel et al., 2014	Germany	12–18	57%	B	A	Alcohol	Self-report	Q	12	13642	.11	–.03	.24	
Heron et al., 2013	UK	12–18	51%	M	A, T, D	Problematic drug use, Non-problematic drug use	Self-report	Self-report	Q	190	4159	.25	.18	.32
Hoffmann & Cerbone, 2002	USA	12–18	Mixed/ boys only/ girls only	B	DD	Drug abuse, Alcohol abuse	Self-report	Self-report	I	96	804	.25	.11	.38
Hussong & Chassin, 1997	USA	12–18	51%	B	AD	Heavy alcohol use	Self-report	Self-report	I	36	293	.08	–.05	.20
Johnson & Leff, 1999	USA	0–5	51%	B	T	IQ	Self-report	Self-report	I	48	122	.03	–.23	.28
Kaplow, et al., 2002	USA	12–18	51%	B	D	Drug use	Self-report	Self-report	I	84	295	.24	.02	.43
Knudsen, et al., 2015	Norway	0–5	52%	M	A	Externalising & Internalising behavior problems	Self-report	Mother	Q	18	49553	.38	.26	.49
Korhonen et al., 2008	Finland	12–18	61%	MF	A, T	Drug use	Self-report	Self-report	Q	72	4138	.14	.08	.21
Kotch et al., 2014	USA	12–18	49%	B	T	Externalising behavior problems	Self-report	Self-report	I	72	915	.10	–.10	.29
Lessard et al., 2014	USA	12–18	52%	MF	T, D, AD	E-cigarette use	Self-report	Self-report	Q	48	136	.10	–.01	.21
Li, et al., 2002	USA	6–11	missing	B	A, T, D, A	Alcohol, Tobacco, Drug use, Any substance use	Self-report	Self-report	Q	1551	.14	.09	.19	
Lindsey et al., 2008	USA	6–11	0%	missing	A	Externalising & Internalising behavior problems	Self-report	Self-report	I	48	514	.15	.01	.29
Loukas, et al., 2001	USA	6–11	52%	B	AD	Externalising behavior	Self-report	Parent	I	36	208	.27	.04	.47
Malone, et al., 2002	USA	12–18	missing	F	AD	Any substance disorder, Alcohol disorder, Drug diagnosis, Nicotine dependence, Conduct disorder, Oppositional defiant disorder, ADHD, Depression, Any disruptive disorder	Self-report	Self-report	I	36	1020	.19	.13	.25
McGrath, et al., 1999	USA	12–18	49%	B	AD	Academic achievement	Self-report	Self-report, Teacher	I	24	454	.13	.03	.22
Miller, 2001	USA	6–11	57%	M	T	Asthma	Self-report	Mother, Other	Q (parents), Other (children)	36	3938	.08	–.05	.21
Morgenstern, et al., 2013	Germany	12–18	missing	B	T	Tobacco, School performance	Self-report	Self-report	Q	36	1320	.12	.03	.21
Naicker, et al., 2012	Canada	12–18	27%	B	A	Emotional disorder	Self-report	Self-report	Q	937	.10	–.10	.29	

(Continued)

Table 1. (Continued.)

Study	Country	Child age (years)	Gender child (% female)	Gender parents	Parental consumption	Child Well-being	Informant (parental consumption)	Informant (child well-being)	Method of assessment	Time lag (months)	Mean <i>N</i>	Mean <i>r</i>	Lower 95% CI	Upper 95% CI
Nigg et al., 2006	USA	12–18	missing	MF	AD	ADHD, Response inhibition, Set shifting and working memory, Conduct symptoms, IQ	Self-report	Self-report, Teacher	I	180	498	.06	.00	.13
Pagani, et al., 2016	Canada	6–11	57%	M	T	Being overweight/obese	Self-report	Other	I (parents), Other (children)	36	1323	.14	.00	.27
Power, et al., 2005	USA	12–18	missing	F	A	Alcohol	Self-report	Self-report	Q	6	743	.00	–.20	.20
Roosa, et al., 1990	USA	12–18	missing	B	AD	Anxiety, Depression	Child	Self-report	Q	3	145	.30	.13	.45
Selya, et al., 2012	USA	12–18	57%	MF	T	Tobacco	Self-report, Child	Self-report	Q	42	1092	.20	.06	.33
Shortt, et al., 2007	Australia	12–18	missing	B	A, T	Alcohol	Self-report	Self-report	Q	12	1166	.13	–.01	.26
Thacher et al., 2016	Sweden	12–18	Girls/ boys analysed separately	MFB	T	Allergies	Child, Spouse	Parents	Q	144	3316	.06	–.02	.13
Thompson & Wilsnack, 1987	USA	missing	0%	MF	A	Alcohol	Child	Self-report	Q	48	712	.09	.06	.13
Ting, et al., 2015	Taiwan	12–18	0%	B	A	Alcohol	Self-report	Self-report	Q	48	496	.13	–.02	.27
Tjora, et al., 2011	Norway	12–18	54%	MF	T	Alcohol	Self-report	Self-report	Q	60	1052	.18	.04	.31
Unger et al., 2006	USA	6–11	53%	B	T	Tobacco		Self-report	Q	12	1940	.10	–.10	.29
Van Der Vorst et al., 2013	Canada	12–18	48%	B	A	Alcohol	Child	Self-report	Q	12	1142	.14	–.06	.33
Vella, et al., 2015	Australian	12–18	53%	M	T	Health-related Quality of Life	Self-report	Self-report	I	120	2785	.25	.12	.38
Windham et al., 2004	USA	0–5	29%	M	A, D	Physical abuse, Self-esteem	Self-report	Self-report	I		541	.09	–.02	.19
Wong, et al., 2011	USA	12–18	47%	MF	AD	Suicidal thoughts, Self-harm	Self-report	Self-report	I	96	392	.02	–.10	.15
Xie, et al., 2013	China	12–18	missing	B	T	Tobacco	Child	Self-report	Q		3572	.08	–.03	.19

Note. *N* = average sample size available for analyses; Gender parents: M = mother, F = father, C = child, MF = mother and father separately, MFB = both parents included but not reported on separately; Parental consumption: T = tobacco, A = alcohol, D = drug use, DD = drug use disorder, AD = alcohol use disorder, A = any substance use; Method of assessment: I = interview, Q = Questionnaire.

Table 2. Results of the three-level meta-analysis models

	<i>k</i> studies	<i>n</i>	Effect Size		Test Statistic
			<i>r</i>	95% CI	
Random-effects model					
Overall effect size	56	220	.15	[.12, .17]	<i>t</i> (219) = 12.94, <i>p</i> < .001
Mixed-effects models Substantive Moderators					
Region					<i>F</i> (2, 208) = 1.01, <i>p</i> = .365
North America	35	138	.13	[.10, .16]	
Europe	9	46	.16	[.12, .21]	
Australasia	9	27	.17	[.12, .22]	
Age of child					<i>F</i> (2, 215) = 0.62, <i>p</i> = .541
0 to 5 years	6	22	.18	[.11, .24]	
6 to 11 years	8	25	.14	[.08, .20]	
12 to 18 years	41	171	.14	[.11, .16]	
Gender of child (% female)	41	181			<i>t</i> (179) = -0.16, <i>p</i> = .873
Gender of parent					<i>F</i> (2, 217) = 0.86, <i>p</i> = .426
Father	25	82	.16	[.13, .19]	
Mother	16	59	.14	[.10, .18]	
Both	31	79	.13	[.10, .17]	
Parental substance abuse					<i>F</i> (3, 205) = 50.15, <i>p</i> = .002
Alcohol ^a	21	83	.13	[.09, .16]	
Tobacco ^b	25	70	.13	[.10, .16]	
Drug use ^{a,b,c,d}	8	14	.25	[.19, .31]	
Alcohol use disorder ^c	11	42	.14	[.09, .19]	
Child well-being dimension					<i>F</i> (2, 214) = 20.65, <i>p</i> = .073
Physical	38	162	.15	[.12, .17]	
Psychological	16	42	.15	[.11, .19]	
Cognitive	4	13	.06	[-.02, .14]	
Methodological Moderators					
Informant on parental substance abuse					<i>t</i> (212) = -10.08, <i>p</i> = .280
Parent self-report	47	159	.15	[.13, .18]	
Child report	7	55	.12	[.06, .17]	
Informant on child well-being					<i>F</i> (2, 203) = 10.53, <i>p</i> = .220
Child self-report	42	166	.13	[.11, .16]	
Parent report	12	33	.17	[.12, .22]	
Teacher report	4	7	.10	[.02, .19]	
Shared informant					<i>t</i> (218) = 10.34, <i>p</i> = .182
Not shared	44	142	.14	[.11, .16]	
Shared	17	78	.17	[.13, .21]	
Mode parental substance abuse					<i>t</i> (169) = -0.29, <i>p</i> = .770
Interview	24	83	.15	[.12, .18]	
Questionnaire	27	88	.14	[.11, .18]	
Mode child well-being					<i>t</i> (174) = 0.02, <i>p</i> = .998
Interview	17	57	.14	[.11, .18]	
Questionnaire	26	119	.14	[.11, .17]	
Time lag association (months)	51	195			<i>t</i> (193) = -0.43, <i>p</i> = .667

Note. Some moderators were missing for certain studies. Each study can contribute multiple effect sizes, thus study sample size across subgroups can exceed the total study sample size for the Level 2 moderators. Within each moderator having more than two subgroups, identical letter superscripts indicate significant ($p < .05$) pairwise comparisons between subgroups.




Figure 2. Funnel plot for study-level mean effect sizes.

Oser, Knudsen, Staton-Tindall, & Leukefeld, 2009), and this suggests that both child well-being and parental substance abuse can be addressed through targeting aspects of the family environment.

Strengths and limitations

This study is the first large-scale review to broadly summarize the longitudinal relationship between parental substance abuse and child well-being. We excluded studies that sampled from specific groups whose membership would be expected to be associated with the child well-being outcome. As such, the reported association might be considered conservative, albeit more representative of the general population by doing so. There was no evidence of publication bias or an effect of methodological (rigor) features on the strength of the association, which further underpins the robustness of the findings. The multilevel approach to meta-analysis allowed us to include all relevant effect sizes, to model between- and within-study heterogeneity, and potential moderators that could explain variance at both levels—features that cannot be addressed with a standard meta-analytic approach. Although the multilevel approach is a powerful technique, some moderator analyses may have suffered from low power due to unbalanced groups. In particular, the subgroup effect of parental substance abuse in very young children was slightly elevated, while the subgroup effect for cognitive well-being was less pronounced. The limited number of studies on these topics could have reduced the power of the omnibus test to statistically detect group differences.

Parents with problematic patterns of using an intoxicating substance are likely to show variation in their use patterns over time, including tolerance and consequent increased consumption of substances to achieve the desired effect. They may also go through repeated withdrawal and relapse. At least for chronic use, there is more than simply the presence or the absence of parental substance abuse that potentially affects child well-being—features that this meta-analysis was unable to consider, as this was not reported in the literature reviewed. Furthermore, there was considerable variation in the way parental substance abuse was measured across studies. Some combined substance abuse into a

nebulous “any substance” category, some measured frequency, and others reported on recency of use. Given the investment in time and resources required by longitudinal studies, it is important that measurement is given careful consideration in future research and a range of metrics are taken to ensure that parental substance abuse can be appropriately captured. It is feasible that consumption heterogeneity in substances used (e.g., polysubstance abuse, frequency, dose, and physical dependence) would provide additional insights.

Although our results highlight that parental substance abuse is detrimental to children and adolescents, our findings are tempered by the limited number of studies that examined cognitive, social, or economic child well-being outcomes. Only three of the five dimensions of child well-being had sufficient data to be analyzed; therefore, the reported association should be interpreted as a deprecated form of child well-being rather than one that encompasses all aspects of well-being as we intended to quantify.

Within primary studies, well-being outcomes have likely been selected on a like-for-like basis such that parental substance abuse (e.g., tobacco) is expected to have implications for children’s health (e.g., respiratory health), and parental alcohol use is expected to have implications for children’s subsequent alcohol use, while educational attainment and other aspects of child well-being have been largely ignored. This is notable because the likely mechanisms—that parental substance abuse affects the parent-child relationship—suggests that a broader set of outcomes should be considered and that these gaps in the literature imply that child well-being is not sufficiently investigated to draw firm conclusions about all facets of well-being.

Moreover, as parental substance abuse measures were variously defined from “any substance abuse” to specific substances, there is potential variability in harm by substance type, other than alcohol and tobacco, which we were unable to capture due to a paucity of studies and poorly operationalized measures. Gender differences were also poorly operationalized, with many studies failing to separately capture father and mother differences between boys and girls. Just a third of the included studies considered mothers and fathers separately, only three studies differentiated between male and female children, and only a quarter considered children under 12 years of age. No study supplemented self-report data with objective outcomes, such as data from school or medical records. Therefore, future research needs to improve efforts to account for the multidimensionality of well-being, substance abuse, and features that could moderate effects, as it is possible that the effect of parental substance abuse is conditional on other factors that have not yet been formally documented. The generalizability of the results may also be limited due to a preponderance of studies emerging from North America. These biases and omissions in the research literature offer important opportunities to improve our understanding of the effects of parental substance abuse on child well-being.

Conclusion

This meta-analysis found that children exposed to parents who consume alcohol (both dependent and non-dependent), tobacco, or other psychoactive drugs experience a detrimental long-term effect on their well-being. However, there is a need for longitudinal studies that more accurately capture the effects of parental psychoactive substance abuse on the full scope of childhood well-being outcomes and integrate substance abuse into models that specify the precise conditions under which parental behavior

determines child well-being. These results can inform the development of family-oriented initiatives directed at improving child well-being that may also assist parental recovery.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0954579419000749>.

References

References marked with an asterisk indicate studies included in the meta-analysis.

- *Adalbjarnardottir, S., & Rafnsson, F. D. (2002). Adolescent antisocial behavior and substance use: Longitudinal analyses. *Addictive Behaviors*, 27, 227–240.
- Bandiera, F. C., Richardson, A. K., Lee, D. J., He, J.-P., & Merikangas, K. R. (2011). Secondhand smoke exposure and mental health among children and adolescents. *Archives of Pediatrics & Adolescent Medicine*, 165, 332–338.
- *Barman, S. K., Pulkkinen, L., Kaprio, J., & Rose, R. J. (2004). Inattentiveness, parental smoking and adolescent smoking initiation. *Addiction*, 99, 1049–1061.
- Barnard, M., & McKeganey, N. (2004). The impact of parental problem drug use on children: What is the problem and what can be done to help? *Addiction*, 99, 552–559.
- *Barnow, S., Schuckit, M., Smith, T., Spitzer, C., & Freyberger, H. J. (2007). Attention problems among children with a positive family history of alcohol abuse or dependence and controls—Prevalence and course for the period from preteen to early teen years. *European Addiction Research*, 13(1), 1–5.
- Beck, F., Guignard, R., & Legleye, S. (2014). Does computer survey technology improve reports on alcohol and illicit drug use in the general population? A comparison between two surveys with different data collection modes in France. *PloS One*, 9(1), e85810.
- Berger, L. M., Paxson, C., & Waldfogel, J. (2009). Income and child development. *Children and Youth Service Reviews*, 31, 978–989.
- *Bor, W., McGee, T. R., & Fagan, A. A. (2004). Early risk factors for adolescent antisocial behaviour: An Australian longitudinal study. *Australian and New Zealand Journal of Psychiatry*, 38, 365–372.
- Bountress, K., & Chassin, L. (2015). Risk for behavior problems in children of parents with substance use disorders. *American Journal of Orthopsychiatry*, 85, 275–286.
- Bradshaw, J. (2016). *The Well-being of Children in the UK*. Bristol: Policy Press.
- *Brook, J. S., Brook, D. W., Arcinibia-Mireles, O., Richter, L., & Whiteman, M. (2001). Risk factors for adolescent marijuana use across cultures and across time. *Journal of Genetic Psychology*, 162, 357–374.
- *Bufferd, S. J., Dougherty, L. R., Olino, T. M., Dyson, M. W., Laptook, R. S., Carlson, G. A., & Klein, D. N. (2014). Predictors of the onset of depression in young children: A multi-method, multi-informant longitudinal study from ages 3 to 6. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 55, 1279–1287.
- *Burke, V., Beilin, L. J., & Dunbar, D. (2001). Family lifestyle and parental body mass index as predictors of body mass index in Australian children: A longitudinal study. *International Journal of Obesity and Related Metabolic Disorders*, 25, 147–157.
- Calhoun, S., Conner, E., Miller, M., & Messina, N. (2015). Improving the outcomes of children affected by parental substance abuse: A review of randomized controlled trials. *Substance Abuse and Rehabilitation*, 6, 15–24.
- *Chaffin, M., Kelleher, K., & Hollenberg, J. (1996). Onset of physical abuse and neglect: Psychiatric, substance abuse, and social risk factors from prospective community data. *Child Abuse and Neglect*, 20, 191–203.
- Cheung, S. F., & Chan, D. K.-S. (2004). Dependent effect sizes in meta analysis: Incorporating the degree of interdependence. *Journal of Applied Psychology*, 89, 780–791.
- Cicero, T. J., Ellis, M. S., Surratt, H. L., & Kurtz, S. P. (2014). The changing face of heroin use in the United States: A retrospective analysis of the past 50 years. *JAMA Psychiatry*, 71, 821–826.
- *Colder, C. R., Chassin, L., Stice, E. M., & Curran, P. J. (1997). Alcohol expectancies as potential mediators of parent alcoholism effects on the development of adolescent heavy drinking. *Journal of Research on Adolescence*, 7, 349–374.
- Collins, L. M., & Horn, J. L. (Eds.) (1991). *Best methods for the analysis of change: Recent advances, unanswered questions, future directions*. Washington: American Psychological Association.
- *Connolly, G. M., Casswell, S., Stewart, J., & Silva, P. A. (1992). Drinking context and other influences on the drinking of 15-year-old New Zealanders. *British Journal of Addiction*, 87, 1029–1036.
- *Crandall, M., Chiu, B., & Sheehan, K. (2006). Injury in the first year of life: Risk factors and solutions for high-risk families. *Journal of Surgical Research*, 133(1), 7–10.
- *Cranford, J. A., Zucker, R. A., Jester, J. M., Puttler, L. I., & Fitzgerald, H. E. (2010). Parental alcohol involvement and adolescent alcohol expectancies predict alcohol involvement in male adolescents. *Psychology of Addictive Behaviors*, 24, 386–396.
- Cumming, G. (2014). The new statistics: Why and how. *Psychological Science*, 25, 7–29.
- De Goede, I. H., Branje, S. J., Delsing, M. J., & Meeus, W. H. (2009). Linkages over time between adolescents' relationships with parents and friends. *Journal of Youth and Adolescence*, 38, 1304–1315.
- De Los Reyes, A., Augenstein, T. M., Wang, M., Thomas, S. A., Drabick, D. A., Burgers, D. E., & Rabinowitz, J. (2015). The validity of the multi-informant approach to assessing child and adolescent mental health. *Psychological Bulletin*, 141, 858–900.
- Diener, E., Oishi, S., & Lucas, R. E. (2003). Personality, culture, and subjective well-being: Emotional and cognitive evaluations of life. *Annual Review of Psychology*, 54, 403–425.
- European Monitoring Centre for Drugs and Drug Addiction (2008). *Drugs and Vulnerable Groups of Young People*. Luxembourg: Office for Official Publications of the European Communities.
- *Dwyer, T., Ponsonby, A.-L., & Couper, D. (1999). Tobacco smoke exposure at one month of age and subsequent risk of SIDS: A prospective study. *American Journal of Epidemiology*, 149, 593–602.
- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ*, 315, 629–634.
- *Eiden, R. D., Edwards, E. P., & Leonard, K. E. (2007). A conceptual model for the development of externalizing behavior problems among kindergarten children of alcoholic families: Role of parenting and children's self-regulation. *Developmental Psychology*, 43, 1187–1201.
- *Ellickson, P. L., Tucker, J. S., Klein, D. J., & McGuigan, K. A. (2001). Prospective risk factors for alcohol misuse in late adolescence. *Journal of Studies on Alcohol*, 62, 773–782.
- Evans, G. W., Li, D., & Whipple, S. S. (2013). Cumulative risk and child development. *Psychological Bulletin*, 139, 1342–1396.
- National Center on Addiction and Substance Abuse (CASA). (2005). *Family matters: Substance abuse and the American family*. Retrieved from https://www.centeronaddiction.org/sites/default/files/Family-matters-substance-abuse-and-the-american-family_0.pdf.
- Faragher, E. B., Cass, M., & Cooper, C. L. (2005). The relationship between job satisfaction and health: A meta-analysis. *Occupational and Environmental Medicine*, 62, 105–112.
- Fearon, R., Bakermans-Kranenburg, M. J., Van Ijzendoorn, M. H., Lapsley, A. M., & Roisman, G. I. (2010). The significance of insecure attachment and disorganization in the development of children's externalizing behavior: A meta-analytic study. *Child Development*, 81, 435–456.
- *Fergusson, D. M., Horwood, L. J., & Lynskey, M. (1994). The childhoods of multiple problem adolescents: A 15-year longitudinal study. *Journal of Child Psychology and Psychiatry and Allied Disciplines*, 35, 1123–1140.
- *Fergusson, D. M., Horwood, L. J., & Lynskey, M. T. (1993). Maternal smoking before and after pregnancy: Effects on behavioral outcomes in middle childhood. *Pediatrics*, 92, 815–822.
- *Fergusson, D. M., Lynskey, M. T., & Horwood, L. J. (1996). Comorbidity between depressive disorders and nicotine dependence in a cohort of 16-year-olds. *Archives of General Psychiatry*, 53, 1043–1047.
- *Frank, D. A., Kuranz, S., Appugliese, D., Cabral, H., Chen, C., Crooks, D., ... Rose-Jacobs, R. (2014). Problematic substance use in urban adolescents: Role of intrauterine exposures to cocaine and marijuana and post-natal environment. *Drug and Alcohol Dependence*, 142, 181–190.

- Friedmann, P. D., Hendrickson, J. C., Gerstein, D. R., & Zhang, Z. (2004). The effect of matching comprehensive services to patients' needs on drug use improvement in addiction treatment. *Addiction*, 99, 962–972.
- Fu, R., Gartlehner, G., Grant, M., Shamliyan, T., Sedrakyan, A., Wilt, T. J., ... Trikalinos, T. A. (2011). Conducting quantitative synthesis when comparing medical interventions: AHRQ and the Effective Health Care Program. *Journal of Clinical Epidemiology*, 64, 1187–1197.
- *Fuller, B. E., Chermack, S. T., Cruise, K. A., Kirsch, E., Fitzgerald, H. E., & Zucker, R. A. (2003). Predictors of aggression across three generations among sons of alcoholics: Relationships involving grandparental and parental alcoholism, child aggression, marital aggression and parenting practices. *Journal of Studies on Alcohol*, 64, 472–483.
- Gallagher, M. W., Lopez, S. J., & Preacher, K. J. (2009). The hierarchical structure of well-being. *Journal of Personality*, 77, 1025–1050.
- Giesbrecht, N., Cukier, S., & Steeves, D. (2010). Collateral damage from alcohol: Implications of 'second-hand effects of drinking' for populations and health priorities. *Addiction*, 105, 1323–1325.
- *Griesler, P. C., Hu, M. C., Schaffran, C., & Kandel, D. B. (2011). Comorbid psychiatric disorders and nicotine dependence in adolescence. *Addiction*, 106, 1010–1020.
- *Gritz, E. R., Prokhorov, A. V., Hudmon, K. S., Jones, M. M., Rosenblum, C., Chang, C. C., ... de Moor, C. (2003). Predictors of susceptibility to smoking and ever smoking: A longitudinal study in a triethnic sample of adolescents. *Nicotine and Tobacco Research*, 5, 493–506.
- *Hanewinkel, R., Sargent, J. D., Hunt, K., Sweeting, H., Engels, R. C., Scholte, R. H., ... Morgenstern, M. (2014). Portrayal of alcohol consumption in movies and drinking initiation in low-risk adolescents. *Pediatrics*, 133, 973–982.
- *Heron, J., Barker, E. D., Joinson, C., Lewis, G., Hickman, M., Munafo, M., & Macleod, J. (2013). Childhood conduct disorder trajectories, prior risk factors and cannabis use at age 16: Birth cohort study. *Addiction*, 108, 2129–2138.
- Higgins, J. P. T., & Green, S. (Ed.) (2011). *Cochrane Handbook for Systematic Reviews of Interventions* (Vol. 4): Chichester, UK: Wiley.
- Hoeve, M., Stams, G. J. J., van der Put, C. E., Dubas, J. S., van der Laan, P. H., & Gerris, J. R. (2012). A meta-analysis of attachment to parents and delinquency. *Journal of Abnormal Child Psychology*, 40, 771–785.
- *Hoffmann, J. P., & Cerbone, F. G. (2002). Parental substance use disorder and the risk of adolescent drug abuse: An event history analysis. *Drug and Alcohol Dependence*, 66, 255–264.
- *Hussong, A. M., & Chassin, L. (1997). Substance use initiation among adolescent children of alcoholics: Testing protective factors. *Journal of Studies on Alcohol*, 58, 272–279.
- Hussong, A. M., Bauer, D. J., Huang, W., Chassin, L., Sher, K. J., & Zucker, R. A. (2008). Characterizing the life stressors of children of alcoholic parents. *Journal of Family Psychology*, 22, 819–832.
- Hussong, A. M., Huang, W., Curran, P. J., Chassin, L., & Zucker, R. A. (2010). Parent alcoholism impacts the severity and timing of children's externalizing symptoms. *Journal of Abnormal Child Psychology*, 38, 367–380.
- *Johnson, J. L., & Leff, M. (1999). Children of substance abusers: Overview of research findings. *Pediatrics*, 103(5 Pt 2), 1085–1099.
- *Kaplow, J. B., Curran, P. J., & Dodge, K. A. (2002). Child, parent, and peer predictors of early-onset substance use: A multisite longitudinal study. *Journal of Abnormal Child Psychology*, 30, 199–216.
- Keys, C. L. (2009). The nature and importance of positive mental health in America's adolescents. In R. Gilman, E. S. Huebner, & M. J. Furlong (Eds.), *Handbook of Positive Psychology in Schools* (pp. 9–23). New York: Routledge.
- *Knudsen, A. K., Ystrom, E., Skogen, J. C., & Torgersen, L. (2015). Maternal heavy alcohol use and toddler behavior problems: a fixed effects regression analysis. *European Child and Adolescent Psychiatry*, 24, 1269–1277.
- *Korhonen, T., Huizink, A. C., Dick, D. M., Pulkkinen, L., Rose, R. J., & Kaprio, J. (2008). Role of individual, peer and family factors in the use of cannabis and other illicit drugs: A longitudinal analysis among Finnish adolescent twins. *Drug and Alcohol Dependence*, 97, 33–43.
- *Kotch, J. B., Smith, J., Margolis, B., Black, M. M., English, D., Thompson, R., ... Bangdiwala, S. I. (2014). Does social capital protect against the adverse behavioral outcomes of child neglect? *Child Abuse Review*, 23, 246–261.
- Kraemer, H. C., Kazdin, A. E., Offord, D. R., Kessler, R. C., Jensen, P. S., & Kupfer, D. J. (1997). Coming to terms with the terms of risk. *Archives of General Psychiatry*, 54, 337–343.
- Kuppens, S., Grietens, H., Onghena, P., & Michiels, D. (2009a). A longitudinal study of childhood social behavior: Inter-informant agreement, inter-context agreement, and social preference linkages. *Journal of Social and Personal Relationships*, 26, 769–792.
- Kuppens, S., Grietens, H., Onghena, P., & Michiels, D. (2009b). Measuring parenting dimensions in middle childhood: Multitrait-multimethod analysis of child, mother, and father ratings. *European Journal of Psychological Assessment*, 25, 133–140.
- Lander, L., Howsare, J., & Byrne, M. (2013). The impact of substance use disorders on families and children: From theory to practice. *Social Work in Public Health*, 28, 194–205.
- *Lessard, J., Henrie, J., Livingston, J. A., Leonard, K. E., Colder, C. R., & Eiden, R. D. (2014). Correlates of ever having used electronic cigarettes among older adolescent children of alcoholic fathers. *Nicotine and Tobacco Research*, 16, 1656–1660.
- *Li, C., Pentz, M. A., & Chou, C. P. (2002). Parental substance use as a modifier of adolescent substance use risk. *Addiction*, 97, 1537–1550.
- *Lindsey, M. A., Browne, D. C., Thompson, R., Hawley, K. M., Graham, J., Weisbart, C., ... Kotch, J. B. (2008). Caregiver mental health, neighborhood, and social network influences on mental health needs among African American children. *Social Work Research*, 32, 79–88.
- Lipari, R. N. and Van Horn, S. L. (2017). *Children living with parents who have a substance use disorder. The CBHSQ Report: August 24, 2017*. Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration, Rockville, MD. Retrieved from https://www.samhsa.gov/data/sites/default/files/report_3223/ShortReport-3223.html
- Littell, R. C., Milliken, G. A., Stroup, W. W., Wolfinger, R. D., & Schabenberger, O. (2006). *SAS for Mixed Models* (2nd Ed.). Cary, NC: SAS Institute.
- *Loukas, A., Fitzgerald, H. E., Zucker, R. A., & von Eye, A. (2001). Parental alcoholism and co-occurring antisocial behavior: Prospective relationships to externalizing behavior problems in their young sons. *Journal of Abnormal Child Psychology*, 29, 91–106.
- Mallett, S., Rosenthal, D., & Keys, D. (2005). Young people, drug use and family conflict: Pathways into homelessness. *Journal of Adolescence*, 28, 185–199.
- *Malone, S. M., Iacono, W. G., & McGue, M. (2002). Drinks of the father: Father's maximum number of drinks consumed predicts externalizing disorders, substance use, and substance use disorders in preadolescent and adolescent offspring. *Alcoholism: Clinical and Experimental Research*, 26, 1823–1832.
- Martin, A., Razza, R. A., & Brooks-Gunn, J. (2012). Specifying the links between household chaos and preschool children's development. *Early Child Development and Care*, 182, 1247–1263.
- McGrath, C. E., Watson, A. L., & Chassin, L. (1999). Academic achievement in adolescent children of alcoholics. *Journal of Studies on Alcohol*, 60, 18–26.
- *McGrath, C. E., Watson, A. L., & Chassin, L. (1999). Academic achievement in adolescent children of alcoholics. *Journal of Studies on Alcohol*, 60, 18–26.
- *Miller, J. E. (2001). Predictors of asthma in young children: Does reporting source affect our conclusions? *American Journal of Epidemiology*, 154, 245–250.
- Mitchell, E. A., Beasley, R., Keil, U., Montefort, S., Odhiambo, J., & Group, I. P. T. S. (2012). The association between tobacco and the risk of asthma, rhinoconjunctivitis and eczema in children and adolescents: Analyses from Phase Three of the ISAAC programme. *Thorax*, 67, 941–949.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & The, P. G. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Medicine*, 6(7), e1000097.
- *Morgenstern, M., Sargent, J. D., Isensee, B., & Hanewinkel, R. (2013). From never to daily smoking in 30 months: The predictive value of tobacco and non-tobacco advertising exposure. *BMJ Open*, 3(6), 1–8.
- Morrison, A., Polisena, J., Husereau, D., Moulton, K., Clark, M., Fiander, M., ... Rabb, D. (2012). The effect of English-language restriction on systematic

- review-based meta-analyses: A systematic review of empirical studies. *International Journal of Technology Assessment in Health Care*, 28, 138–144.
- *Naicker, K., Wickham, M., & Colman, I. (2012). Timing of first exposure to maternal depression and adolescent emotional disorder in a national Canadian cohort. *PLoS ONE*, 7(3), e33422.
- Newman, D. B., Tay, L., & Diener, E. (2014). Leisure and subjective well-being: A model of psychological mechanisms as mediating factors. *Journal of Happiness Studies*, 15, 555–578.
- *Nigg, J. T., Wong, M. M., Martel, M. M., Jester, J. M., Puttler, L. I., Glass, J. M., ... Zucker, R. A. (2006). Poor response inhibition as a predictor of problem drinking and illicit drug use in adolescents at risk for alcoholism and other substance use disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45, 468–475.
- Öberg, M., Jaakkola, M. S., Woodward, A., Peruga, A., & Prüss-Ustün, A. (2011). Worldwide burden of disease from exposure to second-hand smoke: A retrospective analysis of data from 192 countries. *Lancet*, 377 (9760), 139–146.
- Oser, C., Knudsen, H., Staton-Tindall, M., & Leukefeld, C. (2009). The adoption of wraparound services among substance abuse treatment organizations serving criminal offenders: The role of a women-specific program. *Drug and Alcohol Dependence*, 103, S82–S90.
- *Pagani, L. S., Nguyen, A. K., & Fitzpatrick, C. (2016). Prospective associations between early long-term household tobacco smoke exposure and subsequent indicators of metabolic risk at age 10. *Nicotine and Tobacco Research*, 18, 1250–1257.
- Parsons, J. E., Adler, T. F., & Kaczala, C. M. (1982). Socialization of achievement attitudes and beliefs: Parental influences. *Child Development*, 53, 310–321.
- Peterson, A. V., Leroux, B. G., Bricker, J., Kealey, K. A., Marek, P. M., Sarason, I. G., & Andersen, M. R. (2006). Nine-year prediction of adolescent smoking by number of smoking parents. *Addictive Behaviors*, 31, 788–801.
- Podsakoff, P. M., MacKenzie, S. B., Lee, J.-Y., & Podsakoff, N. P. (2003). Common method biases in behavioral research: A critical review of the literature and recommended remedies. *Journal of Applied Psychology*, 88, 879–903.
- Pollard, E. L., & Lee, P. D. (2003). Child well-being: A systematic review of the literature. *Social Indicators Research*, 61, 5978.
- *Power, T. G., Stewart, C. D., Hughes, S. O., & Arbona, C. (2005). Predicting patterns of adolescent alcohol use: A longitudinal study. *Journal of Studies on Alcohol*, 66, 74–81.
- Puttler, L. I., Zucker, R. A., Fitzgerald, H. E., & Bingham, C. R. (1998). Behavioral outcomes among children of alcoholics during the early and middle childhood years: Familial subtype variations. *Alcoholism: Clinical and Experimental Research*, 22, 1962–1972.
- Raudenbush, S. W., & Bryk, A. S. (2002). *Hierarchical Linear Models: Applications and Data Analysis Methods* (2nd ed.). London: Sage.
- *Roosa, M. W., Beals, J., Sandler, I. N., & Pillow, D. R. (1990). The role of risk and protective factors in predicting symptomatology in adolescent self-identified children of alcoholic parents. *American Journal of Community Psychology*, 18, 725–741.
- Rosenthal, R. (1991). *Meta-Analytic Procedures for Social Research*. Newbury Park, CA: Sage.
- Rossow, I., Keating, P., Felix, L., & McCambridge, J. (2016). Does parental drinking influence children's drinking? A systematic review of prospective cohort studies. *Addiction*, 111, 204–217.
- Ryff, C. D., & Keyes, C. L. M. (1995). The structure of psychological well-being revisited. *Journal of Personality and Social Psychology*, 69, 719–727.
- Saulyte, J., Regueira, C., Montes-Martínez, A., Khudyakov, P., & Takkouche, B. (2014). Active or passive exposure to tobacco smoking and allergic rhinitis, allergic dermatitis, and food allergy in adults and children: A systematic review and meta-analysis. *PLoS Medicine*, 11(3), e1001611.
- *Selya, A. S., Dierker, L. C., Rose, J. S., Hedeker, D., & Mermelstein, R. J. (2012). Risk factors for adolescent smoking: Parental smoking and the mediating role of nicotine dependence. *Drug and Alcohol Dependence*, 124, 311–318.
- *Shortt, A. L., Hutchinson, D. M., Chapman, R., & Toumbourou, J. W. (2007). Family, school, peer and individual influences on early adolescent alcohol use: First-year impact of the Resilient Families programme. *Drug and Alcohol Review*, 26, 625–634.
- Siddaway, A. P., Taylor, P. J., & Wood, A. M. (2018). Re-conceptualizing anxiety as a continuum that ranges from high calmness to high anxiety: The joint importance of reducing distress and increasing well-being. *Journal of Personality and Social Psychology*, 114, e1–e11.
- Siddaway, A. P., Wood, A. M., & Taylor, P. J. (2017). The Centre for Epidemiologic Studies-Depression (CES-D) scale measures a continuum from well-being to depression: Testing two key predictions of Positive Clinical Psychology. *Journal of Affective Disorders*, 213, 180–186.
- Siddaway, A. P., Wood, A. M., & Hedges, L. V. (2019). How to do a systematic review: A best practice guide to conducting and reporting narrative reviews, meta-analyses, and meta-syntheses. *Annual Review of Psychology*, 70, 747–770.
- Størksen, I., Røysamb, E., Moum, T., & Tambs, K. (2005). Adolescents with a childhood experience of parental divorce: A longitudinal study of mental health and adjustment. *Journal of Adolescence*, 28, 725–739.
- Straussner, S. L. A. (1994). The impact of alcohol and other drug abuse on the American family. *Drug and Alcohol Review*, 13, 393–399.
- Straussner, S. L. A. (2004). Assessment and treatment of clients with alcohol and other drug abuse problems. In S. L. A. Straussner (Ed.), *Clinical Work with Substance-Abusing Clients* (pp. 3–36). New York: Guildford.
- Straussner, S. L. A., & Fewell, C. H. (2011). *Children of Substance-abusing Parents: Dynamics and Treatment*. New York: Springer Publishing Company.
- Stroup, D. F., Berlin, J. A., Morton, S. C., Olkin, I., Williamson, G. D., Rennie, D., ... Thacker, S. B. (2000). Meta-analysis of observational studies in epidemiology: A proposal for reporting. Meta-analysis of observational studies in epidemiology (MOOSE) group. *JAMA*, 283, 2008–2012.
- *Thacher, J. D., Gruzdeva, O., Pershagen, G., Neuman, A., van Hage, M., Wickman, M., ... Bergstrom, A. (2016). Parental smoking and development of allergic sensitization from birth to adolescence. *Allergy*, 71, 239–248.
- *Thompson, K. M., & Wilsnack, R. W. (1987). Parental influence on adolescent drinking: Modeling, attitudes, or conflict? *Youth and Society*, 19, 22–43.
- *Ting, T. T., Chen, W. J., Liu, C. Y., Lin, Y. C., & Chen, C. Y. (2015). Peer influences on alcohol expectancies in early adolescence: A study of concurrent and prospective predictors in Taiwan. *Addictive Behaviors*, 40, 7–15.
- *Tjora, T., Hetland, J., Aaro, L. E., & Overland, S. (2011). Distal and proximal family predictors of adolescents' smoking initiation and development: A longitudinal latent curve model analysis. *BMC Public Health*, 11, 911–920.
- Torgerson, C. J. (2006). Publication bias: the achilles' heel of systematic reviews? *British Journal of Educational Studies*, 54, 89–102.
- Tourangeau, R., & Yan, T. (2007). Sensitive questions in surveys. *Psychological Bulletin*, 133, 859–883.
- *Unger, J. B., Shakib, S., Gallahe, P., Ritt-Olson, A., Mouttapa, M., Palmer, P. H., & Johnson, C. A. (2006). Cultural/interpersonal values and smoking in an ethnically diverse sample of Southern California adolescents. *Journal of Cultural Diversity*, 13, 55–63.
- Van den Noortgate, W., Lopez-Lopez, J. A., Marin-Martinez, F., & Sanchez-Meca, J. (2013). Three-level meta-analysis of dependent effect sizes. *Behavioral Research Methods*, 45, 576–594.
- *Van Der Vorst, H., Krank, M., Engels, R. C., Pieters, S., Burk, W. J., & Mares, S. H. (2013). The mediating role of alcohol-related memory associations on the relation between perceived parental drinking and the onset of adolescents' alcohol use. *Addiction*, 108, 526–533.
- *Vella, S. A., Magee, C. A., & Cliff, D. P. (2015). Trajectories and predictors of health-related quality of life during childhood. *Journal of Pediatrics*, 167, 422–427.
- Vevea, J. L., & Woods, C. M. (2005). Publication bias in research synthesis: Sensitivity analysis using a priori weight functions. *Psychological Methods*, 10, 428–443.
- Wilson, D. B. (2017). *Comprehensive Meta-Analysis (Version 3)*. Englewood, NJ: Biostat.
- *Windham, A. M., Rosenberg, L., Fuddy, L., McFarlane, E., Sia, C., & Duggan, A. K. (2004). Risk of mother-reported child abuse in the first 3 years of life. *Child Abuse and Neglect*, 28, 645–667.
- *Wong, M. M., Brower, K. J., & Zucker, R. A. (2011). Sleep problems, suicidal ideation, and self-harm behaviors in adolescence. *Journal of Psychiatric Research*, 45, 505–511.
- Wong, W. C., Cheung, C. S., & Hart, G. J. (2008). Development of a quality assessment tool for systematic reviews of observational studies (QATSO)

of HIV prevalence in men having sex with men and associated risk behaviours. *Emerging Themes in Epidemiology*, 5, 23–26.

World Health Organization (2004). *Promoting mental health: Concepts, emerging evidence, practice*. Geneva: World Health Organization.

*Xie, B., Palmer, P., Li, Y., Lin, C., & Johnson, C. A. (2013). Developmental trajectories of cigarette use and associations with multilayered risk factors among Chinese adolescents. *Nicotine and Tobacco Research*, 15, 1673–1681.