

The effectiveness of aquatic plants as surrogates for wider biodiversity in standing fresh waters

Alan Law^{1*}, Ambroise Baker^{2, 3}, Carl Sayer², Garth Foster⁴, Iain D. M. Gunn⁵, Philip Taylor⁵, Zarah Pattison¹, James Blaikie¹, and Nigel J. Willby¹

¹ Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA, UK

² Pond Restoration Research Group, Environmental Change Research Centre, Department of Geography, University College London, London, WC1E 6BT, UK

³ School of Science, Engineering and Design, Teesside University, Middlesbrough, TS1 3BX, UK

⁴ Aquatic Coleoptera Conservation Trust, 3 Eglinton Terrace, Ayr, KA7 1JJ, UK

⁵ Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian, EH26 0QB, UK

* Corresponding author:

alan.law1@stir.ac.uk

Biological and Environmental Sciences, Cottrell building, University of Stirling, Stirling, FK9 4LA, UK.

Keywords: Lake, pond, macrophyte morpho-group, macroinvertebrates, structural equation modelling.

Summary

1. Fresh waters are among the most globally threatened habitats and their biodiversity is declining at an unparalleled rate. In an attempt to slow this decline, multiple approaches have been used to conserve, restore or enhance waterbodies. However, evaluating their effectiveness is time-consuming and expensive. Identifying species or assemblages across a range of ecological conditions that can provide a surrogate for wider freshwater biodiversity is therefore of significant value for conservation management and planning.
2. For lakes and ponds in three contrasting landscapes of Britain (lowland agricultural, eastern England; upland, north-west England; urban, central Scotland) we examined the link between macrophyte species, macrophyte morpho-group diversity (an indicator of structural diversity) and the richness of three widespread aquatic macroinvertebrate groups (molluscs, beetles and odonates) using structural equation modelling. We hypothesised that increased macrophyte richness and, hence, increased vegetation structural complexity, would increase macroinvertebrate richness after accounting for local and landscape conditions.
3. We found that macrophyte richness, via macrophyte morpho-group diversity, were an effective surrogate for mollusc, beetle and odonate richness in ponds after accounting for variation caused by physical variables, water chemistry and surrounding land use. However, only mollusc richness could be predicted by macrophyte morpho-group diversity in lakes, with no significant predicted effect on beetles or odonates.
4. Our results indicate that macrophyte morpho-group diversity can be viewed as a suitable surrogate of macroinvertebrate biodiversity across diverse landscapes, particularly in ponds and to a lesser extent in lakes. This has important implications for the restoration, conservation and creation of standing water habitats and for

49 assessing their effectiveness in addressing the decline of global freshwater
50 biodiversity. Management actions prioritising the development of species-rich and
51 structurally diverse macrophyte assemblages will likely benefit wider freshwater
52 biodiversity.

Introduction

A biological surrogate, indicator or proxy is an individual or group of organisms that can be used to identify a healthy, biodiverse or functional ecosystem, or to infer environmental conditions existing now or in the past. Such surrogates are commonly used in conservation decision making and offer a means of choosing and tracking the effectiveness of management approaches, with the premise that, if the surrogate is protected and conserved, there will be wider biodiversity and ecosystem benefits (Caro, 2010). A further advantage of surrogates is reduced reliance on large-scale, multi-taxon surveys which are time-consuming, expensive and often require specialist knowledge. Quantifying the link between surrogates and wider biodiversity or functioning of an ecosystem is crucial for validation, yet numerous studies conducted across several ecosystems and species have failed to identify consistent, reliable surrogates of either biodiversity, ecosystem function or phylogenetic diversity (Heino, 2015; Rapacciuolo *et al.*, 2018). Despite this, improved ecological knowledge and data accessibility, alongside advancing analytical tools, offer renewed promise in the search for surrogates. This is particularly relevant in freshwater ecosystems as they are one of the most globally threatened habitats due to the scale of humans impacts (Reid *et al.*, 2018; WWF, 2018).

Numerous studies have sought to evaluate surrogacy in freshwaters, with macroinvertebrates receiving most attention. For ponds and rivers there is broad consensus that a few species-rich invertebrate groups (e.g. Coleoptera, Odonata, Mollusca and Trichoptera) are broadly representative of wider macroinvertebrate assemblages (Briers & Biggs, 2003; Bilton *et al.*, 2006; Sánchez-Fernández *et al.*, 2006; Ruhí & Batzer, 2014; Guan *et al.*, 2018). However, where surrogacy across different taxonomic groups has been studied e.g. plants or amphibians to macroinvertebrates, the results have been inconsistent, with relationships variously non-existent (Santi *et al.*, 2010; Guareschi *et al.*, 2015), weak (Heino,

2010; Kirkman *et al.*, 2012; Rooney & Bayley, 2012; Ilg & Oertli, 2017), moderate (Santi *et al.*, 2010; Gioria *et al.*, 2010) or strong (Janssen *et al.*, 2018). Most previous research has concentrated on one or two taxonomic groups, focussing on a single habitat type, distributed over a small geographical range. Therefore, even at small-scales, there is limited evidence of effective surrogates for wider freshwater biodiversity.

Aquatic plants (macrophytes), encompassing bryophytes, macroalgae and vascular plants, are a fundamental component of aquatic food webs and play a central role in nutrient flux within freshwater habitats, linking atmosphere, soil and water. They influence the quality of the surrounding aquatic environment by creating structurally-complex habitats comprised of submerged, floating and emergent vegetation, where differences in leaf and stem architecture (e.g. floating vs. simple linear vs. dendritic leaves) between species, diversifies habitat complexity where it might otherwise be low (Jeppesen *et al.*, 1998). Furthermore, as primary producers, macrophytes influence water chemistry, provide food for grazers, habitats for egg-laying, whilst also mediating predator-prey interactions through provision of refugia for prey and concealment for predators (Diehl & Kornijow, 1998; Jeppesen *et al.*, 1998). A shared response to environmental conditions is often believed to be a key driver of species surrogacy (Gioria, Bacaro & Feehan, 2011; Rooney & Bayley, 2012), but, given the key structuring role of macrophytes, and their potential to operate as ecosystem engineers (Gurnell *et al.*, 2013), it seems highly likely that their presence and richness will directly or indirectly govern the availability of resources to, and environmental suitability for, other species. Since they are taxonomically and ecologically well understood and occur in almost all freshwater habitat types globally, macrophytes may thus be an ideal surrogate for wider freshwater biodiversity.

To our knowledge, the influence of macrophyte richness on multiple aquatic biota, across different freshwater habitats and covering environmentally diverse conditions has not

previously been examined. Therefore, current understanding of the potential value of macrophytes as a surrogate is constrained. In this study, aquatic molluscs, aquatic beetles and odonates were selected as focal biota due to their high taxonomic diversity, widespread distribution in standing fresh waters and because all three groups include species of conservation concern. Our primary objective was to test whether macrophytes act as surrogates for wider freshwater biodiversity across three contrasting (agricultural, upland and urban), but typical aquatic landscapes (so-called ‘hydrosapes’). We did this by assessing the strength of chemical and physical drivers and surrounding land use in explaining waterbody-scale richness of the biota. At the same time, we additionally tested if macrophyte species richness, mediated through morpho-group diversity, could further explain macroinvertebrate richness. We hypothesised that waterbodies with higher macrophyte richness, and, hence, greater macrophyte morpho-group diversity (an indicator of structural diversity), would have greater macroinvertebrate richness, with the former being a stronger predictor than chemical, physical and surrounding land use. However, further macroinvertebrate assemblage-specific effects are expected, reflecting either differences in the degree of dependence on macrophytes for habitat support, or habitat type-specific (pond or lake) differences in the importance of macrophytes as a component of habitat diversity.

Methods

Study areas and data collection

Three contrasting landscapes were chosen within Britain to account for different combinations of stressors associated with different land use types; lowland agricultural (north-east Norfolk, eastern England), upland (Cumbria, north-west England) and urban (Greater Glasgow, central Scotland). Within each of these hydrosapes, 22-29 replicates of both lakes and ponds were sampled. In this study, lakes were defined as waterbodies with

surface area > 1 ha, while ponds were < 1 ha in area and generally shallow (< 2 m max. depth). Both categories included man-made and natural waterbodies. Within each of these waterbodies four taxonomic groups were selected to cover a range of habitat requirements, pollutant sensitivities and dispersal abilities, namely macrophytes (as surrogates), aquatic molluscs, aquatic beetles (hereafter referred to as molluscs and beetles) and odonates (dragon and damselflies). Extensive data on these taxonomic groups were obtained via national recorders (i.e. Aquatic Coleoptera Conservation Trust, British Conchological Society and British Dragonfly Society), while water chemistry, where available, and data on macrophytes from commissioned surveys, was provided by UK environmental agencies or the Joint Nature Conservation Committee (JNCC). All data were closely scrutinised to ensure inter-compatibility, with multi-visit, full inventory surveys prioritised. Only records from the last decade were retained. The availability of multiple recent records of adult odonates influenced site selection because favourable weather conditions for surveying these could not be guaranteed during field campaigns conducted for this study. Where gaps in the data existed or when a greater number of replicate waterbodies were needed, new data were collected during June to August of 2016-17. Several sites had data collected for all species assemblages and 88% of the sites used in the study were visited by the authors to gather additional data for at least one species assemblage or to collect water samples for water chemistry analysis (Table S1).

For each waterbody, the following physical variables were derived from the UK Lakes Portal (<https://eip.ceh.ac.uk/apps/lakes/index.html>); altitude, area, catchment size, perimeter, ratio of waterbody to catchment area and shoreline development index (indicating shape complexity of the shoreline). For water chemistry data provided by UK environmental agencies, a mean value was taken for each variable based on samples collected in summer (June-September). In all other cases we collected a water sample from the middle of each site

and measured conductivity, dissolved oxygen, oxygen saturation, pH and temperature in the field using a HACH HQ30d meter. Alkalinity was also measured in the field by titration using sulphuric acid with a HACH AL-DT kit. A 500 ml subsample was filtered (47 mm glass microfiber, 1.2 μ m pore Whatman GF/C filters) within 12 hours of collection and analysed for major nutrients and metals (see Table S2 for a list of determinands). Chlorophyll *a* was determined by extraction by soaking filters in 90% methanol overnight and quantification by spectrophotometry.

For surveys of biota, exhaustive inventory sampling was conducted for each taxon group covering the complete margin of each waterbody. Macrophytes were recorded from the marginal zone to the maximum growing depth, assisted by use of a double-headed rake and/or a bathyscope for deeper water or where visibility was poor. For ponds, the entire water area was surveyed. For lakes, three or four sectors, each covering 100 m of shoreline, were surveyed to account for variation in exposure, shading, water depth and littoral substrate, following the JNCC survey methodology (Interagency Freshwater Group 2015). Within each sector, five transects were established perpendicular to the shore and four replicate quadrats were sampled per transect at depths of 0.25 m, 0.50 m, 0.75 m and >0.75 m, respectively, giving a total of 60 to 80 quadrats per lake. A boat was used to survey areas that were too deep for survey by wading (>75 cm).

Molluscs, beetles and larval odonates were sampled using a 1 mm mesh pond net. For each waterbody, the number of mesohabitats (e.g. rocky substrate, floating leaved, short/tall emergent, or submerged vegetation) was visually assessed and all were then sampled by sweeping the pond net through the water column and any vegetation present. This was repeated in each mesohabitat until no more new species could easily be found. The sample was live sorted and individuals were identified to species level in the field and released. When individuals could not be identified in the field they were preserved in 70% industrial

178 methylated spirits (IMS) and identified to species-level, wherever possible. Adult odonates
179 were identified visually in the field, assisted by use of binoculars. Where individuals within a
180 taxonomic group were identified to mixed resolution, only the highest resolution records
181 were used.

183 *Land cover and connectivity*

184 Land Cover Maps (Rowland *et al.*, 2017) were used to assess land use within the
185 upstream catchment of each waterbody (representing hydrological connectivity), and within
186 buffers of 50 m, 100 m, 500 m and 1 km surrounding each waterbody (representing riparian
187 and aerial connectivity). To reduce the number of interrelated land cover categories, a series
188 of composites were created; agricultural (arable and horticulture + improved grassland);
189 urban (suburban + urban) and wetland (fen, marsh and swamp + bog). Within each
190 waterbody buffer or catchment, land cover classes were expressed as a percentage of the total
191 buffer or catchment area (minus the area occupied by the focal waterbody). Since freshwater
192 and wetland land cover classes exhibited a high number of zero or low values these classes
193 were transformed to absence (-1) and presence (1) to make their effect sizes directly
194 comparable with those of continuous predictors.

196 *Variable selection and statistical analyses*

197 Species richness was defined as the number of macrophyte or macroinvertebrate
198 species per waterbody (or highest taxonomic resolution). Macrophyte morpho-group
199 diversity was derived by assigning each species to one of 26 morpho-groups based on a
200 library of morphological and regenerative traits (Willby, Abernethy & Demars, 2000), but
201 expanded to incorporate bryophytes, macroalgae and a wide range of emergent species (Table
202 S3). To determine if a sufficient number of waterbodies were surveyed per hydroscape for the

four taxonomic groups, sample coverage was calculated based on incidence data per waterbody using the iNEXT library (Hsieh, Ma & Chao, 2016). Prior to statistical analyses all continuous explanatory variables (excluding pH) were log transformed, mean centred and scaled by 1 SD, to improve comparability between variables and to reduce the effect of outliers (full set of continuous variables given in Figure S1). 64% of ponds sampled (especially those <0.1 ha) did not have definable catchments, so a binary ‘catchment present’ category was created for all ponds. Binary explanatory variables (e.g. catchment present for ponds, outflow and inflow) were transformed to have values of –1 (absent) and 1 (present).

To reduce model complexity principal components analysis (PCA) was applied to separate sets of water chemistry, physical and land use variables to identify those variables that maximised variation amongst sites (Figure S1). All continuous explanatory variables (excluding pH) were log transformed, mean centred and scaled by 1 SD, to improve comparability between variables and to reduce the effect of outliers. Correlations between predictor variables were then assessed in a correlation matrix (Figure S1) and checked for variance inflation (VIF). Where variables were highly correlative ($VIF > 20$) they were removed. The remaining variables were then used as explanatory variables for macrophyte species richness in a linear model (LM) with model-averaging then implemented (Burnham & Anderson, 2002). Variables that significantly explained macrophyte richness, based on the sums of Akaike weights (Figure S1), were then retained.

A conceptual model was developed to incorporate expected relationships between species richness and explanatory variables (Fig. 1). This model was based on the simple hypothesis that connectivity, land use and waterbody physical and water chemistry variables influence macrophyte species richness to a greater extent than macrophyte morpho-group diversity or richness of the macroinvertebrate groups, and that it is predominantly via macrophytes that these environmental effects are transmitted to macroinvertebrates. We also

hypothesised that macrophyte morpho-group diversity would be a more important determinant of macroinvertebrate richness than macrophyte taxonomic richness due to the increased structural complexity that a high richness of macrophyte morpho-groups provides. We used structural equation modelling (SEM) to quantify the direct and indirect effects of these explanatory variables on macrophyte richness, macrophyte morpho-group diversity and macroinvertebrate richness. SEMs are a multivariate technique based on constituent LMs that allow standardised comparisons of direct and indirect relationships. Constituent LMs were created and residuals assessed to determine if they met linear model assumptions and examined for spatial autocorrelation using Moran's I statistic. All constituent LMs met linear model assumptions and no significant patterns in spatial autocorrelation were detected ($P > 0.05$). Bivariate relationships between each response and explanatory variable were explored graphically to identify potential non-linear relationships. Where non-linear relationships were found, the explanatory variable was converted to second degree orthogonal polynomials. No multicollinearity was detected in constituent LMs with a VIF threshold of < 5 . During SEM model evaluation, missing pathways (i.e. previously unconsidered significant relationships) were identified and incorporated into the final SEM. Model fit was assessed using Fisher's C, where values of $P > 0.05$ indicated that the model was supported by the observed data. The term hydroscape ('Agricultural', 'Upland' and 'Urban') was added to each constituent LM, but was never significant and often increased the VIF due to correlations with land use. Hydroscape was then added as a random effect to each constituent LM, but did not improve the AIC. Therefore, the term hydroscape was not included in the final SEMs.

All statistical analysis was conducted using RStudio (R Core Team, 2018) with the libraries: piecewiseSEM (Lefcheck, 2016), sp (Bivand, Pebesma & Gomez-Rubio, 2013), sjPlot (Lüdtke, 2018), MuMIn (Bartoń, 2018), ggbiplot (Vu, 2011), factoextra (Kassambara

252 & Mundt, 2017), FactoMineR (Le, Josse & Husson, 2008), iNEXT (Hsieh *et al.*, 2016) and
253 spdep (Bivand, Hauke & Kossowski, 2015).

254

Results

The *a priori* designation of the three hydrosapes as upland, urban or agricultural was confirmed by analysis of the catchment characteristics of their constituent waterbodies (Table 1).

In total 176, 52, 249 and 35 species of macrophyte, mollusc, beetle and odonates respectively were recorded across the 158 waterbodies, studied via a combination of our surveys and archived data. Estimated sample coverage was generally high (mean = 94%) indicating effective sampling of each taxonomic group per waterbody type per hydroscape (Table 2). Further details of the sampling efficiency and completeness can be found in Figure S2 in the supporting information.

For both lakes and ponds, correlations in raw species richness was compared amongst the taxonomic groups (Figure S3), but none were found to be significant. Therefore, environment variables have to be considered in order to deduce true correlative relationships between the taxonomic groups.

Our conceptual model (Fig. 1) was poorly supported for both lakes and ponds, with multiple missing significant pathways being identified. However, with the addition of these pathways to the SEM (Table S4) the goodness-of-fit for both models reproduced the data well (lakes: Fisher's $C = 162.3$, $df = 164$, $P = 0.523$; ponds: Fisher's $C = 121.2$, $df = 124$, $P = 0.554$). Unstandardised and standardised effect sizes of all explanatory variables for lakes and ponds are provided in Table S5.

In lakes, macrophyte richness was explained principally by water chemistry and to a lesser extent by nearby land use ($R^2 = 0.64$) (Fig. 2). Variables indicative of nutrient-enrichment or poor water quality (nitrate, total phosphorus and water colour) negatively affected macrophyte richness, with nearby agricultural land positively influencing macrophyte richness. Macrophyte morpho-group diversity was, as expected, strongly related

to macrophyte richness. However, the subsequent effect on macroinvertebrates was varied; macrophyte morpho-group diversity positively influenced mollusc richness, but had no effect on beetle and odonate richness. For the latter groups, environmental conditions (i.e. land use and waterbody physical variables) were more influential. Increasing altitude was a strong, negative determinant of both mollusc and odonate richness, with reasonable variance explained for both assemblages ($R^2 = 0.76$ and 0.36). The explained variance in beetle richness was the lowest of all the taxonomic groups ($R^2 = 0.29$) with only wetlands in the catchment and nearby agricultural land positively affecting richness and, to a lesser extent, lakes with relatively large catchments having a negative effect.

For ponds, nearby surrounding land use had no significant impact on macrophyte richness compared to the influence of water chemistry (principally conductivity and pH) and presence of an outflow (Fig. 3). Macrophyte morpho-group diversity was again strongly related to macrophyte richness, whilst ammonium and nearby urban land use also had minor negative effects on morpho-group diversity. The degree of urbanisation within 500 m of a pond had contrasting effects on macroinvertebrate biota, being positive for molluscs, but highly negative for beetles and odonates. A negative effect of altitude was observed for mollusc and beetle richness in ponds, as with lakes. Nevertheless, despite some variation being explained by physical variables, water chemistry and land use, an increased macrophyte morpho-group diversity had a significant positive effect on all macroinvertebrate groups.

Discussion

Simple surrogates for freshwater biodiversity should help to inform choices over the protection, restoration or creation of waterbodies, and in monitoring the effectiveness of related actions. However, few studies have sought out a surrogate appropriate for multiple freshwater habitats and disparate species assemblages over large spatial scales. We found that, regardless of the landscape, high macrophyte richness, specifically via high morpho-group diversity, was a suitable surrogate for a higher richness of multiple macroinvertebrate species assemblages (molluscs, beetles and odonates) in ponds, but only mollusc richness could be predicted by macrophyte morpho-group diversity in lakes.

The drivers of species richness

Land use is often assumed to be a major driver of species composition as it provides a proxy for stressors (e.g. agriculturally-derived nutrients or pollutants originating from urban areas) (Hassall, 2014) or affects spatial processes (altering connectivity both positively and negatively) (Hill *et al.*, 2017). Urbanisation is assumed to be indicative of reduced connectivity due to the density of roads and built-up areas that restrict dispersal between waterbodies (Hassall, 2014). Moreover, previous studies of ponds and rivers indicate that active dispersers were less restricted by habitat structure than passive dispersers (Hill *et al.*, 2017; Sarremejane *et al.*, 2017). In our study, urban land use had a negative effect on actively-dispersing odonates and beetles in ponds, suggesting that an active dispersal ability may be insufficient to counteract effects of urbanisation and the associated changes to local habitat structure that urbanisation produces. However, urban land use was positively associated with passively dispersing molluscs. This latter finding may reflect the increased presence of vectors within the local landscape (for example waterfowl attracted by supplementary feeding may increase bird-mediated dispersal (van Leeuwen *et al.*, 2012;

Simonová *et al.*, 2016)), combined with molluscs' tolerance of productive poorly oxygenated conditions. Alternatively, the increased concentrations of some major ions due to rural and urban run-off may also benefit molluscs since calcium is used for shell construction (Moss, 2017). It was expected that adjacent agricultural land use would negatively affect biodiversity due to increased nutrient or fine sediment inputs, yet agriculture within 500 m of lakes had a slight positive effect on lake macrophyte richness. However, the interpretation that agriculture is positive for biodiversity should be taken with caution, since in the composite LMs that underpin the SEM, agricultural land use in the catchment as a whole had a non-linear relationship with macrophyte richness, becoming negative when agricultural extent exceeded ~40% (though this was not significant in the final model). Freshwaters and wetlands in the catchment or buffers were expected to positively affect biodiversity as they potentially increase connectivity, and therefore resilience, by acting as stepping stones (Biggs *et al.*, 2005). Although we observed a positive effect of nearby wetlands (within a 500 m buffer), or wetlands in the catchment on lake beetles and molluscs, respectively, this was secondary to waterbody-specific influences (e.g. altitude and water chemistry), consistent with other studies (Hill *et al.*, 2017; Thornhill *et al.*, 2017). Water chemistry influenced macrophyte richness in both lakes and ponds, with variables indicative of nutrient-enrichment negatively affecting richness. Alkalinity had a negative effect on lake macrophytes, which was unexpected as previous work has generally shown a positive influence of alkalinity on macrophyte richness (Vestergaard & Sand-Jensen, 2000). The effect we observed was most likely driven by a strong correlation between alkalinity and total oxidised nitrogen or conductivity (Figure S1), indicative of declining water quality (Heegaard *et al.*, 2001). Waterbody chemistry had few direct effects on the studied macroinvertebrate groups and it is therefore likely that macrophytes mediate nutrient-enrichment effects (Declerck *et al.*, 2005).

Identifying a simple surrogate of diverse and complex species assemblages that transcends multiple, potentially interacting variables which vary both temporally and spatially is difficult, with few variables seemingly transferable across habitat types, regions and species assemblages (Batzler, 2013). Macrophyte richness and composition have previously been shown to positively affect macroinvertebrate assemblages in multiple freshwater habitats; ponds (Palmer, 1981; Gioria *et al.*, 2011), wetlands (Kirkman *et al.*, 2012), lakes (Heino & Tolonen, 2017) and rivers (Holmes & Raven, 2014). However, the drivers of species surrogacy are mostly speculative rather than explicitly studied. In our study, the most plausible basis for the surrogacy we observed is that good water quality allows for high macrophyte richness, which leads to a greater diversity of macrophyte morpho-groups and macroinvertebrate richness benefits through provision of increased architectural complexity. These benefits are probably group- or life stage-specific. For example, molluscs may benefit from high macrophyte richness due to increased food resources, reduced predation and increased microhabitat diversity (Brönmark, 1985). Beetles may benefit from the heterogeneous substrate available for egg-laying, refugia and through increased prey availability (Bloechl *et al.*, 2010). Furthermore, adult odonates use emergent macrophytes for perching, egg-laying and emergence (Le Gall *et al.*, 2018), whereas their larvae use submerged macrophytes for shelter and foraging (Goertzen & Suhling, 2013). A greater macrophyte morpho-group richness linked to asynchronous growth peaks may also extend the duration of macrophyte cover (van Donk & Gulati, 1995; Sayer, Davidson & Jones, 2010) which should benefit macroinvertebrates, but this area is relatively unexplored.

It is also possible that some macroinvertebrate groups may influence the richness of others, for example, via predation. However, as positive or negative pathways between any of the macroinvertebrate groups were not identified in our analysis, we can hypothesize that the effect of predation on richness are low, relative to the effect of macrophytes. Differences in

explained variance amongst macroinvertebrates were reasonably consistent across waterbody types, with mollusc richness highest followed by odonates and then beetles. The low explained variance observed for beetles may in part reflect the high species richness found. Beetles are one of the most speciose groups globally with a wide geographical and ecological range (Bilton *et al.*, 2006); moreover, the balance between habitat specialists and generalists will be masked when considering diversity only in terms of species richness.

The strength of the surrogacy between macrophytes and macroinvertebrates differed between waterbody types, with macrophyte richness being a stronger driver of macroinvertebrate richness in ponds than lakes. This pattern may arise because lakes are more likely to support large populations of fish, which are known to exert strong predation pressure on macroinvertebrates (Diehl, 1992; Jones & Sayer, 2003). Molluscs, for example, are commonly consumed by fish with resulting reductions in density, although effects on richness are less understood (Dillon, 2000). Fish could also influence macroinvertebrates indirectly via various cascading effects on macrophyte diversity caused by herbivory (Matsuzaki *et al.*, 2009), zooplanktivory (Jeppesen *et al.*, 1998) or benthivory, particularly in shallow lakes (Kloskowski, 2011). Both abundance of macrophytes and macroinvertebrates will also be affected by waterfowl herbivory and bioturbation (Rodríguez-Pérez & Green, 2012; Wood *et al.*, 2012), with lakes likely to support greater waterfowl densities than ponds. A further factor affecting macroinvertebrate diversity in lakes may be physical disturbance of the shoreline due to wave action, which is much more intense in lakes than ponds due to an increased fetch (Fairchild, Faulds & Matta, 2000). Given that our focal macroinvertebrate groups, molluscs in particular, are poorly stream-lined and prone to being dislodged by currents, their link with macrophyte diversity may reflect a shared need for sheltered marginal habitats. In this study, it is likely that the effects of fish predation or physical disturbance on macroinvertebrate richness is mediated through macrophyte morpho-group

diversity, as found in Cladocera (Burks, Jeppesen & Lodge, 2007), but further study would be useful to tease apart the multiple interacting processes involved (see Dillon (2000) for a review). Moreover, future studies should endeavour to determine fish abundance. As fish can be important drivers of aquatic community composition (Scheffer *et al.*, 2006), their inclusion will undoubtedly improve the predictive power of models and therefore the application of surrogates in other freshwater habitats.

Surrogacy and available statistical tools

The search for widely applicable and robust surrogates of freshwater biodiversity has probably been somewhat confounded by the differing statistical approaches used to detect surrogacy (Gioria *et al.*, 2011). The majority of studies have tested congruence between species assemblages by using multivariate ordination to consider the influence of local environmental variables (Declerck *et al.*, 2005; Bilton *et al.*, 2006; Santi *et al.*, 2010; Gioria *et al.*, 2010; Guareschi *et al.*, 2015). Others have utilised Mantel tests (Heino, 2010; Rooney & Bayley, 2012; Ruhí & Batzer, 2014; Ilg & Oertli, 2017), species correlations (Sánchez-Fernández *et al.*, 2006; Slimani *et al.*, 2019) or a Species Accumulation Index (Kirkman *et al.*, 2012). In addition to the range of analytical methods used, the choice of diversity index for assessing surrogacy also influences outcomes, with alternative measures of alpha diversity (e.g. richness, functional and phylogenetic alpha) varying in their sensitivity to environmental drivers (Heino & Tolonen, 2017). To our knowledge SEMs have not been previously utilised in the quest for surrogacy in freshwater ecology. The advantage of SEMs is that disparate species assemblages can be analysed in relation to environmental variables, unlike most community analyses that can only directly compare two assemblages at a time. Moreover, SEMs standardise across environmental variables without the need for multiple

tests that risk false positives, and can, therefore, elucidate the relative strengths of explanatory variables in driving observed relationships.

Applications

An effective surrogate should be transferable over a broad context and offer a currency that is understandable to a range of stakeholders. According to our findings, macrophytes could meet these criteria in providing an indirect surrogate for molluscs, beetles and dragonflies in ponds and for molluscs in lakes. Macrophyte richness as a freshwater biodiversity surrogate could be applicable from local to landscape scales, and simplify complex patterns and processes. By isolating the effects of multiple environmental and spatial explanatory variables in our dataset we demonstrate statistically that, via the diversity of morpho-group diversity, a greater richness of macrophytes is also broadly indicative of greater richness across disparate macroinvertebrate groups in ponds and molluscs in lakes. From an applied perspective, as macrophytes act as ecosystem architects, our findings suggest that researchers or practitioners can straightforwardly obtain a broad indication of the overall habitat quality and macroinvertebrate biodiversity by monitoring the number of macrophyte species and diversity of macrophyte morpho-groups, especially in the case of ponds. Despite the advantages of surrogates, they cannot replace detailed surveys of taxonomic groups particularly where species are rare, specialists or of conservation interest. Therefore, although our results show that macrophyte morpho-group diversity can be useful to indicate freshwater biodiversity, some caution is required as these results may not be definitive in the broad sense.

It has been argued that declines in macrophyte richness should be viewed as an early warning system for declines in overall macrophyte abundance and hence the quality of the wider environment (Sayer *et al.*, 2010). Hence, we would recommend practitioners and

conservation managers need to be concerned for wider biodiversity if macrophyte richness begins to decrease. The use of macrophytes as freshwater biodiversity surrogates can be important for rapid and cost-effective assessment of conservation and restoration projects, however, they will be most effective where constraints to biodiversity are diagnosed and addressed at site, habitat and landscape-scales. For example, at the site-scale, high grazing pressures may limit macrophyte regeneration from seedbanks and therefore wider biodiversity will only benefit if areas of macrophytes are protected from over-grazing and high disturbance. Additionally, at the habitat or landscape-scale, species translocations may be needed to enhance structural complexity if there are significant barriers to colonisation. However, in using macrophytes as a proxy for wider biodiversity, particularly when assessing habitat restoration, it should be recognised that macrophyte responses to management are complex and can be highly variable (Phillips, Willby & Moss, 2016).

Acknowledgments

This research was financially supported by the Natural Environmental Research Council Highlight Topics grant NE/N006437/1: Hydroscape - connectivity x stressor interaction in freshwater habitats. We also thank our many data providers (Aquatic Coleoptera Conservation Trust, Biological Records Centre, British Conchological Society, British Dragonfly Society, Broads Authority, Buglife, Cumbria Wildlife Trust, ENSIS, Environment Agency, Froglife, Glasgow City Council, Norfolk Biodiversity Information Service, RSPB, Scottish Environment Protection Agency and Upland Waters Monitoring Network) and pay tribute to the numerous volunteers who contribute their data to recording schemes and to the many landowners for allowing access to their sites. We are also grateful to two anonymous reviewers for providing helpful comments on this manuscript.

475

476 **Data Availability Statement**

477 All data referred to in this article and code used in analyses are deposited in DataSTORRE -
478 the University of Stirling research data repository.

479

480 **Conflict of Interest Statement**

481 The authors declare that they have no conflicts of interest in presenting this work for
482 publication.

483

References

- Bartoń K. (2018). *MuMIn: Multi-Model Inference*. R package version 1.42.1. Available at: <https://CRAN.R-project.org/package=MuMIn>
- Batzer D.P. (2013). The seemingly intractable ecological responses of invertebrates in North American Wetlands: A review. *Wetlands*, 33, 1–15. <https://doi.org/10.1007/s13157-012-0360-2>
- Biggs J., Williams P., Whitfield M., Nicolet P. & Weatherby A. (2005). 15 years of pond assessment in Britain: results and lessons learned from the work of Pond Conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 15, 693–714. <https://doi.org/10.1002/aqc.745>
- Bilton D.T., Mcabendroth L., Bedford A. & Ramsay P.M. (2006). How wide to cast the net? Cross-taxon congruence of species richness, community similarity and indicator taxa in ponds. *Freshwater Biology*, 51, 578–590. <https://doi.org/10.1111/j.1365-2427.2006.01505.x>
- Bivand R.S., Hauke J. & Kossowski T. (2015). Comparing Implementations of Estimation Methods for Spatial Econometrics. *Journal of Statistical Software*, 63, 1–36
- Bivand R.S., Pebesma E. & Gomez-Rubio V. (2013). Applied spatial data analysis with R (2nd edition). New York: Springer
- Bloechl A., Koenemann S., Philippi B. & Melber A. (2010). Abundance, diversity and succession of aquatic Coleoptera and Heteroptera in a cluster of artificial ponds in the North German Lowlands. *Limnologica - Ecology and Management of Inland Waters*, 40, 215–225. <https://doi.org/10.1016/j.limno.2009.08.001>

506 Briers R.A. & Biggs J. (2003). Indicator taxa for the conservation of pond invertebrate
 507 diversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 13, 323–330.
 508 <https://doi.org/10.1002/aqc.576>
 509 Brönmark C. (1985). Freshwater snail diversity: effects of pond area, habitat heterogeneity
 510 and isolation. *Oecologia*, 67, 127–131. <https://doi.org/10.1007/BF00378463>
 511 Burks R. L., Jeppesen E., & Lodge D. M. (2001). Littoral zone structures as *Daphnia* refugia
 512 against fish predators. *Limnology and Oceanography*, 46, 230-237.
 513 Burnham K.P. & Anderson D.R. (2002). Model selection and multi-model inference: a
 514 practical information-theoretic approach. New York: Springer
 515 Caro T. (2010). Conservation by proxy: indicator, umbrella, keystone, flagship, and other
 516 surrogate species. Island Press
 517 Declerck S., Vandekerkhove J., Johansson L., Muylaert K., Conde-Porcuna M., Van Der Gucht
 518 K., et al. (2005). Multi-group biodiversity in shallow lakes along gradients of
 519 phosphorus and water plant cover. *Ecology*, 86, 1905–1915
 520 Diehl S. (1992). Fish Predation and Benthic Community Structure - the Role of Omnivory and
 521 Habitat Complexity. *Ecology*, 73, 1646–1661. <https://doi.org/10.2307/1940017>
 522 Diehl S. & Kornijow R. (1998). The influence of submerged macrophytes on trophic
 523 interactions among fish and macroinvertebrates. In: The Structuring Role of
 524 Submerged Macrophytes in Lakes. (Eds E. Jeppesen, M. Søndergaard, M.
 525 Søndergaard & K. Christoffersen), pp. 24–46. New York: Springer.
 526 Dillon R.T. (2000). The ecology of freshwater molluscs. Cambridge University Press
 527 Fairchild G.W., Faulds A.M. & Matta J.F. (2000). Beetle assemblages in ponds: effects of
 528 habitat and site age. *Freshwater Biology*, 44, 523–534.
 529 <https://doi.org/10.1046/j.1365-2427.2000.00601.x>

530 Le Gall M., Fournier M., Chaput-Bardy A. & Husté A. (2018). Determinant landscape-scale
 531 factors on pond odonate assemblages. *Freshwater Biology*, 63, 306-317.
 532 <https://doi.org/10.1111/fwb.13065>

533 Gioria M., Bacaro G. & Feehan J. (2011). Evaluating and interpreting cross-taxon
 534 congruence: Potential pitfalls and solutions. *Acta Oecologica*, 37, 187–194.
 535 <https://doi.org/10.1016/j.actao.2011.02.001>

536 Gioria M., Schaffers A., Bacaro G. & Feehan J. (2010). The conservation value of farmland
 537 ponds: Predicting water beetle assemblages using vascular plants as a surrogate
 538 group. *Biological Conservation*, 143, 1125–1133.
 539 <https://doi.org/10.1016/j.biocon.2010.02.007>

540 Goertzen D. & Suhling F. (2013). Promoting dragonfly diversity in cities: Major determinants
 541 and implications for urban pond design. *Journal of Insect Conservation*, 17, 399–409.
 542 <https://doi.org/10.1007/s10841-012-9522-z>

543 Guan Q, Liu J., Batzer D.P., Lu X. & Wu H. (2018). Snails (Mollusca: Gastropoda) as potential
 544 surrogates of overall aquatic invertebrate assemblage in wetlands of Northeastern
 545 China. *Ecological Indicators*, 90, 193-
 546 200. <https://doi.org/10.1016/j.ecolind.2018.01.069>

547 Guareschi S., Abellan P., Laini A., Green A. J., Sanchez-Zapata J. A., Velasco J., & Millan A.
 548 (2015). Cross-taxon congruence in wetlands: Assessing the value of waterbirds as
 549 surrogates of macroinvertebrate biodiversity in Mediterranean Ramsar sites.
 550 *Ecological Indicators*, 49, 204-215.

551 Gurnell A.M., O'Hare M.T., O'Hare J.M., Scarlett P. & Liffen T.M.R. (2013). The
 552 geomorphological context and impact of the linear emergent macrophyte,
 553 *Sparganium erectum* L.: A statistical analysis of observations from British rivers.

554 *Earth Surface Processes and Landforms*, 38, 1869–1880.

555 <https://doi.org/10.1002/esp.3473>

556 Hassall C. (2014). The ecology and biodiversity of urban ponds. *Wiley Interdisciplinary*

557 *Reviews: Water* 1, 187–206. <https://doi.org/10.1002/wat2.1014>

558 Heegaard E., Birks H.H., Gibson C.E., Smith S.J. & Wolfe-Murphy S. (2001). Species-

559 environmental relationships of aquatic macrophytes in Northern Ireland. *Aquatic*

560 *Botany*, 70, 175–223. [https://doi.org/10.1016/S0304-3770\(01\)00161-9](https://doi.org/10.1016/S0304-3770(01)00161-9)

561 Heino J. (2015). Approaches, potential and pitfalls of applying bioindicators in freshwater

562 ecosystems. In: Indicators and Surrogates of Biodiversity and Environmental Change.

563 (Eds D. Lindenmayer, P. Barton & J. Pierson), pp. 91–100. CRC Press.

564 Heino J. (2010). Are indicator groups and cross-taxon congruence useful for predicting

565 biodiversity in aquatic ecosystems? *Ecological Indicators*, 10, 112–117.

566 <https://doi.org/10.1016/j.ecolind.2009.04.013>

567 Heino J. & Tolonen K.T. (2017). Untangling the assembly of littoral macroinvertebrate

568 communities through measures of functional and phylogenetic alpha diversity.

569 *Freshwater Biology*, 62, 1168–1179. <https://doi.org/10.1111/fwb.12934>

570 Hill M.J., Heino J., Thornhill I., Ryves D.B. & Wood P.J. (2017). Effects of dispersal mode on

571 the environmental and spatial correlates of nestedness and species turnover in pond

572 communities. *Oikos* 126, 1575–1585. <https://doi.org/10.1111/oik.04266>

573 Holmes N. & Raven P. (2014). Rivers: A natural and not-so-natural history. British Wildlife

574 Publishing.

575 Hsieh T.C., Ma K.H. & Chao A. (2016). iNEXT: An R package for rarefaction and extrapolation

576 of species diversity (Hill numbers). *Methods in Ecology and Evolution*, 7, 1451–1456.

577 <https://doi.org/10.1111/2041-210X.12613>

578 Ilg C. & Oertli B. (2017). Effectiveness of amphibians as biodiversity surrogates in pond
 579 conservation. *Conservation Biology*, 31, 437–445.
 580 <https://doi.org/10.1111/cobi.12802>

581 Interagency Freshwater Group. (2015). Common Standards Monitoring Guidance for
 582 Freshwater Lakes. Joint Nature Conservation Committee

583 Janssen A., Hunger H., Konold W., Pufal G. & Staab M. (2018). Simple pond restoration
 584 measures increase dragonfly (Insecta: Odonata) diversity. *Biodiversity and
 585 Conservation*, 27, 2311–2328. <https://doi.org/10.1007/s10531-018-1539-5>

586 Jeppesen E., Søndergaard M., Søndergaard M. & Christoffersen K. (1998). The Structuring
 587 Role of Submerged Macrophytes in Lakes. New York: Springer

588 Jones, J.I. & Sayer, C. D. (2003). Does the fish-invertebrate-periphyton cascade precipitate
 589 plant loss in shallow lakes? *Ecology*, 84, 2155-2167. <https://doi.org/10.1890/02-0422>

590 Kassambara A. & Mundt F. (2017). factoextra: extract and visualize the results of
 591 multivariate data analyses. R package version 1.0.5. Available at: [https://CRAN.R-
 592 project.org/package=factoextra](https://CRAN.R-project.org/package=factoextra)

593 Kirkman L.K., Smith L.L., Quintana-Ascencio P.F., Kaeser M.J., Golladay S.W. & Farmer A.L.
 594 (2012). Is species richness congruent among taxa? Surrogacy, complementarity, and
 595 environmental correlates among three disparate taxa in geographically isolated
 596 wetlands. *Ecological Indicators*, 18, 131–139.
 597 <https://doi.org/10.1016/j.ecolind.2011.10.015>

598 Kloskowski J. (2011). Impact of common carp *Cyprinus carpio* on aquatic communities:
 599 direct trophic effects versus habitat deterioration. *Fundamental and Applied
 600 Limnology*, 178, 245–255. <https://doi.org/10.1127/1863-9135/2011/0178-0245>

601 Le S., Josse J. & Husson F. (2008). FactoMineR: An R package for multivariate analysis.
602 *Journal for Statistical Software*, 25, 1–18

603 Lefcheck J.S. (2016). piecewiseSEM: Piecewise structural equation modeling in R for ecology,
604 evolution, and systematics. *Methods in Ecology and Evolution*, 7, 573–579

605 Lüdtke D. (2018). sjPlot: Data Visualization for Statistics in Social Science. R package
606 version 2.6.0. Available at: <https://CRAN.R-project.org/package=sjPlot>

607 Matsuzaki S.S., Usio N., Takamura N. & Washitani I. (2009). Contrasting impacts of invasive
608 engineers on freshwater ecosystems: an experiment and meta-analysis. *Oecologia*,
609 158, 673–686.

610 Moss B. (2017). Ponds and small lakes. Pelagic Publishing.

611 Palmer M. (1981). Relationship between species richness of macrophytes and insects in
612 some water bodies in the Norfolk Breckland. *Entomologists Monthly Magazine*, 117,
613 35–46

614 Phillips G., Willby N.J. & Moss B. (2016). Submerged macrophyte decline in shallow lakes:
615 What have we learnt in the last forty years? *Aquatic Botany*, 135, 37–45.
616 <https://doi.org/10.1016/j.aquabot.2016.04.004>

617 R Core Team (2018). R: A language and environment for statistical computing. R Foundation
618 for Statistical Computing, Vienna

619 Rapacciuolo G., Graham C.H., Marin J., Behm J.E., Costa G.C., Hedges S.B., et al. (2018).
620 Species diversity as a surrogate for conservation of phylogenetic and functional
621 diversity in terrestrial vertebrates across the Americas. *Nature Ecology & Evolution*,
622 3, 53–61. <https://doi.org/10.1038/s41559-018-0744-7>

623 Reid A.J., Carlson A.K., Creed I.F., Eliason E.J., Gell P.A., Johnson P.T.J., et al. (2018).
 624 Emerging threats and persistent conservation challenges for freshwater biodiversity.
 625 *Biological Reviews*, 94, 849-873. <https://doi.org/10.1111/brv.12480>
 626 Rodríguez-Pérez H., & Green A. J. (2012). Strong seasonal effects of waterbirds on benthic
 627 communities in shallow lakes. *Freshwater Science*, 31, 1273-1288.
 628 Rooney R. C., & Bayley, S. E. (2012). Community congruence of plants, invertebrates and
 629 birds in natural and constructed shallow open-water wetlands: Do we need to
 630 monitor multiple assemblages? *Ecological Indicators*, 20, 42-50.
 631 Rowland C.S., Morton R.D., Carrasco L., McShane G., O'Neil A.W. & Wood C.M. (2017). Land
 632 Cover Map 2015 (vector, GB)
 633 Ruhí A. & Batzer D.P. (2014). Assessing Congruence and Surrogacy Among Wetland
 634 Macroinvertebrate Taxa Towards Efficiently Measuring Biodiversity. *Wetlands*, 34,
 635 1061–1071. <https://doi.org/10.1007/s13157-014-0566-6>
 636 Sánchez-Fernández D., Abellán P., Mellado A., Velasco J. & Millán A. (2006). Are water
 637 beetles good indicators of biodiversity in mediterranean aquatic ecosystems? The
 638 case of the Segura River Basin (SE Spain). *Biodiversity and Conservation*, 15, 4507.
 639 <https://doi.org/10.1007/s10531-005-5101-x>
 640 Santi E., Mari E., Piazzini S., Renzi M., Bacaro G. & Maccherini S. (2010). Dependence of
 641 animal diversity on plant diversity and environmental factors in farmland ponds.
 642 *Community Ecology*, 11, 232–241. <https://doi.org/10.1556/ComEc.11.2010.2.12>
 643 Sarremejane R., Mykrä H., Bonada N., Aroviita J. & Muotka T. (2017). Habitat connectivity
 644 and dispersal ability drive the assembly mechanisms of macroinvertebrate
 645 communities in river networks. *Freshwater Biology*, 62, 1073–1082.
 646 <https://doi.org/10.1111/fwb.12926>

647 Sayer C.D., Davidson T.A. & Jones J.I. (2010). Seasonal dynamics of macrophytes and
 648 phytoplankton in shallow lakes: a eutrophication-driven pathway from plants to
 649 plankton? *Freshwater Biology*, 55, 500–513. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2427.2009.02365.x)
 650 2427.2009.02365.x

651 Scheffer M., van Geest G. J., Zimmer K., Jeppesen E., Sondergaard M., Butler M. G., Hanson
 652 M. A., Declerck S., & De Meester L. (2006). Small habitat size and isolation can
 653 promote species richness: second-order effects on biodiversity in shallow lakes and
 654 ponds. *Oikos*, 112, 227-231.

655 Simonova J., Simon O.P., Kapic S., Nehasil L., & Horsak M. (2016). Medium-sized forest snails
 656 survive passage through birds' digestive tract and adhere strongly to birds' legs:
 657 more evidence for passive dispersal mechanisms. *Journal of Molluscan Studies* 82,
 658 422-426.

659 Slimani N., Sánchez-Fernández D., Guilbert E., Boumaïza M., Guareschi S. & Thioulouse J.
 660 (2019). Assessing potential surrogates of macroinvertebrate diversity in North-
 661 African Mediterranean aquatic ecosystems. *Ecological Indicators*, 101, 324–329.
 662 <https://doi.org/10.1016/j.ecolind.2019.01.017>

663 Thornhill I., Batty L., Death R.G., Friberg N.R. & Ledger M.E. (2017). Local and landscape
 664 scale determinants of macroinvertebrate assemblages and their conservation value
 665 in ponds across an urban land-use gradient. *Biodiversity and Conservation*, 26, 1065–
 666 1086. <https://doi.org/10.1007/s10531-016-1286-4>

667 van Leeuwen C. H. A., van der Velde G., van Lith B., & Klaassen M. (2012). Experimental
 668 quantification of long distance dispersal potential of aquatic snails in the gut of
 669 migratory birds. *PLoS ONE* 7(3): e32292. doi:10.1371/journal.pone.0032292

670 van Donk, E. & Gulati, R.D. (1995). Transition of a lake to turbid state six years after
671 biomanipulation: mechanisms and pathways. *Water Science and Technology*, 32,
672 197-206. [https://doi.org/10.1016/0273-1223\(95\)00699-0](https://doi.org/10.1016/0273-1223(95)00699-0)

673 Vestergaard O. & Sand-Jensen K. (2000). Aquatic macrophyte richness in Danish lakes in
674 relation to alkalinity, transparency, and lake area. *Canadian Journal of Fisheries and*
675 *Aquatic Sciences*, 57, 2022–2031. <https://doi.org/10.1139/f00-156>

676 Vu V.Q. (2011). ggbiplot: A ggplot2 based biplot. R package version 0.55. Available at:
677 <http://github.com/vqv/ggbiplot>

678 Willby N.J., Abernethy V.J. & Demars B.O.L. (2000). Attribute-based classification of
679 European hydrophytes and its relationship to habitat utilization. *Freshwater Biology*,
680 43, 43–74. <https://doi.org/10.1046/j.1365-2427.2000.00523.x>

681 Wood K. A., Stillman R. A., Clarke R. T., Daunt F., & O'Hare, M. T. (2012). The impact of
682 waterfowl herbivory on plant standing crop: a meta-analysis. *Hydrobiologia*, 686,
683 157-167.

684 WWF. 2018. Living Planet Report - 2018: Aiming Higher. Grooten, M. & Almond, R.E.A.(Eds).
685 WWF, Gland, Switzerland.

686

687 Table 1. A summary of environmental characteristics per waterbody type and hydroscape;
688 mean \pm SE (min-max). Land use is representative from within catchments for lakes and the
689 surrounding 500 m buffer for ponds.

Waterbody type	Hydroscape (No. waterbodies surveyed)	Size (ha.)	Altitude (m)	Urban (%)	Agriculture (%)	Freshwater (%)	Wetland (%)
Lake	Upland (n=27)	103.7 \pm 57.1 (1.5 – 1435.8)	166.9 \pm 20.5 (41.0 – 469.0)	0.4 \pm 0.2 (0.0 – 2.8)	14.0 \pm 4.1 (0.0 – 83.2)	7.0 \pm 0.9 (0.8 – 19.4)	0.1 \pm 0.1 (0.0 – 0.3)
	Urban (n=22)	15.3 \pm 4.8 (1.4 – 81.9)	93.3 \pm 11.6 (23.0 – 217.0)	17.1 \pm 5.1 (0.0 – 90.8)	33.2 \pm 4.1 (0.0 – 69.3)	7.5 \pm 1.3 (0.0 – 19.1)	4.2 \pm 1.8 (0.0 – 27.2)
	Agricultural (n=25)	14.5 \pm 3.4 (1.0 – 57.6)	14.6 \pm 4.5 (0.0 – 78.0)	4.2 \pm 1.0 (0.0 – 16.9)	61.7 \pm 5.1 (2.0 – 88.3)	7.9 \pm 2.4 (0.0 – 40.6)	4.9 \pm 2.8 (0.0 – 56.0)
Pond	Upland (n=27)	0.4 \pm 0.1 (0.1 – 1.6)	160.4 \pm 12.3 (64.0 – 306.0)	0.3 \pm 0.1 (0.0 – 2.2)	22.2 \pm 4.7 (0.0 – 75.5)	1.4 \pm 0.5 (0.0 – 12.2)	0.4 \pm 0.2 (0.0 – 5.5)
	Urban (n=26)	0.3 \pm 0.1 (0.1 – 1.2)	92.5 \pm 12.3 (9.0 – 233.0)	39 \pm 5.3 (0.0 – 98.9)	33.1 \pm 4.9 (0.0 – 94.6)	0.5 \pm 0.4 (0.0 – 12.2)	1.1 \pm 0.6 (0.0 – 16.6)
	Agricultural (n=30)	0.2 \pm 0.1 (0.1 – 1.2)	49.2 \pm 5.1 (0.0 – 82.0)	2 \pm 0.5 (0.0 – 13.4)	78.3 \pm 4.4 (14.4 – 99.4)	0.4 \pm 0.2 (0.0 – 4.1)	6.7 \pm 2.9 (0.0 – 58.5)

690

691

692

693

694

695

696

697 Table 2. Summary of species richness and sampling efficiency per waterbody type and
698 hydroscape for each species assemblage. The estimated sample coverage gives an indication
699 of the sampling completeness of each species group per waterbody type per hydroscape.


Waterbody type	Hydroscape (No. waterbodies surveyed)	Species group	Mean richness (range)	Total richness	Estimated sample coverage (%)
Lake	Upland (n=27)	Macrophytes	20 (11-34)	88	95
		Molluscs	4 (0-22)	22	80
		Beetles	13 (3-30)	86	90
		Odonates	6 (2-13)	19	98
	Urban (n=22)	Macrophytes	25 (12-39)	113	95
		Molluscs	8 (1-15)	28	97
		Beetles	16 (6-26)	68	95
		Odonates	5 (1-10)	10	100
	Agricultural (n=25)	Macrophytes	17 (3-29)	87	94
		Molluscs	16 (3-29)	46	99
		Beetles	20 (5-76)	157	87
		Odonates	16 (5-23)	34	98
Pond	Upland (n=27)	Macrophytes	15 (1-25)	86	95
		Molluscs	2 (0-5)	12	90
		Beetles	15 (3-35)	88	94
		Odonates	10 (6-16)	21	99
	Urban (n=26)	Macrophytes	12 (2-19)	84	90
		Molluscs	4 (0-16)	26	90
		Beetles	11 (2-30)	69	95
		Odonates	5 (1-9)	10	100
	Agricultural (n=29)	Macrophytes	11 (1-26)	95	89
		Molluscs	3 (0-12)	29	95
		Beetles	17 (3-50)	130	90

		Odonates	11 (1-25)	29	99
--	--	----------	-----------	----	----

700

701


702 **Figures**



703

704 Figure 1. The conceptual model used to illustrate the direct and indirect relationships between
 705 response variables (macrophyte richness, macrophyte morpho-group diversity, mollusc,
 706 beetle and odonate richness) and explanatory variables (land use, connectivity, physical and
 707 water chemistry metrics).

708



709

710 Fig. 2 Structural equation model (SEM) path diagram for lakes. Arrows are scaled according
 711 to standardised effect sizes, with black arrows indicating positive effects, red arrows negative

and grey arrows indicating specified correlated errors. Explanatory variables with no arrows indicate that they were included in the final SEM but were not significant. Boxes with a superscript represent parameters that had a non-linear relationship with the predictor. Coefficients of determination (R^2) are shown for each response variable. Non-significant relationships ($P > 0.05$) are omitted for clarity.




Fig. 3 Structural equation model (SEM) path diagram for ponds. Arrows are scaled according to standardised effect sizes, with black arrows indicating positive effects and red arrows negative. Explanatory variables with no arrows indicate that they were included in the final SEM but were not significant. Coefficients of determination (R^2) are shown for each response variable. Non-significant relationships ($P > 0.05$) are omitted for clarity.