

# Improving Pacific Oyster (*Crassostrea gigas*, Thunberg, 1793) Production in Mediterranean Coastal Lagoons: Validation of the growth model "ShellSIM" on traditional and novel farming methods

Philip Graham<sup>ab</sup>, Gianni Brundu<sup>b</sup>, Maria Scolamacchia<sup>a</sup>, Angelica Giglioli<sup>c</sup>, Piero Addis<sup>c</sup>, Yuri Artioli<sup>d</sup>, Trevor Telfer<sup>a</sup>, Stefano Carboni<sup>a\*</sup>

<sup>a</sup>University of Stirling, Institute of Aquaculture, Pathfoot Building FK94LA, Stirling, UK.

<sup>b</sup>IMC - International Marine Centre, Loc. Sa Mardini, 09170 Oristano, Italy.

<sup>c</sup>Università degli studi di Cagliari, Department of Life Science and Environment, Via Fiorelli 1, 09126 Cagliari, Italy.

<sup>d</sup>PML - Plymouth Marine Laboratory, Prospect Place, Plymouth PL13DH, UK.

## Abstract

Bivalve farming is a major European aquaculture activity, representing 48.5% of total biomass produced. Italy is one of the largest consumers of oysters but local production does not meet the market demand. Italy has approximately 384,000 ha of shallow lagoons in its coastal area, already devoted to extensive aquaculture activities, which could also represent potential locations for Pacific oyster (*Crassostrea gigas*, Thunberg, 1793) farming.

The aim of this study is to enhance Pacific oyster farming in shallow coastal lagoons by testing novel farming technologies and validating an existing bioenergetic growth model (ShellSIM).

Commercial performance of Pacific oysters and associated environmental parameters were monitored in two Sardinian coastal lagoons (San Teodoro and Santa Gilla, Italy). Oyster growth and survival were compared during a production cycle for two rearing systems: traditional systems (floating bags or lanterns) and Ortac units. The latter has not been previously tested in coastal lagoons. Measured performances were compared with ShellSIM predictions to evaluate the model's ability to predict growth and the potential production in other coastal lagoons.

Results showed that at the end of a six months cycle the oysters mean weight and Condition Index were significantly higher ( $p$  value $< 0.05$ ) in floating bags than in Ortac, ( $55.8 \pm 0.9$  g and  $50.1 \pm 1.3$  g;  $4.6 \pm 0.1$  and  $3.9 \pm 0.1$  respectively). Also, the minimum commercial size (40 g) was reached by 98 % and 68 % of the oyster farmed in floating bags and Ortac units respectively. On the other hand, oysters reared in the Ortac showed a higher survival than in the floating bags ( $95.8 \pm 0.9$  % and  $82.1 \pm 3.4$  %, respectively).

ShellSIM growth predictions were highly correlated with the observed data in both lagoons. However, high values for root mean square deviation (RMSD) indicated that ShellSIM predictions were significantly validated for San Teodoro lagoon but not for Santa Gilla suggesting further tailoring to some environmental conditions to produce more realistic growth predictions.

Results of this study indicate that both floating bags and Ortac system should be employed during the production cycle to maximise oysters' survival and growth performances. Furthermore, this study provides a new validated tool to farmers and stakeholders to monitor oysters' performances and estimate productivity in local waters.

41

## 42 **Keywords**

43 Pacific oysters farming; Shellfish growth model; Farming technologies;

44

## 45 **Abbreviations**

46 POS 1, POS 2 and POS 3 are the three chosen experimental position

47

## 48    **1. Introduction**

49    Italy is one of the main seafood consumers in Europe and amongst the World's top 10 importers,  
50    estimated at 5.6 million US dollars in 2016(FAO, 2016). Different species of shellfish, crustaceans  
51    and fish are farmed using both extensive and intensive methods.

52    In 2016 shellfish farming was the main aquaculture industry, contributing to over 64% of the total  
53    Italian production. This country is the largest producer of Manila clam (*Venerupis philippinarum*;  
54    Adams and Reeve, 1850) and the third producer of Mediterranean mussel (*Mytilus*  
55    *galloprovincialis*; Lamarck, 1819) in Europe. A smaller production includes grooved carpet shell  
56    (*Ruditapes decussatus*; Linnaeus, 1758) and Pacific oyster (*Crassostrea gigas*; Thunberg, 1793)  
57    (Eurofish, 2016; FAO, 2016). Pacific oyster is native to Japan and coastal regions of Asia, and due  
58    to its wide adaptation range at different environmental conditions, is the most widespread cultured  
59    oyster species in the world (Shatkinet *al.*, 1997).

60    In 2016, Europe produced 77,000 tonnes of Pacific oysters, 145 of which were of Italian origin (30  
61    tonnes by a single Sardinian company (FAO 2011-2018Fishstat.J). Italy is one of the largest  
62    consumers of oysters in Europe importing 6,500 tonnes per year primarily from France; this could  
63    represent an opportunity to diversify Italian shellfish farming in the future (Sardegnaagricoltura.it,  
64    2016, FAO, 2016). Sardinia has approximately 10,000 ha of shallow coastal lagoons. This surface  
65    represents 2.6% of the total lagoons area in Italy (Bazzoniet *al.*, 2013). Many of these lagoons are  
66    used for extensive finfish farming, but could be potential sites for Pacific oyster farming.

67    Currently, in the world, three main oyster farming methods are used depending on environmental  
68    conditions such as water depth, tidal range, water exchange rates and bottom substrates: off-bottom  
69    culture, on-bottom culture and suspended culture (Buestelet *al.*, 2009). In Sardinian lagoons  
70    suspended culture is the most commonly used method due to the local environmental conditions.  
71    More specifically, floating bags are designed to keep the oyster growing at the water surface where  
72    most of the food is available. These are manufactured in square and diamond mesh patterns (from 4

73 to 23mm), suspended on the surface thanks to two floaters which allow periodic exposure of the  
74 oysters to the air to reduce biofouling and strengthen the adductor muscle.

75 Amongst suspended oyster culture methods, several new farming tools have been recently  
76 developed, for example Ortac units (ABBLOX), OysterGro© (OysterGro) and Zapco Tumbler  
77 (Zapco Aquaculture). These systems aim at improving oyster production by reducing manual  
78 labour, increasing growth rates and improving oysters' quality (i.e. shell shape). The Ortac system  
79 has been employed in this study. The Ortac system consists of baskets made of polypropylene  
80 plastic and divided in two halves. These operate attached to a trestle and, due to their shape, an up-  
81 welling water flow is passively generated by the surrounding water currents. Furthermore, thanks to  
82 the constant movement under currents actions, this system has been designed to reduce fouling  
83 therefore requiring less handling.

84 Aside from environmental conditions, the use of different grow-out gears affects oyster  
85 performances as suggested by the recent study from Rankin *et al.*, (2018).

86 To date, only one independent trial has been conducted in Scotland to compare growth, survival and  
87 physiological performances of *Ostrea edulis* between Ortac and traditional bag systems (Francouer,  
88 2017). Results of this study indicated that there were no significant differences in growth between  
89 *Ostrea edulis* reared in the two different systems (Ortac units and traditional bags) but higher  
90 survival was observed within the Ortac units. The study presented here is the first investigation and  
91 comparison of the performance of the Ortac system in warmer climates with a smaller tidal range.

92 Much effort has been dedicated to generate and validate growth models for bivalves (Pouvrerau *et*  
93 *al.*, 2006). Most of the energy budget models predicting growth are net production models, which  
94 assume that energy is immediately available for the animal maintenance while the rest is used for  
95 growth or deposited as a reserve. Others are based on a dynamic energy budget approach (DEB)  
96 where energy is first stored as a reserve and then used for different metabolic processes at a  
97 catabolic rate (Kooijman, 2000; Pouvreau *et al.*, 2006; Ren and Ross, 2001; Beadman *et al.*, 2002).

98 Most shellfish energy budget models are only able to simulate growth for locations where they have

99 been calibrated, therefore restricting their use in areas with different environmental conditions  
100 (Hawkins *et al.*, 2013; Dowd, 1997). ShellSIM growth model has been calibrated for 16 shellfish  
101 species in different locations throughout Europe, the U.S.A, China, New Zealand, Malaysia and  
102 Australia. This includes *Mytilus edulis* and *Crassostrea gigas* (ShellSIM, 2011; Hawkins *et al.*,  
103 2013).

104 ShellSIM is based on the principles of energy balance:

105

106 
$$\text{Net energy balance} = \text{Energy ingested} - (\text{Energy egested} + \text{Energy excreted} + \text{Energy expended})$$

107

108 This was developed as a tool to be used by farmers, scientist and environmental regulators  
109 (Hawkins *et al.*, 2013). Consequently, this growth model was considered to be appropriate to  
110 provide growth forecasts in Sardinian coastal lagoons with suitable validation for local conditions.

111 The aim of this study is to validate this existing bioenergetic growth model in two ecologically  
112 different Mediterranean coastal lagoons and for three different oyster farming systems: the Ortac  
113 units, the traditional floating bags and the lantern nets, and furthermore to compare the production  
114 efficiency between the traditional and new farming tools (Ortac and floating bags).

115

## 116 **2. Materials and Methods**

### 117 **2.1. Growth Trial: Ortac vs floating bags**

118 This trial was performed between June 2017 and December 2017 in the lagoon of San Teodoro  
119 (north-east Sardinia: 40°48' 38.08''N, 9°40'26.99''E). A total of 2,400 triploid Pacific oyster seeds  
120 ( $1.7 \pm 0.1$  g,  $2.9 \pm 0.2$  cm), from a French hatchery located in the Loire region of France, were  
121 randomly divided between 6 Ortac units and 6 Floating bags (200 individual per unit, mean total  
122 biomass per unit was  $260.7 \pm 5.6$  g). Thirty oysters from each unit were tagged with an underwater  
123 curing epoxy resin (AquaScape) and biometric parameters were measured every two weeks (i.e.  
124 weight, length, depth and width) using a portable scale (Steinberg SBS-LW-2000A, 0.01g) and

125 callipers (METRICA, 0.05 mm). At each sampling point, mortality was also recorded and 5 oysters  
126 per unit were selected for dry weight measurements (Mo and Neilson, 1994) and Condition Index  
127 (CI) calculations using the protocols described by Mo and Neilson (1994) for the dry weight and  
128 Davenport and Chen (1987) and Walne and Mann (1975) for CI calculations:

129

$$130 \quad CI = (\text{Dry weight meat (g)} / \text{Dry weight meat (g)} + \text{Dry weight shell (g)}) \times 100$$

131

132 The oyster culture systems were positioned in two rows of three Ortac units and three floating bags  
133 (fig.1). Ortac units were mounted onto two trestles (3 units per trestle), floating bags were attached  
134 to ropes as in the usual commercial setting of the Compagnia Ostricola Mediterranea, host of these  
135 trials.

136 Oysters in both systems were cultured following the standard conditions of the company, with 24hrs  
137 of air exposure every two weeks to prevent biofouling, changing of the floating bags mesh (4, 9, 14  
138 and 19mm) according to oysters' size, and based on the increasing Pacific Oysters biomass. Grading  
139 was performed when the biomass in each farming unit reached about 4kg live weight and generally  
140 once every three months for both Ortac units and floating bags in order to keep similar biomass in  
141 both systems.

142

## 143 2.2. ShellSIM validation for floating Bag units (San Teodoro Lagoon)

144 A survey of the dominant currents in the area was conducted during the neap (minimum) and spring  
145 (maximum) tides using GPS tracking drifters (drogues) (Cromeey and Black, 2005). The drifters  
146 were released at the same time at three different points, one hour before the tidal peak until one hour  
147 after. During the survey the wind direction and speed was recorded by a fixed weather station (La  
148 Crosse WS3650). These information were fed to ShellSIM model to account for the hydrodynamic  
149 and ecological conditions within the lagoon.

Three floating bags per each area were stocked with *C. gigas* ( $838 \pm 36.4$  g,  $811.5 \pm 17.8$  g and  $709.8 \pm 40.1$  g total biomass) with a mean size in weight and length of  $4.5 \pm 0.3$  g and  $4.0 \pm 0.2$  cm (POS 1),  $4.5 \pm 0.3$  g and  $3.9 \pm 0.2$  cm (POS 2),  $3.9 \pm 0.2$  g and  $3.9 \pm 0.2$  cm (POS 3). The oysters were cultured following the standard procedures described above.

Sampling for oyster growth was performed monthly for 5 months. Each month 80 individuals unit<sup>-1</sup> were randomly measured for wet weight, 30 of which also measured for length, depth and width. Other 10 individuals unit<sup>-1</sup> were collected for dry weights measurements.

Environmental data: temperature (T, °C), salinity (Sal, ‰), dissolved oxygen (DO, mgL<sup>-1</sup>), total particulate matter (TPM, mg/L), particulate organic matter (POM, mgL<sup>-1</sup>), particulate organic carbon (POC, mg m<sup>-3</sup>) and Chlorophyll-a (Chl-a, µgL<sup>-1</sup>) were collected in the immediate vicinity of the farming gears. Temperature, salinity and dissolved oxygen were collected at a depth of 15 cm, with a multi-parametric probe (HACH HQ40d) and data loggers (HOBO: UTBI-001, U26-001 and U24-002-C respectively for T, DO and Sal). Temperature data loggers were set-up to take measurements every 30 minutes, while the Sal and DO probes measured values every 2 hrs.

Water for the TPM, POM, POC was collected using 1L pre-rinsed in sample water plastics bottles, while 5L pre-rinsed in sample water plastic bottles were used to collect water for Chl-a analysis. Laboratory analysis for TPM and POM were performed according to Hawkins *et al.* (2013), while Chl-a analysis according to Lorenzen (1967). For POC measurements water was collected in 1L pre-rinsed plastic bottles then filtered in 47-mm diameter GF/F filters previously combusted at 450 °C. POC samples were analysed with a CEI Flash smart elemental analyser. The average values of each environmental parameter were used to run the model, excluding September 2017, when no data were collected due to farmers' activities and weather constraints.

172

### 2.3. ShellSIM validation for lantern systems (Santa Gilla Lagoon)

In order to validate the growth model in a different location, a growth trial was performed between May 2017 and September 2017, in Santa Gilla lagoon ( $39^{\circ}12'28.2''$ N  $9^{\circ}05'53.5''$ E). Three lanterns

176 with five compartments each and a mesh size of 3.5 x 5 mm, were stocked with 500 oysters per  
177 compartment (mean weight =  $4.4 \pm 0.1$  g; mean length =  $3.6 \pm 0.6$  cm). The oysters were farmed  
178 following the standard production protocols, grading and changing the mesh size according to  
179 oysters' size and biomass. Growth was measured monthly when 70 individuals per lantern were  
180 randomly sampled and weighted, 30 of which were also measured for shell length, depth and width.  
181 Furthermore, 10 individuals per lantern were collected for dry weight measurements.  
182 Environmental data sampling and analysis were conducted as described above. The monthly means  
183 of all the environmental data were used to run the growth model.

184

#### 185 2.4. ShellSIM Validation for Ortac units and floating Bags (San Teodoro Lagoon)

186 In order to validate the model for different gear types, a new experiment was set up in the lagoon of  
187 San Teodoro (July 2017 – December 2017) where the model performance was also tested on a  
188 different farming system, the Ortac units.

189 Farming methods, growth measurements, sample collection and analysis of all environmental  
190 parameters were conducted as described previously. A bi-weekly mean of all the environmental data  
191 were used to run the ShellSIM, except for November and December, when data were collected only  
192 once per month due to farmers' activities and weather constraints.

193

#### 194 2.5. Statistical Analysis

195 Prior to analyses, data were tested for normality and homogeneity of variance. Weight gain,  
196 biometrics measures differences, survival rate and Condition index differences, over time were  
197 analysed by general linear model followed by a Tukey post-hoc test where significant differences  
198 occurred.

199 End points of all biometrics measures, survival rate and condition index, were analysed by one-way  
200 ANOVA followed by post-hoc Tukey's Multiple Comparison tests where significant differences  
201 occurred.



202 To assess fitness between the prediction made by ShellSIM and observed data, Taylor diagrams and  
203 skill scores (S) were used (Taylor, 2001). A Taylor diagram is a way to show graphically how well a  
204 model prediction fits the observed data, using correlation, centred root mean square difference  
205 (RMSD) and amplitude of their variation (standard deviations). The skill score proposed by Taylor  
206 (2001) quantifies model performance against observed data.

207

### 208 **3. Results**

#### 209 **3.1. Growth Trial: Ortac vs floating bags**

210 At the end of the production cycle (October to December), the Pacific oysters farmed in the floating  
211 bags had a significantly higher weight and shell depth ( $p$  value = 0.001, 0.001 respectively) to those  
212 in the Ortac units ( $55.8 \pm 0.9$  g,  $50.1 \pm 1.3$  g;  $26.6 \pm 0.2$  g,  $24.2 \pm 0.3$  g) (figs. 2a, 2c). Oysters farmed  
213 inside the Ortac units showed instead a significant higher growth in shell length ( $86.9 \pm 1$ ,  $75.4 \pm$   
214  $0.6$ ,  $p$  value = 0.001, Ortac and floating bags respectively), and shell width ( $46.2 \pm 0.5$  mm,  $44.6 \pm$   
215  $0.4$  mm,  $p$  value = 0.017) (fig.2b,d).

216 Survival was significantly higher ( $p$  value= 0.001) in the Ortac units compared to the floating bags  
217 ( $95.8 \pm 0.9\%$ ,  $82.1 \pm 3.4$  %) (fig.3). The highest mortality occurred between June and July ( $3.8 \pm 1$   
218  $\%$ ,  $16.3 \pm 3.3$  % Ortac units and floating bags respectively).

219 The condition index at the end of the production cycle was significantly higher ( $p$  value = 0.001) in  
220 the floating bags compared to the Ortac units ( $4.6 \pm 0.1$ ,  $3.9 \pm 0.1$ ) (fig.4) and the smallest  
221 commercial size (40 g) was reached by the 98 % and 69 % of the oyster farmed in the floating bags  
222 and Ortac units respectively (fig.5).

223

#### 224 **3.2. ShellSIM validation in San Teodoro Lagoon**

225 Three areas with different current speed (POS1  $0.15 \text{ m s}^{-1}$ , POS2  $0.07 \text{ m s}^{-1}$ ; POS 3  $0.04 \text{ m s}^{-1}$ ) were  
226 identified in San Teodoro. A decreasing speed gradient from the sea mouth to the internal part of

the lagoon was identified. Consequently, these areas were used as experimental locations to monitor the oysters' growth and the environmental parameters required by the growth model.

Environmental data are illustrated in Table 1. ShellSIM predicted, during a 6 months production cycle, a final weight and length of 19.7 g, 48.4 g and 121.6 g; 6.0 cm, 8.3 cm and 11.5 cm, respectively for POS1, POS2 and POS3.

The measured weight and length at the end of this production cycle, was  $16.4 \pm 1.1$  g,  $46.9 \pm 2.1$  g,  $48.9 \pm 1.5$  g and  $5.4 \pm 0.3$  cm,  $8.2 \pm 0.3$  cm,  $9 \pm 0.2$  cm, respectively in POS1, POS2 and POS3.

Figure 6 shows that measured growth in weight and length, fitted the predicted growth curve in POS2, while in POS1 and POS3 ShellSIM overestimate the final mean growth in weight and length respectively 20.5, 12.1, 148.8 and 27.9 %. The calculated skill score for the three different areas indicate the best fitting between observed and predicted measures of weight and length, respectively in POS2 ( $S=1$ ;  $S=1$ ), POS1 ( $S=0.87$ ;  $S=0.81$ ) and POS3 ( $S=0.42$ ;  $S=0.79$ ).

Standard deviation, Centred Root Mean square difference (RMSD), correlation and the overall skill score of the performance of the predicted growth curve to fit the observed data in the lagoon of San Teodoro are shown in Figure 7 and Table 2.

### 3.3. ShellSIM validation in Santa Gilla Lagoon

Environmental data collected in the lagoon of Santa Gilla and their seasonal variations are illustrated in Table 3.

The measured growth in weight and length ( $79.5 \pm 1.8$  g and  $9.1 \pm 0.1$  cm) did not fit the predicted growth curve (data not shown), and the calculated skill score indicates a very poor fit between observed and predicted measures of weight and length, respectively  $S=0.003$  and  $S=0.17$  (Table 2).

Standard deviation, Centred Root Mean square difference (RMSD) and correlation are shown in Figure 8 and Table 2.

### 3.4 ShellSIM Validation on Ortac vs floating bags

Environmental data collected to run ShellSIM and their changes are shown in Table 4.

In this trial, ShellSIM was run in POS2 for two different farming systems. It predicted a growth of 48.6 g and 8.3 cm in weight and length respectively for the Ortac system, and a growth of 49.1 g and 8.2 cm for the floating bags over a 6 months production cycle. At the end of this production cycle, the measured weight and length were  $50.1 \pm 1.3\text{g}$  and  $8.7 \pm 0.1\text{cm}$  for the Ortac and  $55.8 \pm 0.9\text{ g}$  and  $7.5 \pm 0.1\text{ cm}$  for the floating bags. Figure 9 shows that during the production cycle, the measured mean weight and length in the Ortac units and floating bags were underestimated by ShellSIM, except for the final length farmed in the floating bags which was accurate. Indeed, in November there was a change in trend of the model prediction, from underestimation to overestimation (the model overestimated the final mean length of 8.2 %), while the final weight was still overestimated by 11.9%.

Moreover, Figure 9 shows that ShellSIM at the end point of the production cycle of the oysters reared inside the Ortac units, unlike the rest of the predictions, slightly underestimated growth in weight and length by 3% and 4.4% respectively.

The calculated skill score indicates that the best fitting between observed and predicted measures of weight and length was respectively obtained in Ortac ( $S=0.95$ ,  $S=0.93$ ) compared to floating bags ( $S=0.90$ ,  $S=0.89$ ). Standard deviation, Centred Root Mean square difference (RMSD), correlation and the overall skill score of the performance of the predicted growth curve to fit the observed data in this trial in the SanTeodoro lagoon are shown in Figure 10 and Table 2.

272

#### 273 4. Discussion and conclusions

274 The results of this study provide new information to improve *C. gigas* farming and a growth  
275 prediction tool in shallow coastal lagoons. The higher survival rate in the Ortac units for the first two  
276 months and the higher growth in weight and CI in the floating bags, suggest a potential mixed use of  
277 the two systems during the production cycle. Specifically, the Ortac units may be employed when  
278 Pacific oysters are more susceptible to stress and during the stressful period (e.g smaller size and

hottest periods) and floating bags thereafter. As a result, by combining the two farming gears within one production cycle, it would be possible to reduce the capital costs of equipment by reducing the need for several meshes sizes in the floating bag system, and achieve a significantly higher survival rate from seed to market size individuals.

There was no statistically significant difference in growth between Ortac units and floating bags, but at the end of the production cycle there was a significant higher mean weight in the floating bags than in the Ortac units. Comparison of these results with previous studies is difficult due to difference in culture techniques, local environment, species used, initial oyster size and the production season.

Many studies report that the shell morphology in bivalves is influenced by population density, predation responses, handling and grow-out methods (Telesca *et al.*, 2018; Seed, 1968; Brake *et al.*, 2003; Kubeet *al.*, 2011; Griffiths and Buffenstein, 1981; Van Erkom Schurink and Griffiths, 1993; Bayne, 2000; Sheridan *et al.*, 1996). As in these studies, we observed a difference in shape between the animals reared inside the floating bags or Ortac units, with the latter showing longer and wider shells compared to the former which were instead thicker and with a higher C.I.

The morphological differences found between individuals farmed in Ortac and floating bags are probably due to the shape of these different tools, and consequently the different interaction of these with the currents. Under low current speed typical of shallow lagoons, the shape of the Ortac units may have not promoted the rocking motion required to generate enough rubbing between oysters and the farming gear, causing less shell chipping, which is widely recognised as a factor promoting shell depth and a higher meat content (Brake *et al.*, 2003; Holliday, 1991; O'Meley, 1995; Robert *et al.*, 1993).

Moreover, the fact that the animals did not move enough inside the Ortac units probably induced those in the innermost part to grow more in length and width in order to increase the filtering surface. Nonetheless, results of this study are comparable with those obtained by Francouer *et al.*, (2017). In his study, performed in Scotland, oysters' growth differences were found to be not

305 statistically significant between the two systems, consistently with data reported here. Mortality may  
306 depend on the farming system (Pernet *et al.*, 2012). Improved survival in the Ortac system could be  
307 due to the shading effect provided by a more solid structure, which would shelter farmed individuals  
308 from direct sunlight and desiccation, particularly during the earlier part of the growth cycle and  
309 during air exposure periods (Potter and Hill, 1982; Spencer-Davies, 1970). Moreover, different  
310 studies report that one of the stress factors associated to mortalities is temperature, and sudden small  
311 changes may have a large effect on the survival of bivalves (Pernet *et al.*, 2018; Pernet *et al.*, 2012;  
312 Petton *et al.*, 2015; Le Deuff *et al.*, 1994, Le Deuff *et al.*, 1996; Sauvage *et al.*, 2009; Kennedy and  
313 Mihursky, 1971). Again the more solid structure of the Ortac may have promoted more stable  
314 temperature and reduced stress.

315 Results of this study indicate that the predicted growth by ShellSIM fitted well with field  
316 measurements in the lagoon of San Teodoro. However, results from the growth trial in Santa Gilla  
317 lagoon demonstrate that the model would require further tailoring to local conditions to produce  
318 realistic growth projections. In particular, we tested the hypothesis that ShellSIM assumptions on the  
319 conversion of food concentration into available/digestible energy for the oysters, may not apply to  
320 Santa Gilla Lagoon. In order to do this, we run the model reducing the amount of POC available to  
321 one quarter of the measured POC and the model prediction was more accurate ( $S = 0.97$  and  $S =$   
322  $0.95$  respectively for weight and length). Indeed, POC can be considered as a very heterogeneous  
323 nutrient source composed by different materials with large variations in digestible energy content  
324 (Mazzola and Sarà, 2001; Lawacz, 1977; Watanabe and Kuwae, 2015).

325 Further studies to identify the real digestible energy content of the Particulate Organic Carbon in  
326 Santa Gilla area is required to modify the model assumption and improve its performances. Our data  
327 also suggest that seasonality and farming system used can influence the accuracy of ShellSIM  
328 providing scope for further tailoring of the model to reflect gear types and local environmental  
329 conditions.

330 During the first-year trial in the lagoon of San Teodoro the measured growth closely fitted the  
331 predicted growth in POS2, while in POS1 ShellSIM slightly overestimated and in POS3  
332 considerably overestimated the growth, both in weight and length. Similar results in POS2 were  
333 observed in the second-year validation trial. The growth in weight and length of the oyster was  
334 different between the two farming tools, with a higher growth in weight recorded for oysters reared  
335 in the floating bags and a higher growth in length for oysters reared in the Ortac units. In this trial,  
336 ShellSIM underestimated the weight and length during the production cycle except at the end point  
337 where it only slightly underestimated weight and length in the Ortac units providing a better  
338 accuracy at harvest time. While in the floating bags the final mean weight was underestimated and  
339 the length was overestimated.

340 These overestimation and underestimation can be potentially associated with a less than optimal  
341 rearing method (the Ortac), combined with the potential different production capacity of each  
342 farming areas within the lagoon. Furthermore, as reported by several authors, the grow-out methods  
343 employed could affect oyster growth (Bayne, 2000; Sheridan *et al.*, 1996). ShellSIM does not  
344 consider different grow-out methods in its variables, possibly generating the discrepancy between  
345 observed and predicted growth measured in this study. Overall, ShellSIM predictions correspond  
346 with the growth trends observed by the farmers over the years (POS3 with higher growth rates and  
347 POS1 with lower growth rates) suggesting the good accuracy of the model with the general growth  
348 dynamics in the different areas of San Teodoro lagoon. This is reflected in the calculated skill  
349 scores, for both validation trials in the fore mentioned lagoon.

350 Taken together, the results of this study provide information to improve bivalve growth prediction  
351 tools for Mediterranean lagoons. They could be applied to study the productivity of different sites to  
352 potentiate the oyster's aquaculture industry and for coastal spatial planning. Moreover, the presented  
353 results indicate that Ortac units improve the oyster's survival in the production early stage. The use  
354 of Ortac units also reduces reliance on multiple mesh bags therefore simplifying production  
355 protocols.

356

357 **Acknowledgments**

358 This study is part of the Sardinian project "OSTRINNOVA, Valorisation of sustainable production  
359 of oysters in the shellfish production system in Sardinia", funded by the Sardinia Region, Det. DG n  
360 ° 566 of 27/04/2016. Many thanks to Alessandro Gorla and Francesca Gargiulli and all the staff of  
361 the Compagnia Ostricola Mediterranea and the cooperativa lo Squalo, for the hospitality, help and  
362 availability during the field work. The authors would like to thank Ian Washbourne for the POC  
363 analysis in Stirling University, Andrea Cucco and Marilisa Lo Prejato for their help with Taylor's  
364 diagrams and Tony Legg for providing the Ortac units. Yuri Artioli's contribution has been partly  
365 supported by the Natural Environment Research Council single centre national capability  
366 programme – Climate Linked Atlantic Sector Science. Finally, we would like to thank the  
367 anonymous reviewers who helped improve the original manuscript.

368




Figure 1. Diagram of the experimental layout of the Ortac and Floating Bags (FB) in the San Teodoro Lagoon.





Figure 1. (a) Difference growth in weight between *C. gigas* farmed in two different tools (Ortac units and Floating bags). (b) Difference growth in length between *C. gigas* farmed in two different tools (Ortac units and Floating bags). (c) Difference growth in width between *C. gigas* farmed in two different tools (Ortac units and Floating bags). (d) Difference growth in depth between *C. gigas* farmed in two different tools (Ortac units and Floating bags). Stars indicate where significant difference occurs. ( $p$ -value < 0.05). Data are presented as mean  $\pm$  SE; n=6.






380

381 *Figure 2.* Comparison of survival rate between *C. gigas* farmed inside the Ortac units and Floating bags. Stars indicate where  
 382 significant difference occurs. (p-value < 0.05). Data are presented as mean ± SE; n=6.

383



384

385 *Figure 3.* Comparison of condition index (CI) calculated as (Dry weight Meat (g) / Dry weight Meat + Dry weight shell) \* 100,  
 386 between *C. gigas* farmed inside the Ortac units and Floating bags. Stars indicate where significant difference occurs. (p-value < 0.05).  
 387 Data are presented as mean ± SE; n=6.

388



Figure 4. Comparison of size class distribution between *C. gigas* farmed inside the Ortac units and Floating bags.






Figure 5. ShellSIM growth prediction compared to the measured oyster growth in weight and length, during a production cycle performed in three different areas (POS1, POS2 and POS3) of the San Teodoro lagoon. Measured growth data are presented as mean  $\pm$  SE; n=3.

SHELLSIM VALIDATION IN SAN TEODORO LAGOON (g)



SHELLSIM VALIDATION IN SAN TEODORO LAGOON (cm)




397


398 Figure 6. Taylor diagrams representing how closely model performance (B) match the observed data (A). The similarity between  
 399 model prediction and observed data is quantified in terms of their correlation, the amplitude of their variation (normalised standard  
 400 deviation) and their root mean square difference (RMSD) (dashed circular arcs). The left panel contain the results for the ShellSIM  
 401 validation in the san Teodoro lagoon in terms of predicting the overall growth in weight of the *C.gigas* farmed inside the floating  
 402 bags. The right panel contain the results for the ShellSIM validation in the san Teodoro lagoon in terms of predicting the overall  
 403 growth in length of the *C.gigas* farmed inside the floating bags.

404

SHELLSIM VALIDATION IN SANTA GILLA LAGOON (g)




SHELLSIM VALIDATION IN SANTA GILLA LAGOON (cm)



405

406 Figure 8. Taylor diagrams representing how closely model performance (B) match the observed data (A). The similarity between  
 407 model prediction and observed data is quantified in terms of their correlation, the amplitude of their variation (normalised standard  
 408 deviation) and their root mean square difference (RMSD) (dashed circular arcs). The left panel contain the results for the ShellSIM  
 409 validation in the Santa Gilla lagoon in terms of predicting the growth in weight of the *C.gigas*. The right panel contain the results for  
 410 the ShellSIM validation in the Santa Gilla lagoon in terms of predicting the growth in length of the *C.gigas*.




412

413 *Figure 9.*ShellSIM growth prediction compared to the measured oyster growth in weight and length,during a production cycles  
414 performed in to two different farming systems (Ortac units and Floating bags) in the San Teodoro lagoon (July 2017 – December  
415 2017).Measured growth data are presented as mean  $\pm$  SE; n=6.

416

SHELLSIM VALIDATION ORTAC AND FLOATING BAGS (g)



SHELLSIM VALIDATION ORTAC AND FLOATING BAGS (cm)




Figure 10. Taylor diagrams representing how closely modelled performances (B) matched the observed data (A). The similarity between model prediction and observed data is quantified in terms of their correlation, the amplitude of their variation (normalised standard deviation) and their root mean square difference (RMSD) (dashed circular arcs). The left panel contain the results for the ShellSIM validation in the San Teodoro lagoon on Ortac and Floating bags in terms of predicting the growth in weight of the *C.gigas*. The right panel contain the results for the ShellSIM validation in the San Teodoro lagoon on Ortac and Floating bags in terms of predicting the growth in length of the *C.gigas*.

Table 1. Summary of the environmental data collected to run ShellSIM. These data were collected during the production cycles started in August 2016, in three different areas (POS1, POS2 and POS3) of the San Teodoro lagoon. Data are presented as mean  $\pm$  SE.

	<i>T °C</i>	<i>Sal</i> ‰	<i>DO</i> mg/L	<i>TPM</i> mg/L	<i>POM</i> mg/L	<i>POC</i> mg/m <sup>3</sup>	<i>Chl-a</i> µg/L
<b>August 2016</b>							
<b>POS 1</b>	27.1 $\pm$ 0.1	39.3 $\pm$ 0.1	7.2 $\pm$ 0.2	31.5 $\pm$ 12.4	5.3 $\pm$ 1.9	848.2 $\pm$ 18.6	2.1 $\pm$ 0.2
<b>POS 2</b>	27.2 $\pm$ 0.1	39.7 $\pm$ 0.1	7.5 $\pm$ 0.1	15.5 $\pm$ 1.6	3.2 $\pm$ 0.3	1213.1 $\pm$ 67.9	4.3 $\pm$ 0.7
<b>POS 3</b>	27.2 $\pm$ 0.2	39.5 $\pm$ 0.1	6.8 $\pm$ 0.1	19.4 $\pm$ 1.1	3.6 $\pm$ 0.2	1421.9 $\pm$ 68.5	4.1 $\pm$ 0.1
<b>October 2016</b>							
<b>POS 1</b>	23.5 $\pm$ 0.1	39.2 $\pm$ 0.1	8.7 $\pm$ 0.1	5.2 $\pm$ 0.2	1 $\pm$ 0.1	206.9 $\pm$ 6	0.3 $\pm$ 0.1
<b>POS 2</b>	24.1 $\pm$ 0.1	38.8 $\pm$ 0.1	8.5 $\pm$ 0.1	5 $\pm$ 0.2	1 $\pm$ 0.1	211.3 $\pm$ 18.1	0.3 $\pm$ 0.1
<b>POS 3</b>	23.1 $\pm$ 0.3	38.8 $\pm$ 0.1	8.9 $\pm$ 0.3	21.1 $\pm$ 2.4	3.9 $\pm$ 0.2	1192.5 $\pm$ 55.8	3.7 $\pm$ 0.3

**November 2016**

<b>POS 1</b>	$17.1 \pm 0.1$	$39.2 \pm 0.1$	$9.3 \pm 0.1$	$0.6 \pm 0.03$	$0.5 \pm 0.1$	$167.3 \pm 9.1$	$0.4 \pm 0.02$
<b>POS 2</b>	$15.8 \pm 0.5$	$37.7 \pm 0.5$	$8.8 \pm 0.2$	$2.3 \pm 0.1$	$1 \pm 0.1$	$485.2 \pm 33.9$	$2.8 \pm 0.2$
<b>POS 3</b>	$14.5 \pm 0.2$	$38.8 \pm 0.1$	$9.6 \pm 0.1$	$3.0 \pm 0.04$	$1.1 \pm 0.1$	$473.4 \pm 20.4$	$2.9 \pm 0.1$

**December 2016**

<b>POS 1</b>	$16 \pm 0.1$	$36.4 \pm 0.2$	$10 \pm 0.2$	$1.7 \pm 0.1$	$0.6 \pm 0.1$	$232.8 \pm 21$	$0.7 \pm 0.1$
<b>POS 2</b>	$17.5 \pm 0.3$	$37 \pm 0.3$	$10.7 \pm 0.4$	$1.0 \pm 0.1$	$0.6 \pm 0.1$	$199.1 \pm 28.4$	$0.4 \pm 0.1$
<b>POS 3</b>	$15.1 \pm 0.3$	$36.2 \pm 0.2$	$9.2 \pm 0.6$	$4.8 \pm 0.2$	$1.2 \pm 0.1$	$408.4 \pm 24$	$0.9 \pm 0.03$

Table 2. Summary of how well observed data match predicted data by ShellSIM in terms of their correlation, their root-mean square difference (RMSD), the ratio of their variances and skill score (Taylor, 2001).

**SHELLSIM VALIDATION IN SAN TEODORO LAGOON (FLOATING BAGS)**

	<i>St.dev Obs.</i>	<i>St.dev Pred.</i>	<i>RMSD</i>	<i>Correlation</i>	<i>Skill score</i>
<b>POS1 (g)</b>	0.3	0.43	0.14	0.98	0.87
<b>POS1 (cm)</b>	0.27	0.43	0.16	1	0.81
<b>POS2 (g)</b>	0.35	0.36	0.02	1	1
<b>POS2 (cm)</b>	0.36	0.37	0.02	1	1
<b>POS3 (g)</b>	0.14	0.38	0.24	0.99	0.42
<b>POS3 (cm)</b>	0.24	0.38	0.16	0.97	0.79
<b>OVERALL (g)</b>	0.32	0.4	0.2	0.87	0.83
<b>OVERALL (cm)</b>	0.31	0.4	0.17	0.92	0.87

**SHELLSIM VALIDATION ON ORTAC AND FLOATING BAGS IN SAN TEODORO LAGOON**

<b>ORTAC (g)</b>	0.32	0.30	0.1	0.95	0.95
<b>ORTAC (cm)</b>	0.31	0.28	0.11	0.93	0.93
<b>FLOATING BAGS (g)</b>	0.33	0.27	0.12	0.94	0.9
<b>FLOATING BAGS (cm)</b>	0.27	0.31	0.13	0.9	0.89
<b>OVERALL (g)</b>	0.32	0.29	0.11	0.94	0.93
<b>OVERALL (cm)</b>	0.29	0.30	0.14	0.89	0.9
<b>SHELLSIM VALIDATION IN SANTA GILLA LAGOON</b>					
<b>SANTA GILLA (g)</b>	0.01	0.4	0.39	0.97	0
<b>SANTA GILLA (cm)</b>	0.08	0.38	0.3	0.96	0.18

433

434

435

436 *Table 3. Summary of the environmental data used to run ShellSIM. These data were collected during the production cycles started in*  
437 *June 2017, in the Santa Gilla lagoon. Data are presented as mean ± SE.*

	<b><i>T °C</i></b>	<b><i>Sal ‰</i></b>	<b><i>DO mg/L</i></b>	<b><i>TPM mg/L</i></b>	<b><i>POM mg/L</i></b>	<b><i>POC mg/m<sup>3</sup></i></b>	<b><i>Chl-a µg/L</i></b>
<b><i>June 2017</i></b>	24.7 ± 0.02	37.6 ± 0.03	7 ± 0.03	2.3 ± 0.2	0.9 ± 0.1	413.1 ± 8.2	0.7 ± 0.1
<b><i>July 2017</i></b>	24.8 ± 0.02	43.3 ± 0.1	6.1 ± 0.04	4.6 ± 1.1	1 ± 0.1	349.5 ± 3.3	0.7 ± 0.1
<b><i>August 2017</i></b>	27 ± 0.01	35.4 ± 0.03	5.2 ± 0.04	6.6 ± 0.9	1.4 ± 0.1	451.4 ± 12.6	1.8 ± 0.1
<b><i>September 2017</i></b>	24.4 ± 0.02	36 ± 0.03	6 ± 0.03	8.1 ± 0.7	2.1 ± 0.3	561.6 ± 19.9	1.9 ± 0.1

438

439

440 *Table 4. Summary of the environmental data used to run ShellSIM. These data were collected during the production cycles started in*  
441 *July 2017, in the San Teodoro lagoon. Data are presented as mean ± SE.*

<b><i>T °C</i></b>	<b><i>Sal ‰</i></b>	<b><i>DO mg/L</i></b>	<b><i>TPM mg/L</i></b>	<b><i>POM mg/L</i></b>	<b><i>POC mg/m<sup>3</sup></i></b>	<b><i>Chl-a µg/L</i></b>
--------------------	---------------------	-----------------------	------------------------	------------------------	------------------------------------	--------------------------

<b>July 2017</b>	$27.5 \pm 0.2$	$38.9 \pm 0.4$	$9.6 \pm 0.3$	$6.1 \pm 4.0$	$1.7 \pm 1.2$	$358.2 \pm 62.7$	$1.7 \pm 0.7$
<b>August 2017</b>	$28.4 \pm 0.3$	$41.1 \pm 0.4$	$7.8 \pm 0.3$	$3.9 \pm 0.8$	$1.5 \pm 0.2$	$557.6 \pm 94.2$	$3.1 \pm 1.1$
<b>September 2017</b>	$21.5 \pm 0.1$	$40.2 \pm 1.1$	$9.2 \pm 0.1$	$5.4 \pm 0.3$	$1.5 \pm 0.2$	$491.1 \pm 30.1$	$1.7 \pm 0.6$
<b>October 2017</b>	$18 \pm 2.1$	$40.7 \pm 0.1$	$8.3 \pm 0.3$	$5.2 \pm 0.3$	$1.4 \pm 0.1$	$769.2 \pm 99$	$2.8 \pm 0.6$
<b>November 2017</b>	$18.3 \pm 0.3$	$38.4 \pm 0.3$	$9.4 \pm 0.1$	$23.2 \pm 3.3$	$3 \pm 0.4$	$151.3 \pm 23.9$	$0.5 \pm 0.03$
<b>December 2017</b>	$14.7 \pm 0.1$	$36.7 \pm 0.5$	$10 \pm 0.1$	$12.3 \pm 9.5$	$1.9 \pm 1.3$	$222.4 \pm 67.4$	$0.8 \pm 0.3$

442

443

#### 444 Reference List

- 445 1. © FAO 2011-2018. Fisheries and aquaculture software. FishStatJ - software for fishery  
446 statistical time series. In: *FAO Fisheries and Aquaculture Department* [online]. Rome.  
447 Updated 21 July 2016. [Cited 1 November 2018]. <http://www.fao.org/fishery/>
- 448 2. Bayne, B. L. (2000). Relations between variable rates of growth, metabolic costs and growth  
449 efficiencies in individual Sydney rock oysters (*Saccostrea commercialis*). *Journal of*  
450 *Experimental Marine Biology and Ecology* **251**, 185-203.
- 451 3. Bazzoni, A.M., Pulina, S., Padedda, B.M., Satta, C.T., Lugliè, A., Sechi, N., Facca, C.  
452 (2013). Water quality evaluation in Mediterranean lagoons using the Multimetric  
453 Phytoplankton Index (MPI): Study cases from Sardinia. *Transitional Waters Bulletin* **7 (1)**,  
454 64-76.
- 455 4. Beadman, H.A., Willows, R.I., Kaiser, M.J. (2002) Potential application of mussel  
456 modelling. *Helgoland Marine Research* **56 (1)**, 76-85.



5. Brake, J., Evans, F., Langdon, C. (2003). Is beauty in the eye of the beholder? Development of a simple method to describe desirable shell shape for the Pacific oyster industry. *Journal of Shellfish Research***22**, 767-772.
6. Buestel, D., Ropert, M., Prou, J., Goulletquer, P. (2009) History, Status, and Future of Oyster Culture in France. *Journal of Shellfish Research***28** (4), 813-820.
7. Cromey C.J., Black K.D. (2005). Modelling the Impacts of Finfish Aquaculture. In: Hargrave B.T. (eds) Environmental Effects of Marine Finfish Aquaculture. Handbook of Environmental Chemistry, vol 5M. Springer, Berlin, Heidelberg.
8. Davenport, J., Chen, X. (1987) A comparison of methods for the assessment of condition in the mussel (*Mytilus edulis* L.). *Journal of Molluscan Studies***53**, 293-297.
9. Dowd, M. (1997) On predicting the growth of cultured bivalves. *Ecological Modelling***104**, 113-131.
10. Eurofish (2016). Eurofish International Organisation. Developing fisheries and aquaculture in Europe. [Online] Available from: <https://www.eurofish.dk/italy> [Accessed: 19<sup>th</sup> March 2019].
11. FAO (2016) Food and Agriculture Organization of the United Nations [Online] Available from: <http://www.fao.org/3/i9540EN/i9540en.pdf> [Accessed: 12<sup>th</sup> June 2018].
12. Francouer, (2017). Assessment of productivity of novel oyster farming methods. MSc Thesis, Institute of Aquaculture, University of Stirling FK9 4LA, UK.
13. Griffiths, R. J., Buffenstein, R. (1981). Aerial exposure and energy input in the bivalve *Choromytilus meridionalis* (Kr.). *Journal of Experimental Marine Biology and Ecology***52**, 219-229.
14. Hawkins, A.J.S., Pascoe, P.L., Parry, H., Brinsley, M., Black, K.D., McGonigle, C., Moore, H., Newell, C.R., O'Boyle, N., O'Carroll, T., O'Loan, B., Service, M., Small, A.C., Zhang, X.L., Zhu, M.Y. (2013) Shellsim: A Generic Model of Growth

and Environmental Effects Validated Across Contrasting Habitats in Bivalve Shellfish. *Journal of Shellfish Research***32** (2), 237-253.

15. Holliday, J. E. (1991). Nursery Culture of the Sydney Rock Oyster *Saccostrea commercialis* and the Pacific Oyster *Crassostreagigas*. Fishing Industry Research and Development Council Final Report. NSW Agriculture and Fisheries, Brackish Water Fish Culture Research Station, Salamander Bay, NSW.
16. Kennedy, V. S., Mihursky, J. A. (1971). Upper temperature tolerance of some estuarine bivalves. *Chesapeake Science***12** (4), 193-204.
17. Kooijman, S.A.L.M. (2000) Dynamic Energy and Mass Budgets in Biological Systems. Cambridge University Press, Cambridge (2000) 424 pp.
18. Kube, P., Cunningham, M., Dominik, S., Parkinson, S., Finn, B., Henshall, J., Bennett, R., Hamilton, M. (2011). *Enhancement of the Pacific oyster selective breeding program*. Hobart: FRDC and Seafood CRC Project.
19. Lawacz, W. (1977). Dissolved and Particulate Organic Carbon (DOC and POC) in Ecological Studies. *Acta Hydrochimica et Hydrobiologica***5**, 363-367.
20. Le Deuff, R. M., Nicolas, J.L., Renault, T., Cochenec, N. (1994). Experimental transmission of a herpes-like virus to axenic larvae of Pacific oyster, *Crassostreagigas*. *Bulletin of the European Association of Fish Pathologists***14**, 69-72.
21. Le Deuff, R. M., Renault, T., Gérard, A. (1996). Effects of temperature on herpes-like virus detection among hatchery-reared larval Pacific oyster *Crassostreagigas*. *Diseases of Aquatic Organisms***24**, 149-157.
22. Lorenzen, C.J. (1967) Determination of chlorophyll and phaeopigments: spectrophotometric equations. *Limnology and Oceanography***12** (2), 343-346.
23. Mazzola, A., Sarà, G. (2001). The effect of fish farming organic waste on food availability for bivalve mollusc (Gaeta Gulf, Central Tyrrhenian, MED): stable carbon isotopic analysis. *Aquaculture***192** (2-4), 361-379.

24. Mo, C., Neilson, B. (1994) Standardization of oyster soft tissue dry weight measurements. *Water research***28** (1), 243-246.
25. O'Meley, C. M. (1995). *Effects of Shell Abrasion and Aerial Exposure on the performance of Pacific Oysters Crassostreagigas (Thunberg, 1793) Cultured in Tasmania, Australia*. BSc (Hons), University of Tasmania.
26. Pernet, F., Barret, J., Le Gall, P., Corporeau, C., Dégremont, L., Lagarde, F., Pépin, J. F., Keck, N. (2012). Mass mortalities of Pacific oysters *Crassostreagigas* reflect infectious diseases and vary with farming practices in Mediterranean Thau lagoon, France. *Aquaculture Environment Interactions***2**, 215-237.
27. Pernet, F., Fuhrmann, M., Petton, B., Mazurié, J., Bouget, J-F., Fleury, E., Daigle, G., Gernez, P. (2018). Determination of risk factors for herpesvirus outbreak in oysters using a broad-scale spatial epidemiology framework. *Scientific Reports***8**.
28. Petton, B., Boudry, P., Alunno-Bruscia, M., Pernet, F. (2015). Factors influencing disease-induced mortality of Pacific oysters *Crassostreagigas*. *Aquaculture Environment Interactions***6**, 205-222.
29. Potter, M. A., Hill, B. J. (1982). Heat Mortality in the Sydney Rock Oyster, *Saccostrea (crassostrea) commercialis* and the effectiveness of some control methods. *Aquaculture***29**, 101-108.
30. Pouvreau, S., Bourles, Y., Lefebvre, S., Gangnery, A., Bruscia, M.A. (2006) Application of a dynamic energy budget model to the Pacific oyster, *Crassostreagigas*, reared under various environmental conditions. *Aquaculture***56**(2), 156-167.
31. Rankin, C., Moltsghaniwskyj, N., Morton, J., Wilkie, E. (2018). Shell shape and meat condition in selectively bred Sydney rock oysters, *Saccostrea glomerata* (Gould, 1850): The influence of grow-out methods. *Aquaculture Research***49** (3), 1189-1199.

32. Ren, J.S., Ross, A.H. (2001) A dynamic energy budget model of the Pacific oyster *Crassostreagigas*. *Ecological Modeling***142**, 105-120.
33. Robert, R., Trut, G., Borel, M., Maurer, D. (1993). Growth, fatness and gross biochemical composition of the Japanese oyster, *Crassostreagigas*, in Stanway cylinders in the Bay of Arcachon, France. *Aquaculture***110**, 249-261.
34. Sardegnaagricoltura. (2016) Sardegna Agricoltura, il sistema agricolo della Sardegna. [Online] Available from:  
[http://www.sardegnaagricoltura.it/documenti/14\\_43\\_20140613123850.pdf](http://www.sardegnaagricoltura.it/documenti/14_43_20140613123850.pdf)  
[Accessed: 01th November 2018].
35. Sauvage, C., Pépin, J.F., Lapègue, S., Boudry, P., Renault, T. (2009). Ostreid herpes virus 1 infection in families of the Pacific oyster, *Crassostreagigas*, during a summer mortality outbreak: difference in viral DNA detection and quantification using real-time PCR. *Virus Research***142**, 181-187.
36. Seed, R. (1968) Factors Influencing Shell Shape in the Mussel *Mytilus Edulis*. *Journal of the Marine Biological Association of the United Kingdom***48 (3)**, 561-584.
37. Shatkin, G., Shumway, S.E., Hawes, R. (1997) Considerations regarding the possible introduction of the pacific oyster (*Crassostreagigas*) to the gulf of maine: a review of global experience. *Journal of shellfish research***16 (2)**, 463-447.
38. Shellsim (2011) ShellSIM modelling shellfish growth [Online] Available from:  
[www.shellsim.com](http://www.shellsim.com) [Accessed: 01<sup>th</sup> November 2018].
39. Sheridan, A. K., Smith, I. R., Nell, J. A. (1996). Reducing the impact of environmental variation in a growth rate improvement program for the Sydney rock oyster *Saccostrea commercialis*. *Aquaculture***143**, 145-154.
40. Spencer-Davies, P. (1970). Physiological ecology of *Patella* IV. Environmental and limpet body temperatures. *Journal of the Marine Biological Association of the United Kingdom***50**, 1069-1077.

41. Taylor, K. E., (2001). Summarizing multiple aspects of model performance in a single diagram. *Journal of geophysical research***10**, 7183-7192.
42. Telesca, L., Michalek K., Sanders T., Peck L. S., Thyrring J., Harper E.M. (2018) Blue mussel shell shape plasticity and natural environments: a quantitative approach. *Scientific reports***8**, 1-15.
43. Van ErkomSchurink, C., Griffiths, C. L. (1993). Factors affecting relative rates of growth in four South African mussel species. *Aquaculture***109**, 257-273.
44. Walne, P. R., Mann, R. (1975) Growth and biochemical composition in *Ostrea edulis* and *Crassostrea gigas*. In Barnes H. (ed) *Proceeding of the ninth European marine biology symposium*, Aberdeen: Aberdeen University Press, pp. 587-607.
45. Watanabe, K., Kuwae, T. (2015). How organic carbon derived from multiple sources contributes to carbon sequestration processes in a shallow coastal system. *Global Change Biology***21**, 2612-2623.