

Complexity revealed in the greening of the Arctic

Reference

Myers-Smith, I.H., Kerby, J.T., Phoenix, G.K. *et al.* Complexity revealed in the greening of the Arctic. *Nat. Clim. Chang.* **10**, 106–117 (2020). <https://doi.org/10.1038/s41558-019-0688-1>

Weblink

<https://www.nature.com/articles/s41558-019-0688-1>

Isla H. Myers-Smith^{1*}, Jeffrey T. Kerby^{2,3*}, Gareth K. Phoenix⁴, Jarle W. Bjerke⁵, Howard E. Epstein⁶, Jakob J. Assmann^{1,7}, Christian John³, Laia Andreu-Hayles⁸, Sandra Angers-Blondin¹, Pieter S.A. Beck⁹, Logan T. Berner¹⁰, Uma S. Bhatt¹¹, Anne D. Bjorkman^{12,13}, Daan Blok¹⁴, Anders Bryn¹⁵, Casper T. Christiansen¹⁶, J. Hans C. Cornelissen¹⁷, Andrew M. Cunliffe¹⁸, Sarah C. Elmendorf¹⁹, Bruce C. Forbes²⁰, Scott J. Goetz¹⁰, Robert D. Hollister²¹, Rogier de Jong²², Michael M. Loranty²³, Marc Macias-Fauria²⁴, Kadmiel Maseyk²⁵, Signe Normand⁷, Johan Olofsson²⁶, Thomas C. Parker²⁷, Frans-Jan W. Parmentier^{28,29,30}, Eric Post³, Gabriela Schaepman-Strub³¹, Frode Stordal²⁸, Patrick F. Sullivan³², Haydn J. D. Thomas¹, Hans Tømmervik⁵, Rachael Treharne⁴, Craig E. Tweedie³³, Donald A. Walker³⁴, Martin Wilmking³⁵, Sonja Wipf³⁶

* Joint first authors

1. School of GeoSciences, University of Edinburgh, Edinburgh, EH9 3FF, UK

2. Neukom Institute for Computational Science and the Institute for Arctic Studies, Dartmouth College, 78 College Street, Hanover, NH, 03755, USA

3. Department of Wildlife, Fish, and Conservation Biology, University of California, Davis, One Shields Ave., Davis, CA, 95616, USA

4. Department of Animal and Plant Sciences, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK

5. Norwegian Institute for Nature Research, FRAM – High North Research Centre for Climate and the Environment, P.O. Box 6606 Langnes, NO-9296 Tromsø, Norway

6. Department of Environmental Sciences, University of Virginia, Charlottesville, VA, 22904-4123, USA
7. Section for Ecoinformatics and Biodiversity and Arctic Research Center, Department of Bioscience, University of Aarhus, Ny Munkegade 114, Building 1540, DK-8000 Aarhus C, Denmark
8. Lamont-Doherty Earth Observatory of Columbia University, 61 Route 9W, Palisades, NY, 10964, USA
9. European Commission, Joint Research Centre, 21027 Ispra VA, Italy
10. School of Informatics, Computing and Cyber Systems, Northern Arizona University, AZ, 86011, USA
11. Department of Atmospheric Sciences, University of Alaska Fairbanks, 2156 Koyukuk Drive, Fairbanks, AK, 99775-7320, USA
12. Senckenberg Gesellschaft für Naturforschung, Biodiversity and Climate Research Centre, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany
13. Department of Biological and Environmental Sciences, University of Gothenburg, PO Box 461, Gothenburg SE-405 30, Sweden
14. Netherlands Organisation for Scientific Research, Laan van Nieuw Oost-Indië 300, NL-2593 CE The Hague, The Netherlands
15. Natural History Museum, University of Oslo, PB 1172 Blindern, 0318 Oslo, Norway
16. NORCE Norwegian Research Centre & Bjerknes Centre for Climate Research, Jahnebakken 5, 5007 Bergen, Norway
17. Systems Ecology, Department of Ecological Science, Faculty of Science, Vrije Universiteit, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands
18. School of Geography, University of Exeter, Exeter, EX4 4RJ, UK
19. Institute for Arctic and Alpine Research, University of Colorado, Boulder, CO, 80303, USA
20. Arctic Centre, University of Lapland, Box 122, FIN-96101 Rovaniemi, Finland
21. Biology Department, Grand Valley State University, 1 Campus Drive, Allendale, MI, 49401, USA
22. Department of Geography, University of Zürich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

23. Department of Geography, Colgate University, 13 Oak Dr, Hamilton, NY, 13346, USA

24. School of Geography & the Environment, University of Oxford, S Parks Rd, Oxford, OX1 3QY,
UK

25. School of Environment, Earth and Ecosystem Sciences, The Open University, Milton Keynes,
MK7 6AA, UK

26. Umeå University, Department of Ecology and Environmental Sciences, KB.H4, Linnaeus
Väg 6, 90187 Umeå, Sweden

27. Biological and Environmental Sciences, School of Natural Sciences, University of Stirling,
Stirling, FK9 4LA, UK

28. Department of Geosciences, University of Oslo, Postboks 1022 Blindern, 0315 Oslo, Norway

29. Department of Physical Geography and Ecosystem Science, Lund University, Sölvegatan 12,
223 62 Lund, Sweden

30. Department of Arctic and Marine Biology, UiT-The Arctic University of Norway, N-9037,
Tromsø, Norway

31. Department of Evolutionary Biology and Environmental Studies, University of Zurich,
Winterthurerstr. 190, 8057 Zurich, Switzerland

32. Environment and Natural Resources Institute, University of Alaska Anchorage, 3211
Providence Dr., Anchorage, AK, 99508, USA

33. Department of Biological Sciences and the Environmental Science and Engineering Program,
University of Texas at El Paso, 500 W University Ave, El Paso, TX 79968, USA

34. Institute of Arctic Biology, University of Alaska Fairbanks, 2140 Koyukuk Drive, Fairbanks, AK,
99775, USA

35. Institute of Botany and Landscape Ecology, University Greifswald, Soldmannstrasse 15, D-
17487 Greifswald, Germany

36. WSL Institute for Snow and Avalanche Research SLF, Team Mountain Ecosystems,
Fluelastrasse 11, 7260 Davos Dorf, Switzerland

Editor's Summary

As tundra ecosystems respond to rapid Arctic warming, satellites records suggest a widespread greening. This Perspective highlights the challenges of interpreting complex Arctic greening trends and provides direction for future research by combining ecological and remote sensing approaches.

Abstract

As the Arctic warms, vegetation is responding and satellite measures indicate widespread greening at high latitudes. This 'greening of the Arctic' is among the world's most significant large-scale ecological responses to global climate change. However, a consensus is emerging that the underlying causes and future dynamics of so-called Arctic greening and browning trends are more complex, variable, and inherently scale dependent than previously thought. Here, we summarize the complexities of observing and interpreting high-latitude greening to identify key priorities for future research. Incorporating satellite and proximal remote sensing with *in-situ* data, while accounting for uncertainties and scale issues will advance the study of past, present, and future Arctic vegetation change.

Main text

The Arctic has warmed at more than twice the rate of the rest of the planet in recent decades^{1,2}. Over the past forty years, satellite-derived vegetation indices have indicated widespread change at high latitudes^{3–16}. Satellite records allow for the quantification of change in places that are otherwise unevenly sampled by *in-situ* ecological observations¹⁷. Positive trends in satellite-derived vegetation indices (often termed Arctic greening)¹⁵ are generally interpreted as signs of *in-situ* increases in vegetation height, biomass, cover and abundance^{5,18,19} associated with warming^{5,14}. In the most recent Intergovernmental Panel on Climate Change report, tundra vegetation change including greening trends derived from satellite records²⁰ was identified as one of the clearest examples of the terrestrial impacts of climate change. Large-scale vegetation-climate feedbacks at high latitudes associated with greening could alter global soil carbon storage and the surface energy budget^{21,22}. In recent years, slowing or reversal of apparent greening from satellite studies have been reported in some regions (sometimes termed Arctic browning)^{3,4,12,13,15,23,24}. This slowdown is seemingly at

odds with earlier responses to long-term warming trends^{3,25}. Research now indicates substantial heterogeneity in vegetation responses to climate change in the Arctic^{18,19,26,27}. However, the mechanistic links between satellite records and *in-situ* observations^{3,6,24} remain unclear due to conceptual and technical barriers in their analysis and combined interpretation.

A review of Arctic greening

The terms Arctic ‘greening’ and ‘browning’ can have different meanings in the remote sensing and ecology literatures. From a remote sensing perspective, ‘greening’ (hereafter spectral greening) generally refers to a positive trend^{4,5,7,8,10,13–15}, and ‘browning’ (hereafter spectral browning) generally refers to negative trend in satellite-derived vegetation indices^{3,4,12,13,15,23,24}. Less frequently, greening is also used to describe advances in the seasonal timing of these vegetation proxies^{4,28}. From a field-ecology perspective, greening (hereafter vegetation greening) and browning (hereafter vegetation browning) refer to field-observed changes in vegetation^{4,12,13,24}. Historically, the general terms greening and browning were thus used to describe both a proxy of vegetation change and/or vegetation change itself depending on context. This lack of precise usage causes conceptual misunderstandings about Arctic greening and attribution to the drivers of change. Here, we present the current understanding of Arctic spectral and vegetation greening and browning to lay the foundations for a consensus between the remote sensing and field ecology perspectives.

Vegetation indices as proxies of vegetation productivity

Long-term trends in global vegetation dynamics are most commonly quantified from time series of spectral vegetation indices derived from optical satellite imagery (Figure 1). These indices are designed to isolate signals of leaf area and green vegetation cover from background variation by emphasizing reflectance signatures in discrete regions of the radiometric spectrum^{6,29–32}. Common vegetation indices include the Normalized Difference Vegetation Index (NDVI, Figure 2), Enhanced Vegetation Index (EVI) and Soil Adjusted Vegetation Index (SAVI), among others^{33–35}. NDVI correlates with biophysical vegetation properties like Leaf Area Index (LAI) and the fraction of Absorbed Photosynthetically Active Radiation (fAPAR)^{14,36–39}. However, these vegetation indices were not developed in polar contexts⁴⁰ and are only proxies of photosynthetic activity rather than

143 direct measurements of biological productivity^{33,39,41}. NDVI is the most commonly used vegetation
144 index because it is simple to calculate with spectral bands monitored since the launch of early-
145 generation Earth-observing satellites in the 1970s (Figure 2) and is perhaps best defined as a
146 measure of above-ground vegetation greenness.

147
148 The longest-term openly-available NDVI datasets have been produced from satellite-based sensors
149 with broad spatial coverages and different sampling frequencies. The most common datasets
150 include: 1) the Advanced Very-High-Resolution Radiometer (AVHRR – 1982 to present) on board
151 NOAA satellites, 2) the Moderate-resolution Imaging Spectroradiometer (MODIS – 2000 to present)
152 on board NASA satellites, and 3) NASA-USGS Landsat sensors (1972 to present). Most studies of
153 long-term trends calculate annual measures of maximum NDVI to derive change over space and
154 time, though time-integrated approaches are also used^{30,42–44}. However, trends in NDVI data
155 produced from different satellite datasets or using different methods do not always correspond at a
156 given location^{6,45,46} (Figure 1a,c). Thus, it can be challenging to distinguish ecological change from
157 differences due to methods and sensor/platform-related issues when interpreting localised spectral
158 greening or browning signals (Table 1, Figure 2).

Ecological factors influencing greening and browning trends

The ecological processes underlying spectral greening or browning measured by satellites are diverse and may unfold across overlapping scales, extents and timeframes. In tundra ecosystems, vegetation changes linked to spectral greening could include: encroachment of vegetation on previously non-vegetated land surfaces^{18,47}, changes in community composition – such as tundra shrub expansion^{5,19,27}, and/or changes in plant traits such as height^{48,49}, leaf area, or phenology^{50–52}. Tall shrub tundra typically has a higher NDVI than other tundra plant types^{49,53,54}, and bare ground²⁹ has a much lower NDVI than vegetated tundra (Figure 2). Spectral browning could be related to a variety of factors including for example loss of photosynthetic foliage¹² or increases in bare ground cover due to permafrost thaw⁵⁵ (Figure 1). Thus, changes in the species composition, growth form and traits of plant communities can influence greening and browning trends.

Physical factors influencing greening and browning trends

Widespread non-biological changes in high-latitude ecosystems could confound and decouple spectral greening or browning trends from changes in plant productivity (Table 1). Land cover, topography, and associated soil moisture, surface water, land-surface disturbances and snow-melt dynamics can all influence the measured spectral greenness of landscapes^{56–63} and likely influence greening trends. For example, changes in the extent of summer snow patches⁶³, surface water⁶⁰ or surface soil moisture⁵⁹ that are often associated with landscape-scale topographic variation could influence the measured NDVI of the land surface. At high latitudes, optical satellite sensors are only effective for a short annual window due to the prolonged polar night, while low sun angles and persistent cloud cover reduce data quality in the summer season (Table 1). The unique physical properties of high-latitude ecosystems in addition to the constraints of polar remote sensing are often underemphasized in remote sensing studies of Arctic vegetation change.

Arctic browning and heterogeneity of spectral greening trends

Not all areas of the Arctic are spectrally greening (Figure 1), and in recent years spectral browning and heterogeneity of spectral greening trends have been highlighted^{3,4,12,13,23}. Ecological explanations for vegetation browning include for example the sudden loss of photosynthetically active foliage due to extreme climatic events^{64–67}, biological interactions (e.g., disease or herbivore outbreaks)^{68–70}, permafrost degradation^{23,55} (Figure 1), increases in standing dead biomass⁷¹, coastal erosion⁷², salt inundation⁷³, altered surface water hydrology^{74,75} or fire^{9,76,77}. Spectral browning, however, could be attributed to reduced productivity caused by adverse changes in growing conditions such as lower water availability, shorter growing seasons³ or nutrient limitation²⁷. Nonetheless, long-term spectral greening trends remain far more pervasive than spectral browning in tundra ecosystems. Figures vary from 42% greening and 2.5% browning from 1982 to 2014 in the GIMMS3g AVHRR dataset⁷⁸, 20% greening and 4% browning from 2000 to 2016 in Landsat data¹⁵ and estimates of 13% greening and 1% browning for the MODIS trends calculated for 1000 random points in the tundra polygon in Figure 1 from 2000 to 2018. At circumarctic scales, the magnitude, spatial variability, and proximal drivers of patterns and trends of spectral greening versus browning are not well understood.

Correspondence between satellite and ground-based observations

Evidence for correspondence among *in-situ* vegetation change and trends in satellite-derived vegetation indices is mixed^{47,79–81}. NDVI trends across satellite datasets do not necessarily directly correspond with one another^{6,9}, nor does any one sensor or vegetation index combination correspond directly with *in-situ* vegetation change⁴⁷. For example, NDVI has been related to interannual variation in radial shrub growth^{5,10,82}, yet how radial growth links to change in leaf area, aboveground biomass, or landscape measures of productivity is not always clear^{83–85} (Figure 3). AVHRR NDVI greening trends did not correspond with the lack of change observed with Landsat NDVI data and *in-situ* plant composition between 1984 and 2009 in North Eastern Alaska⁴⁷. Direct comparisons of productivity changes from vegetation cover estimates^{18,86}, biomass harvests⁵³ or shrub growth⁸⁷ are complicated by the lack of annual-resolution *in-situ* data and low sampling replication across the landscape. We attribute the mixed evidence for correspondence between *in-*

situ and satellite-derived measures of tundra vegetation change and greening to the complexities of existing terminology, challenges of interpretation of spectral vegetation indices at high latitudes, and the scaling issues as outlined below.

In addition to productivity analyses, changes in growing season length and advances in plant phenology have been documented using both satellite^{43,78,88–91} and ground-based datasets, and here also paired comparisons do not always correspond (Figure 4). Measures of longer growing seasons have been attributed to earlier snowmelt and/or earlier leaf emergence in spring⁹², and longer periods of photosynthetic activity or later snowfall in autumn⁹³. However, few studies have monitored both leaf emergence and senescence of tundra plants *in situ* and so far provide no evidence for an increasing growing period at specific sites^{94,95}. In addition, community-level analyses indicate shorter flowering season lengths around the tundra biome⁵⁰. Shifts in plant phenology with warming⁵⁰ could also be linked to changing species composition or diversity^{18,48,86}, thus influencing the phenological diversity across the landscape^{96,97}. Satellite records may not capture the ecological dynamics of vegetation phenology at high latitudes, as snow cover can obscure the plant seasonal signal and deciduous plants only make up a portion of the vegetated land cover. Thus, uncertainty remains whether satellite-derived changes in circumarctic phenology represent a longer snow-free period uncoupled from the vegetation response or an actual realized longer growing season of plants^{94,98–100}.

Clarifying the terminology

To distinguish spectral greening and browning events from longer-term trends, we propose clarified definitions of events and trends. For an individual pixel, we define the *spectral trend* as an increase or decrease in NDVI (or other spectral vegetation index) over decadal time scales and a *spectral event* as a temporal outlier in the vegetation index relative to the long-term trend. Trends should be determined using a Theil-Sen estimator or similar robust statistical test for analyses of satellite data^{30,101}. We define a *spectral greening trend* as an increase of the vegetation index over decadal time scales. *In situ*, we interpret a *vegetation greening trend* as improved conditions for photosynthesis, reduced resource limitation and/or positive responses to disturbance in plant

communities, resulting in greater aboveground biomass, leaf area, productivity or changes in plant community composition. We define a *spectral browning trend* as a decrease in the vegetation index over decadal time scales. A *vegetation browning trend* may correspond with an *in-situ* change in vegetation productivity due to plant dieback or loss of vegetation cover through biotic or abiotic disturbances. We define *spectral greening events* as short-term increases in vegetation index greenness that can be attributed to an ecological process such as revegetation of ground cover after fire and *spectral browning events* as short-term decreases in the vegetation index that can be attributed to a disturbance such as permafrost thaw or plant dieback. The definitions we propose here distinguish between slower acting climatic or biotic drivers of greening or browning trends versus event-driven changes caused by weather, biotic pulses, or other regional events such as fire.

Differentiating events and trends

In any measure of remotely sensed or field-based greening separate consideration of trends and events will increase ecological interpretability (Figure 5). Spectral greening and browning trends operate at any spatial scale, from localised patches to landscapes or even biome extents over decades. In contrast, spectral greening and browning events, such as those caused by vegetation dieback or rapid vegetation increase after disturbance, are often restricted to patch and regional scales over shorter durations. Events often have more limited extents relative to trends due to their proximal causes, like changes in herbivory or precipitation. Broader scale events are also possible (e.g. globally synchronized reductions in vegetation productivity caused by changes in insolation related to an intense volcanic eruption¹⁰²). Therefore, greening or browning events might be embedded within overall spectral greening or browning trends, both temporally and/or spatially, without necessarily driving them (Figure 5). Examining the trend direction, magnitude and variance around the fit over time can shape more detailed investigations into the ecological interpretation of Arctic spectral greening trends.

The influence of baselines and temporal sampling

The baseline to which we compare productivity change will influence our interpretation of trends¹⁰³. Spectral greening or browning trends and events may result in threshold changes where on-the-

ground productivity does not return to the longer-term baseline (Figure 5; e.g., pulse in recruitment at treeline¹⁰⁴ or shrubline¹⁰⁵ or a large fire⁷⁷). In both satellite datasets and field observations, the baseline conditions are often constrained by the limitations of data availability rather than any deliberately selected starting point⁶. The low temporal sampling frequency of a few days to a few weeks of many legacy remote-sensing datasets (e.g., AVHRR, MODIS, Landsat, etc.) also introduces temporal scale-dependent effects that may be magnified in Arctic systems (Table 1). For example, comparisons of phenology across latitudes can be less reliable at higher versus lower latitudes due to shorter growing seasons and therefore fewer satellite data collection points for use in change detection analyses^{42,88,89}. Metrics based on the annual maximum NDVI of a given pixel are more likely to be influenced by temporal sampling artefacts at high latitudes than those that integrate productivity estimates through time, such as the growing season integrated NDVI (GSINDVI)⁴², time-integrated NDVI (TiNDVI)⁴³ or early growing season integrated NDVI indices⁴⁴. Trends in either instance could be observed or not observed due to statistical reasons related to sample size and/or the strength or linearity of the trend. Thus, simple linear analyses of annual greenness metrics derived from satellite data may not always capture real-world ecological change (Figure 5).

Challenges in the interpretation of vegetation indices

In addition to the need for more clearly defined terms, challenges remain in the ecologically meaningful interpretation of long-term trends in optical satellite data, especially at high latitudes. The statistical relationship between a vegetation index and biomass, leaf area, phenology, or any other measures of productivity can vary due to a suite of intrinsic (e.g., sensor design, quality flagging algorithms), extrinsic (e.g., atmospheric conditions, sun angle, snow cover)^{6,106} and biological factors¹⁰⁷ (Table 1). For example, the centre wavelength and width of spectral bands (e.g., in the red or near-infrared) used to generate vegetation indices were designed for different purposes in different sensors (Figure 2). While the NDVI formula may be the same, the covered spectral wavelength ranges differ between different datasets¹⁰⁸ (Figure 2b). Thus, the datasets may be more or less sensitive to specific non-vegetative influences, such as atmospheric scattering or the magnitude of spectral mixing associated with non-vegetated surfaces⁵⁷. Spectral unmixing is the process of decomposing the spectral signature of a mixed pixel into the abundances of a set of endmember categories¹⁰⁹. Longer-term vegetation change is difficult to resolve from cross-sensor comparisons among different satellite datasets or even among intercalibrations of the same sensor type (Figure 1). For these reasons, caution is warranted when comparing vegetation indices derived from different satellite products or even versions of the same product with different atmospheric corrections, quality assessments, and spatial/temporal compositing approaches^{6,108}. Differences in NDVI signal processing are actively studied by the remote-sensing community (Table 1), but could be better accounted for or quantified in Arctic greening studies.

Nonlinearities in NDVI as a vegetation proxy

Direct interpretations of vegetation changes from spectral data are contingent on the local relationship between NDVI and *in-situ* vegetation. The statistical relationships between vegetation indices and measures of Arctic vegetation biomass are nonlinear^{29,110} (Figure 2). This nonlinearity presents challenges for trend interpretation that are illustrated in Figure 2a. Here, an absolute increase in biomass for a 'low biomass' community towards a 'moderate biomass' community would result in a positive NDVI trend, but that same absolute biomass increase from moderate to high biomass would show virtually no trend in NDVI due to saturation (Figure 2). Thus, the relationship to

common ecological variables like changes in biomass or shrub ring widths (Figure 4) can be obscured by nonlinearities. Because the greening and browning terms are tied to changes in vegetation proxies, rather than direct biological measures, a lack of correspondence could occur between remotely-sensed vegetation proxies and *in-situ* vegetation change (Figure 2, 4 and 5). Such potential discrepancies exemplify why caution should be used when interpreting linear trends in proxies like NDVI (Figure 1) that are nonlinearly related to vegetation productivity without the use of *in-situ* data to corroborate conclusions.

Scaling issues in Arctic greening analyses

Scale and hierarchies present a longstanding challenge in the interpretation of remotely-sensed vegetation proxies^{111–113} (Figure 5). All long-term vegetation proxy time series (Landsat, MODIS, AVHRR) spatially aggregate spectral data to pixels (i.e., grains) that span hundreds of square metres to tens of square kilometres. The spectral signatures of plants and non-vegetative features in a landscape are reduced to a single value. The loss of variability within pixels masks information useful for the attribution of greening signals to processes across ecological hierarchies from populations and communities to ecosystems (Table 1, Figure 3 and 5). For example, within a single AVHRR GIMMS3g pixel, a subselection of 1 x 1 km pixels are upscaled to 8 x 8 km³². Within this aggregated pixel, ecological contributions to spectral greening signals such as increased shrub cover on south-facing slopes or revegetation of drained lake beds may be mixed with browning signals from for example disturbances such as retrogressive thaw slumps or vegetation trampling by herbivores (Figure 1). High-latitude pixels may also contain shadows caused by low-sun angle, patchy snow- and/or cloud-cover (Table 1). Thus, the emergent time series from such a pixel describes no single vegetation dynamic or environmental factor, but rather their integrated spectral responses. Broad-scale patterns of spatial variability in greening and browning across pixels are also influenced by grain size¹¹³ (Figure 1, 2, 5). Finer resolution satellites such as Landsat can reduce, but not necessarily eliminate such spectral mixing¹⁵. However, the extent to which the sometimes-contradictory greening and browning signals found across different spectral datasets can be attributed to the influence of the scale of measurement is poorly understood.

Complexities of capturing phenology

Measuring landscape phenology with satellite data presents additional challenges to ecological interpretation of Arctic greening (Table 1). The variability of timing of satellite imagery from year to year particularly at high latitudes⁹¹ can confound measures of phenology (known as phenometrics). Cloud or fog cover is highly variable and sensitive to changing sea ice conditions in coastal Arctic sites⁴⁴. Seasonal variation in cloud and fog cover influences both data availability and image compositing approaches in many phenology products⁹¹. In addition, vegetation metrics from early spring are much more likely to be influenced by snow, standing water or low sun angle than those closer to peak biomass in mid- to late-summer^{8,54,59}. However, early spring is a critical period for establishing a baseline for curve fitting or thresholding used to derive phenometrics. Ultimately no phenometric is best suited to all Arctic environments or time periods¹¹⁴. Snow regimes and land cover variability differ annually and regionally and thus phenometrics using coarse-grain imagery integrate different abiotic and biotic signals at different points in space and time¹¹⁴. Phenological differences of days to weeks or even months can result from analyses using different methods and metrics for the same datasets at the same location¹¹⁵. These relative differences are of substantial ecological importance given the short growing seasons of the Arctic^{78,114} (Figure 4). Circumarctic analyses of vegetation indices generally indicate that phenological shifts in the spectral greenness of the land surface are widespread^{78,88–90}. However, the magnitude and extent of spatial and temporal scaling issues in high-latitude remotely-sensed phenology trends warrant further consideration and research¹¹².

Towards a consensus perspective on Arctic greening

The fields of remote sensing and field-based ecology will benefit from jointly addressing the complexities of interpreting spectral and vegetation greening and browning trends. Analyses from one satellite platform or one specific ecological context is not sufficient to disentangle Arctic greening complexity. The required next steps will be an integration of perspectives and approaches through existing and new international research efforts to address the following critical research gaps:

1. *Addressing scale issues by integrating proximal remote sensing and in-situ observations into circumarctic greening analyses*

Analyses of observations across scales will allow us to bridge the gap and improve our mechanistic understanding of the links between *in-situ* vegetation dynamics and broader remotely-sensed patterns and trends. New instruments for carrying out *in-situ* and proximal remote-sensing observations for comparison with satellite data are developing rapidly. However, we must urgently develop standardized field data collection protocols. In order to facilitate future synthesis, we need to incorporate data from long-term ecological monitoring^{12,18,86,94}, historical imagery¹¹⁶, phenocam networks¹¹⁷, flux towers¹¹⁸, high-resolution imagery such as from aircraft, towers, and drones¹¹⁹ and satellites.

2. *Incorporation of heterogeneity and uncertainty into analyses to improve confidence in detection of Arctic greening trends*

New finer spatial or temporal resolution data will inform analyses of historic greening trends. Current circumarctic Landsat analyses are shedding light on greening trends by exploiting finer spatial resolution data while accounting for the lower temporal resolution of observation records¹⁵. Recent and ongoing release of finer-resolution satellite datasets (e.g., EU-funded Sentinel missions, Digital Globe, Planet constellations) and data products (e.g., the Arctic Digital Elevation Model) will provide finer spatial (2-10 m) and/or temporal resolution (1-5 days) data across the Arctic¹²⁰. We can gain a better understanding of past spectral greening signals from legacy satellite datasets by conducting standardized reprocessing with for example statistical methods incorporating uncertainty in observations such as image quality information, improved atmospheric corrections and snow detection.

3. *Inclusion of new observational tools beyond optical vegetation indices to clarify the mechanistic links between spectral greening and vegetation change*

In addition to incorporating finer resolution datasets, new types of data collection can inform our understanding of what greening patterns and trends represent. Emerging remote sensing campaigns using hyperspectral sensors or those that can measure Solar-Induced Fluorescence (SIF)¹²¹ will

provide new insights into vegetation dynamics. However, future sensor development across satellite, aircraft and near-surface platforms should be designed to maximize comparability. In addition to new data collection, novel data integration approaches, for example those employing machine learning, will provide greater insights into biome-scale analyses linking remote sensing observations with ecological change in high-latitude ecosystems^{21,122}.

Conclusions

Recent research has highlighted the complexity in observed Arctic greening and browning trends. Although satellite data have been used to detect and attribute global change impacts and resulting climate feedbacks in Arctic ecosystems^{20,22}, numerous questions and uncertainties remain. The three major challenges in resolving these uncertainties are: 1) improving the clarity of the definitions of widely used terminology associated with greening and browning phenomena, 2) promoting the understanding of the strengths and limitations of vegetation indices when making ecological interpretations and, 3) better incorporating and accounting for different scales of observation and uncertainty in analyses of changing tundra productivity and phenology. New sensors and better access to legacy data are improving our ability to remotely sense vegetation change. However, new data alone will not provide solutions to many of the longstanding conceptual and technical challenges. The complexity of Arctic greening will only be fully understood through multidisciplinary efforts spanning the fields of ecology, remote sensing, earth system science and computer science. As a field, we need to look forwards to quantify contemporary and future change, but also backwards by conducting reanalyses of historical data. Ultimately, we urgently need a deeper understanding of the relationships between patterns and processes in greening and browning dynamics to improve estimates of the globally-significant climate change feedbacks in high-latitude ecosystems²⁰.

Table 1. A variety of geophysical^{13,106,123}, environmental^{44,60,61} and ecological^{12,47,49,54,57,110} factors can influence the magnitude and direction of change in vegetation indices and are particularly problematic at high latitudes⁶. The effects include: 1) Radiometric effects: differences among satellite datasets including band widths, atmospheric effects, cloud-screening algorithms, sensor degradation, orbital shift and bidirectional reflectance distribution functions originating from differences in field of view and sun geometries. 2) Spectral mixing: the blending of sub-pixel spatial heterogeneity that can influence the overall pixel signal (Figure 2). 3) Adjacency effects: the reflectance of surrounding pixels that can influence the signal of a given pixel (Figure 2). And, 4) a variety of environmental and ecological factors from snow melt and soil moisture dynamics to composition of evergreen versus deciduous or vascular versus non-vascular plants.

Factors influencing vegetation indices	Specific effects	Influence on apparent greening patterns and trends
Low sun angle	Radiometric effects	At high latitudes, low sun angles and cloud shadows can have a greater influence on vegetation indices relative to lower latitudes ⁶² . NDVI varies with sun angle, an effect magnified in spring and autumn ⁶² . Shadows also reduce NDVI and may be difficult to detect in coarse grained imagery ⁴⁴ .
Cloud cover	Radiometric effects, Spectral mixing, Adjacency effects	Thin cloud, fog and smoke can influence imagery, reducing NDVI. Cloud and fog are particularly problematic in coastal regions and can vary greatly between image acquisitions ⁴⁴ . Cloud-screening algorithms differ among satellite datasets (in part as a function of available spectral bands), and partly cloudy or hazy conditions are particularly difficult for screening algorithms to detect consistently. In addition, the foggiess of Arctic locations can vary over time due to changing temperatures ⁴⁴ and/or sea ice conditions ¹²⁴ .
Standing water	Spectral mixing, Adjacency effects	Standing water ⁶⁰ can influence comparisons of vegetation indices across space and may not be detectable in coarse-grained imagery, despite influencing spectral signatures. NDVI values of water are generally low, however shallow water or standing water intermixed with vegetation or algal growth may not be identified as water by quality filters and may have higher NDVI. Water within a pixel may lead to artificially low NDVI values and can influence estimates of NDVI change over time. This is especially relevant to the Arctic during the spring and summer as snow melts and turns into ephemeral ponds and lakes whose spectral signatures will be mixed with nearby vegetation ¹²⁵ . NDVI signals could be driven by changes in standing water over time associated with changing precipitation, permafrost conditions, and/or warming rather than by changes in vegetation ^{56,57,60,125,126} .
Snow patches	Spectral mixing, adjacency effects	Sub-pixel sized snow patches will decrease the NDVI for a given tundra area ⁵⁷ . NDVI values of snow are strongly negative. Earlier snow loss or later snow return may drive a strong positive trend in NDVI. Longer persistence of snow on the landscape in patches may not be filtered by quality algorithms, yet could still lead to lower NDVI values.
	Snow versus phenology dynamics	Surface reflectance just after snow off is commonly used as the baseline when fitting phenology models. This approach masks the effects of sub-nivean phenological progression and/or may overemphasise the role of snow-off or snow-on dates as a driver of plant phenology ^{57,63} .

Soil moisture	Spectral mixing	Soil moisture can influence the reflectance of vegetated tundra surfaces ^{58,59} . NDVI values are sensitive to soil moisture, which may or may not covary with vegetation change ¹²⁵ . Furthermore, NDVI is relatively insensitive to changes in very sparsely vegetated (e.g., the High Arctic ¹²⁷) and very densely vegetated (e.g., forest or shrubland ¹²⁸) environments.
	Plant water content	Mosses can absorb water and thus influence surface reflectance of landscapes independent of vascular plant phenology and productivity ¹²⁶ .
Short growing season	Timing of image acquisition	Trends in NDVI metrics and growing season length can be influenced by the timing of data acquisition. To compare spatial patterns in vegetation indices among sites, images are required from the same time within the growing season and the same time points within the day ¹²⁶ . However, the short growing seasons at high latitudes make image acquisition particularly challenging. Satellites have different temporal frequencies for overpasses thus influencing comparisons. Growing season length decreases at higher latitudes, thus the impact of missing data is of a greater magnitude as latitude increases.
Rapid plant phenology	Chosen phenometric	The specific metrics used to quantify phenology will influence the resulting patterns observed ⁹¹ . Combining datasets with different spatial and temporal resolutions can limit comparisons (Figure 2). Variation in phenology metrics due to curve-fitting methods can exceed variation in measured phenology signals. Thus, using the same phenological functions across large geographic and ecological gradients, such as across the high latitudes, may introduce biases and/or errors.
	Phenological diversity	Changes in phenology of individual species or plants growing in particular microclimates can lead to shifts in landscape phenology ⁵⁰ .
Plant traits and functional groups or types	Isolating changes in plant productivity and canopy structure versus composition	Vegetation indices are related to radiation absorbed by green foliage (APAR), canopy structure, species composition, leaf-level traits and biomass ^{37,39} (Figure 2). However, how vegetation indices and ecological properties covary across diverse Arctic ecosystems is not well established. Other factors including bare ground cover, canopy structure, etc. that influence vegetation indices must be accounted for to isolate productivity change from other land surface changes.
	Vascular and deciduous versus non-vascular and evergreen plants	Non-vascular or evergreen plants can obscure the deciduous vascular plant seasonal signal ^{49,81} . Tundra without vascular plants can additionally have a substantial cover of biological soil crust communities consisting of lichens, cyanobacteria, mosses and green algae that may also influence NDVI ^{107,126} .


Satellite records indicate greening trends across the circumpolar Arctic

Figure 1. Arctic greening, which varies across space and time and among satellite datasets, is driven by both actual in-situ change and, in part, by the challenges of satellite data interpretation and integration. Trends in maximum NDVI vary spatiotemporally and the magnitude of changes is different depending on what satellite imagery is analysed (a and c, data subsetted to temporally overlapping years; b and d, GIMMS3gv1 1982 to 2015 and MODIS MOD13A1v6 2000 to 2018). Regional trends may summarise localised greening, for example shrub encroachment (e) and browning such as permafrost thaw (g) occurring at the pixel scale on Qikiqtaruk - Herschel Island in the Canadian Arctic (f). NDVI trends (a and c) were calculated using robust regression (Theil-Sen estimator) in the Google Earth Engine¹²⁹. Dashed line indicates the Arctic Circle and the black outlined polygon (a and c) and green 'Tundra' line (b and d) indicates the Arctic tundra region from the Circumpolar Arctic Vegetation Map (www.geobotany.uaf.edu/cavm/). The inset map in d indicates the regions for the mean trends for yellow 'Eurasia' and blue 'North America' polygons.

NDVI can vary across datasets due to NDVI biomass relationships, bandwidths of sensors and data quality issues




Figure 2. Ecological interpretation of trends in the Normalized Difference Vegetation Index

(NDVI) requires a consideration of non-ecological factors. NDVI, calculated as the difference between red and near infrared bands (NIR), has a non-linear relationship with several common metrics of plant productivity, like biomass and LAI (a). Satellite platforms have different spectral band widths which can influence calculations of NDVI despite shared centre wavelengths (b). NDVI values from commonly available satellite data products and drone datasets (c) differed substantially across products and across plots of three different vegetation types (e) during the period of peak biomass in 2017 on Qikiqtaruk – Herschel Island, Yukon. Here, factors such as a lack of atmospheric correction (f), cloud or fog contamination (g), sub-pixel mixing (h), different plot grain sizes of data in more or less heterogeneous vegetation cover and timing of data acquisition could have all influenced NDVI values. Data were analysed and extracted for 30 x 30 m plots from 13th July to 4th August in 2017 using the Google Earth Engine¹²⁹ for the MODIS MYD13A1v6 (pixel size = 500 m x 500 m) and Landsat 8 (pixel size = 30 m x 30 m) NDVI product, and the top-of-atmosphere Sentinel-2 NDVI product without atmospheric corrections (pixel size = 10 m x 10 m) NDVI, and Pix4D-processed drone data collected using a radiometrically calibrated four-band multispectral sensor (Sequoia, pixel size = 12 cm x 12 cm) on an FX-61 fixed-wing platform with the High-latitude Drone Ecology Network protocols (<https://arcticdrones.org/>). We purposefully present data with quality and processing issues above to highlight the challenges in quantifying NDVI in regional-to-global studies where data quality issues may be spatially or temporally variable among locations.

Spatial heterogeneity in landcover can influence NDVI ~ vegetation relationships

Qikiqtaruk, Canada – low landscape-level heterogeneity and increasing shrub abundance and variable radial growth




Figure 3. Sub-pixel spatial heterogeneity in vegetative greening and browning cannot be accurately captured at coarser grains. Landscape patterns (a, e), trends (b, f), and variability (d, h) in NDVI may not represent *in-situ* observations of vegetation change. NDVI trends and interannual variability had mixed correspondence with increases in shrub abundance (c, g) and interannual variability in shrub growth on Qikiqtaruk – Herschel Island, Yukon⁹⁴ (c, point framing in twelve 1-m² plots; d, *Salix pulchra* = 21, <https://github.com/ShrubHub/QikiqtarukHub>) and Kangerlussuaq, Greenland^{84,130} (g, 13 0.25-m² plots; h, *Betula nana* = 42, *Salix glauca* = 32, <https://arcticdata.io/catalog/view/doi:10.18739/A24X0Q>, <https://arcticdata.io/catalog/view/doi:10.18739/A28Q18>, <https://arcticdata.io/catalog/view/doi:10.5065/D6542KRH>). Errors are standard error bars around mean values (c, g) and 95% credible intervals for a Bayesian hierarchical model of the relationship between detrended annual growth rings and NDVI with shrub individual and year as random effects (d, h). Detrending was done using a spline fit from the dplR package in R. Credible intervals for model slopes overlapped with zero (d, h). Marginal R^2 values indicate the variance in detrended ring widths explained by detrended NDVI (d, h). Landscape NDVI patterns (a and e) were measured using a Parrot Sequoia and FX-61 fixed wing platform according to High-latitude Drone Ecology Network protocols in the summer of 2017 (<https://arcticdrones.org/>) and analysed using the Pix4D

489 software. Coarser-grain NDVI time series (MODIS MOD13A1v6, 500m pixels) were calculated using
490 Google Earth Engine¹²⁹ and the Phenex package in R.



Plant phenology does not always match land-surface greenness across the growing season

Figure 4. Satellite-derived phenology estimates do not always match with in-situ plant phenology observations. Satellite-observed snow-free season length of the land surface (here defined as the period with NDVI greater than 50% of the max NDVI, b and c) might not directly correspond to the growing season of vascular plants in tundra ecosystems, particularly in autumn (a). Snow-melt dynamics can obscure the plant phenology signal and non-vascular or evergreen plants can obscure the deciduous vascular plant seasonal signal. Plant phenology data were collected at 20 monitoring plots on Qikiqtaruk-Herschel Island for the species *Salix arctica*, which makes up approximately 30% of the cover in the grass- and forb-dominated vegetation type. Analyses indicate that both leaf emergence and senescence have become earlier, resulting in no change in realized growing season length despite substantial increases in the snow-free period of the land surface⁹⁴ (a – c, <https://github.com/ShrubHub/QikiqtarukHub>). Satellite data are MODIS MOD13A1v6 extracted for the pixel containing the phenology transects using Google Earth Engine¹²⁹ and the Phenex package in R (b and c).

Greening and browning complexity can be addressed by incorporating scale and clarifying ambiguity in terminology

a. Arctic greening patterns and trends are influenced by issues of scale



b. Spatial heterogeneity in NDVI greening/browning patterns can influence greening/browning trends over time




Figure 5. Arctic greening is influenced by both issues of measurement scale and inference across ecological hierarchies. Spectral resolution (Figure 2), extent (Figure 1), spatial resolution (Figure 2), landscape-level heterogeneity (Figure 3), temporal resolution (Figure 4), and ecological factors all influence the interpretation of greening trends (a). Within-pixel changes in land surface greening and browning events and trends can translate into different greening and browning patterns as their effects are scaled up (b). Ecological processes that comprise greening and browning trends include a combination of events, such as a pulse of plant recruitment or growth, a dieback of plants due to an extreme winter climate event, herbivore or disease outbreak or other disturbance and subsequent recovery. Longer-term change such as increasing shrub cover or progression of permafrost disturbances can also influence real-world NDVI time series. These different factors add

516 complexity to the interpretation of Arctic greening trends. The scale and hierarchy of observations
517 need to be incorporated into and/or accounted for in future analyses of Arctic greening.

Corresponding authors

Correspondence to Isla Myers-Smith (isla.myers-smith@ed.ac.uk) and Jeffrey Kerby (jeffrey.t.kerby@dartmouth.edu).

Author Contributions

IHM-S and JTK conducted the analyses and wrote the manuscript with contributions from all authors. GKP, JWB and HE contributed substantially to early versions of the manuscript. IHM-S, JTK, JJA, AMC, CJ, SA-B, HJDT and ESP collected drone and *in-situ* data. This paper results from two collaborations: the sTundra working group at the German Centre for Integrative Biodiversity Research (iDiv) led by IHM-S, SCE and ADB and the ‘Event Drivers of Arctic Browning Workshop’ at the University of Sheffield led by GKP.

Funding

Data collection on Qikiqtaruk-Herschel Island was funded by the UK Natural Environment Research Council (NERC) NE/M016323/1 [to IMS] and a National Geographic Society grant CP-061R-17 and a Parrot Climate Innovation Grant [to JTK]. Data collection at Kangerlussuaq, Greenland was supported by the US National Science Foundation (NSF) grants 0724711, 0713994, 0732168, 0902125, 1107381, 1525636, 1748052 and the National Geographic Society [to EP], as well as an Arctic Institute of North America Grant-in-Aid [to CJ]. The sTundra working group was supported by sDiv, the Synthesis Centre of the German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig (DFG FZT 118). The Event Drivers of Arctic Browning workshop was funded by P3-Plant Production and Protection (<http://p3.sheffield.ac.uk/>). Several members of the team are supported by the NASA ABoVE program (<https://above.nasa.gov/>). Additional funding was provided by the Research Council of Norway grant 287402 [to JWB and HT] and 294948 [to FS, JWB, AB, HT, and FJWP], the NERC doctoral training partnership grant NE/L002558/1 [to JJA and HJDT], the US NSF grants PLR-1504134, AGS 15-02150, PLR16-03473 [to LAH], the Natural Sciences and Engineering Research Council of Canada and the Canadian Centennial Scholarship Fund [to SAB], the Academy of Finland decision 256991 and JPI Climate 291581 [to BCF], the NASA ABoVE grants NNX17AE44G and NNX17AE13G [to SJG & LTB], NSF grants PLR-0632263, PLR-0856516, PLR-

1432277, PLR-1504224, PLR-1836839 [to RDH], the US NSF grant PLR-1417745 [to MML], an NERC IRF NE/L011859/1 [to MMF], Independent Research Fund Denmark 7027-00133B and Villum Fonden VKR023456 to [SN], the Norwegian Research Council grants 230970 and 274711 and the Swedish Research Council registration 2017-05268 [to FJWP], University of Zurich Research Priority Program on Global Change and Biodiversity [to GS-S] and the US NSF grants OPP-1108425 and PLR-1108425 [to PFS].

Acknowledgements

We thank John Gammon and Matthias Forkel for their very thoughtful and constructive reviews of the manuscript. We thank the Inuvialuit and Greenlandic People for the opportunity to conduct field research on their land.

Data availability

Data come from publicly available remote sensing and ecological datasets including:

MODIS (<https://modis.gsfc.nasa.gov/>), GIMMS3g.v1 (<https://nex.nasa.gov/nex/projects/1349/>), the High Latitude Drone Ecology Network (<https://arcticdrones.org/>), shrub abundance, annual growth ring and phenology datasets (<https://github.com/ShrubHub/QikiqtarukHub>, <https://arcticdata.io/catalog/view/doi:10.18739/A24X0Q>, <https://arcticdata.io/catalog/view/doi:10.18739/A28Q18>, <https://arcticdata.io/catalog/view/doi:10.5065/D6542KRH>).

Code availability

Code is available in a GitHub repository (<https://github.com/ShrubHub/GreeningHub>).

References

1. IPCC. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. (2013).

2. Post, E. et al. The polar regions in a 2°C warmer world. *Science Advances* **5**, eaaw9883 (2019).

Reviews how Arctic warming has outpaced the global average with wide reaching consequences for tundra ecosystems.

3. Bhatt, U. S. et al. Recent declines in warming and vegetation greening trends over Pan-Arctic tundra. *Remote Sens.* **5**, 4229–4254 (2013).

Reports of declines in spectral greening trends across Pan-Arctic tundra ecosystems.

4. de Jong, R., de Bruin, S., de Wit, A., Schaepman, M. E. & Dent, D. L. Analysis of monotonic greening and browning trends from global NDVI time-series. *Remote Sens. Environ.* **115**, 692–702 (2011).

5. Forbes, B. C., Macias-Fauria, M. & Zetterberg, P. Russian arctic warming and ‘greening’ are closely tracked by tundra shrub willows. *Glob. Change Biol.* **16**, 1542–1554 (2010).

Links spectral greening with both warming and tundra shrub growth in the Russian Arctic.

6. Guay, K. C. et al. Vegetation productivity patterns at high northern latitudes: a multi-sensor satellite data assessment. *Glob. Change Biol.* **20**, 3147–3158 (2014).

Reports conflicting patterns in spectral greening trends from different satellite datasets highlighting critical uncertainties in the links between satellite-observed spectral greening trends and *in situ* vegetation change.

7. Jia, G. J., Epstein, H. E. & Walker, D. A. Greening of arctic Alaska, 1981–2001. *Geophys. Res. Lett.* **30**, HLS 3-1 (2003).

8. Jia, G. J., Epstein, H. E. & Walker, D. A. Vegetation greening in the Canadian Arctic related to decadal warming. *J. Environ. Monit.* **11**, 2231 (2009).

9. Ju, J. & Masek, J. G. The vegetation greenness trend in Canada and US Alaska from 1984–2012 Landsat data. *Remote Sens. Environ.* **176**, 1–16 (2016).

- 597 10. Macias-Fauria, M., Forbes, B. C., Zetterberg, P. & Kumpula, T. Eurasian Arctic greening
598 reveals teleconnections and the potential for structurally novel ecosystems. *Nature Clim.*
599 *Change* **2**, 613–618 (2012).
- 600 11. Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G. & Nemani, R. R. Increased plant growth
601 in the northern high latitudes from 1981 to 1991. *Nature* **386**, 698–702 (1997).
- 602 **One of the first studies to report spectral greening trends for Northern high-latitude**
603 **ecosystems.**
- 604 12. Phoenix, G. K. & Bjerke, J. W. Arctic browning: extreme events and trends reversing arctic
605 greening. *Glob. Change Biol.* **22**, 2960–2962 (2016).
- 606 **Reviews Arctic browning from an ecological perspective capturing the differences between**
607 **events and trends in spectral greening and browning.**
- 608 13. Zhang, Y., Song, C., Band, L. E., Sun, G. & Li, J. Reanalysis of global terrestrial vegetation
609 trends from MODIS products: Browning or greening? *Remote Sens. Environ.* **191**, 145–155
610 (2017).
- 611 14. Zhu, Z. et al. Greening of the Earth and its drivers. *Nature Clim. Change* **6**, 791–795 (2016).
- 612 15. National Academies of Sciences, E. Understanding Northern Latitude Vegetation Greening
613 and Browning: Proceedings of a Workshop. (2019). doi:10.17226/25423.
- 614 16. Huang, M. et al. Velocity of change in vegetation productivity over northern high latitudes. *Nat.*
615 *Ecol. Evol.* **1**, 1649 (2017).
- 616 17. Metcalfe, D. B. et al. Patchy field sampling biases understanding of climate change impacts
617 across the Arctic. *Nat. Ecol. Evol.* **2**, 1443 (2018).
- 618 18. Elmendorf, S. C. et al. Plot-scale evidence of tundra vegetation change and links to recent
619 summer warming. *Nature Clim. Change* **2**, 453–457 (2012).
- 620 19. Myers-Smith, I. H. et al. Shrub expansion in tundra ecosystems: dynamics, impacts and
621 research priorities. *Environ. Res. Lett.* **6**, 045509 (2011).
- 622 20. IPCC Working Group II. Climate Change 2014: Impacts, Adaptation, and Vulnerability. (2014).
- 623 21. Forkel, M. et al. Enhanced seasonal CO₂ exchange caused by amplified plant productivity in
624 northern ecosystems. *Science* **351**, 696–699 (2016).

22. Pearson, R. G. et al. Shifts in Arctic vegetation and associated feedbacks under climate change. *Nature Clim. Change* **3**, 673–677 (2013).
23. Lara, M. J., Nitze, I., Grosse, G., Martin, P. & McGuire, A. D. Reduced arctic tundra productivity linked with landform and climate change interactions. *Sci. Rep.* **8**, 2345 (2018).
24. Miles, V. V. & Esau, I. Spatial heterogeneity of greening and browning between and within bioclimatic zones in northern West Siberia. *Environ. Res. Lett.* **11**, 115002 (2016).
25. Piao, S. et al. Evidence for a weakening relationship between interannual temperature variability and northern vegetation activity. *Nat. Commun.* **5**, 5018 (2014).
26. Bjorkman, A. D. et al. Status and trends in Arctic vegetation: Evidence from experimental warming and long-term monitoring. *Ambio* (2019) doi:10.1007/s13280-019-01161-6.
27. Martin, A. C., Jeffers, E. S., Petrokofsky, G., Myers-Smith, I. & Macias-Fauria, M. Shrub growth and expansion in the Arctic tundra: an assessment of controlling factors using an evidence-based approach. *Environ. Res. Lett.* **12**, 085007 (2017).
28. Piao, S., Friedlingstein, P., Ciais, P., Viovy, N. & Demarty, J. Growing season extension and its impact on terrestrial carbon cycle in the Northern Hemisphere over the past two decades. *Global Biogeochem. Cycles* **21**, GB3018 (2007).
29. Reynolds, M. K., Walker, D. A. & Maier, H. A. NDVI patterns and phytomass distribution in the circumpolar Arctic. *Remote Sens. Environ.* **102**, 271–281 (2006).
30. Forkel, M. et al. Trend change detection in NDVI time series: Effects of inter-annual variability and methodology. *Remote Sens.* **5**, 2113–2144 (2013).
- Highlights how the analyzed satellite dataset, the corresponding spatiotemporal resolution, and the applied statistical method influence the detection of spectral greening trends.**
31. Tucker, C. J. et al. An extended AVHRR 8-km NDVI dataset compatible with MODIS and SPOT vegetation NDVI data. *Int. J. Remote Sens.* **26**, 4485–4498 (2005).
32. Pinzon, J. E. & Tucker, C. J. A non-stationary 1981–2012 AVHRR NDVI3g time series. *Remote Sens.* **6**, 6929–6960 (2014).
33. Huete, A. et al. Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sens. Environ.* **83**, 195–213 (2002).

- 653 34. Brown, T. B. et al. Using phenocams to monitor our changing Earth: toward a global phenocam
654 network. *Front. Ecol. Environ.* **14**, 84–93 (2016).
- 655 35. Myneni, R. B., Hall, F. G., Sellers, P. J. & Marshak, A. L. The interpretation of spectral
656 vegetation indexes. *IEEE Trans. Geosci. Remote Sens.* **33**, 481–486 (1995).
- 657 36. Forkel, M. et al. Codominant water control on global interannual variability and trends in land
658 surface phenology and greenness. *Glob. Change Biol.* **21**, 3414–3435 (2015).
- 659 37. Myneni, R. B. & Williams, D. L. On the relationship between FAPAR and NDVI. *Remote Sens.*
660 *Environ.* **49**, 200–211 (1994).
- 661 38. Zhu, Z. et al. Global data sets of vegetation leaf area index (LAI)_{3g} and fraction of
662 photosynthetically active radiation (FPAR)_{3g} derived from global inventory modeling and
663 mapping studies (GIMMS) normalized difference vegetation index (NDVI_{3g}) for the period
664 1981 to 2011. *Remote Sens.* **5**, 927–948 (2013).
- 665 39. Sellers, P. J. Canopy reflectance, photosynthesis, and transpiration, II. The role of biophysics
666 in the linearity of their interdependence. *Remote Sens. Environ.* **21**, 143–183 (1987).
- 667 40. Benedetti, R. & Rossini, P. On the use of NDVI profiles as a tool for agricultural statistics: The
668 case study of wheat yield estimate and forecast in Emilia Romagna. *Remote Sens. Environ.*
669 **45**, 311–326 (1993).
- 670 41. Goetz, S. J. & Prince, S. D. Modelling terrestrial carbon exchange and storage: Evidence and
671 implications of functional convergence in light-use efficiency. in *Advances in Ecological*
672 *Research* (eds. Fitter, A. H. & Raffaelli, D.) vol. 28 57–92 (Academic Press, 1999).
- 673 42. Wang, L. & Fensholt, R. Temporal changes in coupled vegetation phenology and productivity
674 are biome-specific in the Northern Hemisphere. *Remote Sens.* **9**, 1277 (2017).
- 675 43. Bhatt, U. S. et al. Changing seasonality of panarctic tundra vegetation in relationship to climatic
676 variables. *Environ. Res. Lett.* **12**, 055003 (2017).
- 677 44. Karlsen, S. R., Anderson, H. B., Wal, R. van der & Hansen, B. B. A new NDVI measure that
678 overcomes data sparsity in cloud-covered regions predicts annual variation in ground-based
679 estimates of high arctic plant productivity. *Environ. Res. Lett.* **13**, 025011 (2018).

- 680 45. Fensholt, R. & Proud, S. R. Evaluation of Earth Observation based global long term vegetation
681 trends — Comparing GIMMS and MODIS global NDVI time series. *Remote Sens. Environ.*
682 **119**, 131–147 (2012).
- 683 46. Tian, F. et al. Evaluating temporal consistency of long-term global NDVI datasets for trend
684 analysis. *Remote Sens. Environ.* **163**, 326–340 (2015).
- 685 47. Pattison, R. R., Jorgenson, J. C., Raynolds, M. K. & Welker, J. M. Trends in NDVI and tundra
686 community composition in the Arctic of NE Alaska between 1984 and 2009. *Ecosystems* **18**,
687 707–719 (2015).
- 688 48. Bjorkman, A. D. et al. Plant functional trait change across a warming tundra biome. *Nature*
689 **562**, 57–62 (2018).
- 690 49. Boelman, N. T., Gough, L., McLaren, J. R. & Greaves, H. Does NDVI reflect variation in the
691 structural attributes associated with increasing shrub dominance in arctic tundra? *Environ.*
692 *Res. Lett.* **6**, 035501 (2011).
- 693 50. Prevéy, J. S. et al. Warming shortens flowering seasons of tundra plant communities. *Nat.*
694 *Ecol. Evol.* **3**, 45 (2019).
- 695 51. Oberbauer, S. F. et al. Phenological response of tundra plants to background climate variation
696 tested using the International Tundra Experiment. *Phil. Trans. R. Soc. B* **368**, (2013).
- 697 52. Post, E., Steinman, B. A. & Mann, M. E. Acceleration of phenological advance and warming
698 with latitude over the past century. *Sci. Rep.* **8**, 3927 (2018).
- 699 53. Berner, L. T., Jantz, P., Tape, K. D. & Goetz, S. J. Tundra plant above-ground biomass and
700 shrub dominance mapped across the North Slope of Alaska. *Environ. Res. Lett.* **13**, 035002
701 (2018).
- 702 54. Blok, D. et al. The response of Arctic vegetation to the summer climate: relation between shrub
703 cover, NDVI, surface albedo and temperature. *Environ. Res. Lett.* **6**, 035502 (2011).
- 704 55. Grosse, G., Goetz, S., McGuire, A. D., Romanovsky, V. E. & Schuur, E. A. G. Changing
705 permafrost in a warming world and feedbacks to the Earth system. *Environ. Res. Lett.* **11**,
706 040201 (2016).
- 707 56. Brown, R., Derksen, C. & Wang, L. Assessment of spring snow cover duration variability over
708 northern Canada from satellite datasets. *Remote Sens. Environ.* **111**, 367–381 (2007).

57. Gamon, J. A., Huemmrich, K. F., Stone, R. S. & Tweedie, C. E. Spatial and temporal variation in primary productivity (NDVI) of coastal Alaskan tundra: Decreased vegetation growth following earlier snowmelt. *Remote Sens. Environ.* **129**, 144–153 (2013).

Highlights how spectral greening trends can be associated with soil moisture and growing degree days rather than earlier snowmelt emphasizing the complexity in Arctic greening and environmental change.

58. Goswami Santonu, Gamon John A. & Tweedie Craig E. Surface hydrology of an arctic ecosystem: Multiscale analysis of a flooding and draining experiment using spectral reflectance. *J. Geophys. Res. Biogeosciences* **116**, (2011).

59. Liu, N., Budkewitsch, P. & Treitz, P. Examining spectral reflectance features related to Arctic percent vegetation cover: Implications for hyperspectral remote sensing of Arctic tundra. *Remote Sens. Environ.* **192**, 58–72 (2017).

60. Reynolds, M. K. & Walker, D. A. Increased wetness confounds Landsat-derived NDVI trends in the central Alaska North Slope region, 1985–2011. *Environ. Res. Lett.* **11**, 085004 (2016).

61. Riihimäki, H., Heiskanen, J. & Luoto, M. The effect of topography on arctic-alpine aboveground biomass and NDVI patterns. *Int. J. Appl. Earth Obs.* **56**, 44–53 (2017).

62. Stow, D. A. et al. Remote sensing of vegetation and land-cover change in Arctic Tundra Ecosystems. *Remote Sens. Environ.* **89**, 281–308 (2004).

Highlights the issues with satellite records of spectral greening due to cloud contamination, low sun angle, surface water and other factors.

63. Wang, S. et al. Limitations and challenges of MODIS-derived phenological metrics across different landscapes in pan-Arctic regions. *Remote Sens.* **10**, 1784 (2018).

64. Bjerke, J. W. et al. Record-low primary productivity and high plant damage in the Nordic Arctic Region in 2012 caused by multiple weather events and pest outbreaks. *Environ. Res. Lett.* **9**, 084006 (2014).

65. Bokhorst, S. et al. Impacts of extreme winter warming in the sub-arctic: growing season responses of dwarf shrub heathland. *Glob. Change Biol.* **14**, 2603–2612 (2008).

66. Richardson, A. D. et al. Ecosystem warming extends vegetation activity but heightens vulnerability to cold temperatures. *Nature* **560**, 368 (2018).

67. Bjerke, J. W. et al. Understanding the drivers of extensive plant damage in boreal and Arctic ecosystems: Insights from field surveys in the aftermath of damage. *Sci. Total Environ.* **599–600**, 1965–1976 (2017).
68. Jepsen, J. U. et al. Ecosystem impacts of a range expanding forest defoliator at the forest-tundra ecotone. *Ecosystems* **16**, 561–575 (2013).
69. Lund, M. et al. Larval outbreaks in West Greenland: Instant and subsequent effects on tundra ecosystem productivity and CO₂ exchange. *Ambio* **46**, 26–38 (2017).
70. Post, E. S., Pedersen, C., Wilmers, C. C. & Forchhammer, M. C. Phenological sequences reveal aggregate life history response to climatic warming. *Ecology* **89**, 363–370 (2008).
71. Johnson, D. R. et al. Exclusion of brown lemmings reduces vascular plant cover and biomass in Arctic coastal tundra: resampling of a 50 + year herbivore exclosure experiment near Barrow, Alaska. *Environ. Res. Lett.* **6**, 045507 (2011).
72. Fritz, M., Vonk, J. E. & Lantuit, H. Collapsing Arctic coastlines. *Nature Clim. Change* (2017) doi:10.1038/nclimate3188.
73. Lantz, T. C., Kokelj, S. V. & Fraser, R. H. Ecological recovery in an Arctic delta following widespread saline incursion. *Ecol. Appl.* **25**, 172–185 (2015).
74. Nitze, I. et al. Landsat-based trend analysis of lake dynamics across northern permafrost regions. *Remote Sens.* **9**, 640 (2017).
75. Smith, L. C., Sheng, Y., MacDonald, G. M. & Hinzman, L. D. Disappearing Arctic Lakes. *Science* **308**, 1429–1429 (2005).
76. Rocha, A. V. et al. The footprint of Alaskan tundra fires during the past half-century: implications for surface properties and radiative forcing. *Environ. Res. Lett.* **7**, 044039 (2012).
77. Mack, M. C. et al. Carbon loss from an unprecedented arctic tundra wildfire. *Nature* **475**, 489–492 (2011).
78. Park, T. et al. Changes in growing season duration and productivity of northern vegetation inferred from long-term remote sensing data. *Environ. Res. Lett.* **11**, 084001 (2016).
- Reports that spectral greening trends are more prevalent than spectral browning trends and that there are strong regional differences in satellite-derived estimates of growing season duration and productivity.**

79. Fraser, R., Olthof, I., Carrière, M., Deschamps, A. & Pouliot, D. A method for trend-based change analysis in Arctic tundra using the 25-year Landsat archive. *Polar Rec.* **48**, 83–93 (2012).
80. Frost, G. V., Epstein, H. E. & Walker, D. A. Regional and landscape-scale variability of Landsat-observed vegetation dynamics in northwest Siberian tundra. *Environ. Res. Lett.* **9**, 025004 (2014).
81. Reynolds, M. K., Walker, D. A., Verbyla, D. & Munger, C. A. Patterns of Change within a Tundra Landscape: 22-year Landsat NDVI Trends in an Area of the Northern Foothills of the Brooks Range, Alaska. *Arctic, Antarctic, and Alpine Research* **45**, 249–260 (2013).
82. Weijers, S., Pape, R., Löffler, J. & Myers-Smith, I. H. Contrasting shrub species respond to early summer temperatures leading to correspondence of shrub growth patterns. *Environ. Res. Lett.* **13**, 034005 (2018).
83. Brehaut, L. & Danby, R. K. Inconsistent relationships between annual tree ring-widths and satellite-measured NDVI in a mountainous subarctic environment. *Ecol. Indic.* **91**, 698–711 (2018).
84. Gamm, C. M. et al. Declining growth of deciduous shrubs in the warming climate of continental western Greenland. *J. Ecol.* **106**, 640–654 (2018).
85. Ropars, P. et al. Different parts, different stories: climate sensitivity of growth is stronger in root collars vs. stems in tundra shrubs. *Glob. Change Biol.* **23**, 3281–3291 (2017).
86. Elmendorf, S. C. et al. Experiment, monitoring, and gradient methods used to infer climate change effects on plant communities yield consistent patterns. *PNAS* **112**, 448–452 (2015).
87. Myers-Smith, I. H. et al. Climate sensitivity of shrub growth across the tundra biome. *Nature Clim. Change* **5**, 887–891 (2015).
88. Zeng, H., Jia, G. & Epstein, H. Recent changes in phenology over the northern high latitudes detected from multi-satellite data. *Environ. Res. Lett.* **6**, 045508 (2011).
89. Zeng, H., Jia, G. & Forbes, B. C. Shifts in Arctic phenology in response to climate and anthropogenic factors as detected from multiple satellite time series. *Environ. Res. Lett.* **8**, 035036 (2013).

- 795 90. Park, T. et al. Changes in timing of seasonal peak photosynthetic activity in northern
796 ecosystems. *Glob. Change Biol.* **25**, 2382–2395 (2019).
- 797 91. Helman, D. Land surface phenology: What do we really ‘see’ from space? *Sci. Total Environ.*
798 **618**, 665–673 (2018).
- 799 92. Callaghan, T. V. et al. Multiple effects of changes in Arctic snow cover. *Ambio* **40**, 32–45
800 (2011).
- 801 93. Liu, Q. et al. Extension of the growing season increases vegetation exposure to frost. *Nat.*
802 *Commun.* **9**, 426 (2018).
- 803 94. Myers-Smith, I. H. et al. Eighteen years of ecological monitoring reveals multiple lines of
804 evidence for tundra vegetation change. *Ecol. Monogr.* **89**, e01351 (2019).
- 805 95. Westergaard-Nielsen, A. et al. Transitions in high-Arctic vegetation growth patterns and
806 ecosystem productivity tracked with automated cameras from 2000 to 2013. *Ambio* **46**, 39–52
807 (2017).
- 808 96. Oehri, J., Schmid, B., Schaepman-Strub, G. & Niklaus, P. A. Biodiversity promotes primary
809 productivity and growing season lengthening at the landscape scale. *PNAS* **114**, 10160–10165
810 (2017).
- 811 97. Armstrong, J. B., Takimoto, G., Schindler, D. E., Hayes, M. M. & Kauffman, M. J. Resource
812 waves: phenological diversity enhances foraging opportunities for mobile consumers. *Ecology*
813 **97**, 1099–1112 (2016).
- 814 98. Khorsand Rosa, R. et al. Plant phenological responses to a long-term experimental extension
815 of growing season and soil warming in the tussock tundra of Alaska. *Glob. Change Biol.* **21**,
816 4520–4532 (2015).
- 817 99. Sweet, S. K., Gough, L., Griffin, K. L. & Boelman, N. T. Tall deciduous shrubs offset delayed
818 start of growing season through rapid leaf development in the Alaskan Arctic tundra. *Arct.*
819 *Antarct. Alp. Res.* **46**, 682–697 (2014).
- 820 100. Sweet, S. K., Griffin, K. L., Steltzer, H., Gough, L. & Boelman, N. T. Greater deciduous shrub
821 abundance extends tundra peak season and increases modeled net CO₂ uptake. *Glob.*
822 *Change Biol.* **21**, 2394–2409 (2015).

- 823 101. Verbesselt, J., Hyndman, R., Newnham, G. & Culvenor, D. Detecting trend and seasonal
824 changes in satellite image time series. *Remote Sens. Environ.* **114**, 106–115 (2010).
- 825 102. Tucker, C. J. et al. Higher northern latitude normalized difference vegetation index and growing
826 season trends from 1982 to 1999. *Int. J. Biometeorol.* **45**, 184–190 (2001).
- 827 103. Høye, T. T., Post, E., Meltofte, H., Schmidt, N. M. & Forchhammer, M. C. Rapid advancement
828 of spring in the High Arctic. *Curr. Biol* **17**, R449–R451 (2007).
- 829 104. Harsch, M. A., Hulme, P. E., McGlone, M. S. & Duncan, R. P. Are treelines advancing? A
830 global meta-analysis of treeline response to climate warming. *Ecol. Lett.* **12**, 1040–1049
831 (2009).
- 832 105. Myers-Smith, I. H. & Hik, D. S. Climate warming as a driver of tundra shrubline advance. *J.*
833 *Ecol.* **106**, 547–560 (2017).
- 834 106. Wang, D. et al. Impact of sensor degradation on the MODIS NDVI time series. *Remote Sens.*
835 *Environ.* **119**, 55–61 (2012).
- 836 107. Williams, L. et al. Biological soil crusts of Arctic Svalbard and of Livingston Island, Antarctica.
837 *Polar Biol.* **40**, 399–411 (2017).
- 838 108. Gallo, K., Ji, L., Reed, B., Dwyer, J. & Eidenshink, J. Comparison of MODIS and AVHRR 16-
839 day normalized difference vegetation index composite data. *Geophys. Res. Lett.* **31**, (2004).
- 840 109. Shi, C. & Wang, L. Incorporating spatial information in spectral unmixing: A review. *Remote*
841 *Sens. Environ.* **149**, 70–87 (2014).
- 842 110. Reynolds, M. K., Walker, D. A., Epstein, H. E., Pinzon, J. E. & Tucker, C. J. A new estimate of
843 tundra-biome phytomass from trans-Arctic field data and AVHRR NDVI. *Remote Sens. Lett.* **3**,
844 403–411 (2012).
- 845 111. Anderson, C. B. Biodiversity monitoring, earth observations and the ecology of scale. *Ecol.*
846 *Lett.* **21**, 1572–1585 (2018).
- 847 112. Estes, L. et al. The spatial and temporal domains of modern ecology. *Nat. Ecol. Evol.* **2**, 819
848 (2018).
- 849 113. Woodcock, C. E. & Strahler, A. H. The factor of scale in remote sensing. *Remote Sens.*
850 *Environ.* **21**, 311–332 (1987).

- 851 114. Buitenwerf, R., Rose, L. & Higgins, S. I. Three decades of multi-dimensional change in global
852 leaf phenology. *Nature Clim. Change* **5**, 364–368 (2015).
- 853 115. White, M. A. et al. Intercomparison, interpretation, and assessment of spring phenology in
854 North America estimated from remote sensing for 1982–2006. *Glob. Change Biol.* **15**, 2335–
855 2359 (2009).
- 856 116. Tape, K. D., Sturm, M. & Racine, C. H. The evidence for shrub expansion in Northern Alaska
857 and the Pan-Arctic. *Glob. Change Biol.* **12**, 686–702 (2006).
- 858 117. Richardson, A. D. et al. Tracking vegetation phenology across diverse North American biomes
859 using PhenoCam imagery. *Sci. Data* **5**, 180028 (2018).
- 860 118. Euskirchen, E. S., Bret-Harte, M. S., Shaver, G. R., Edgar, C. W. & Romanovsky, V. E. Long-
861 Term Release of Carbon Dioxide from Arctic Tundra Ecosystems in Alaska. *Ecosystems* **20**,
862 960–974 (2017).
- 863 119. Duffy, J. P. et al. Location, location, location: considerations when using lightweight drones in
864 challenging environments. *Remote Sens. Ecol. Cons.* **0**, (2017).
- 865 120. Dash, J. & Ogutu, B. O. Recent advances in space-borne optical remote sensing systems for
866 monitoring global terrestrial ecosystems. *Prog. Phys. Geog.* **40**, 322–351 (2016).
- 867 121. Luus, K. A. et al. Tundra photosynthesis captured by satellite-observed solar-induced
868 chlorophyll fluorescence. *Geophys. Res. Lett.* **44**, 1564–1573 (2017).
- 869 122. Forkel, M. et al. Identifying environmental controls on vegetation greenness phenology through
870 model–data integration. *Biogeosciences* **11**, 7025–7050 (2014).
- 871 123. Tian, F. et al. Evaluating temporal consistency of long-term global NDVI datasets for trend
872 analysis. *Remote Sens. Environ.* **163**, 326–340 (2015).
- 873 124. Wilson, A. M. & Jetz, W. Remotely sensed high-resolution global cloud dynamics for predicting
874 ecosystem and biodiversity distributions. *PLOS Biology* **14**, e1002415 (2016).
- 875 125. Arndt, K. et al. Arctic greening associated with lengthening growing seasons in Northern
876 Alaska. *Environ. Res. Lett.* (2019) doi:10.1088/1748-9326/ab5e26.
- 877 126. May, J. L., Parker, T., Unger, S. & Oberbauer, S. F. Short term changes in moisture content
878 drive strong changes in Normalized Difference Vegetation Index and gross primary productivity
879 in four Arctic moss communities. *Remote Sens. Environ.* **212**, 114–120 (2018).

- 880 127. Vickers, H. et al. Changes in greening in the high Arctic: insights from a 30 year AVHRR max
881 NDVI dataset for Svalbard. *Environ. Res. Lett.* **11**, 105004 (2016).
- 882 128. Loranty, M. M. et al. Vegetation indices do not capture forest cover variation in upland Siberian
883 larch forests. *Remote Sens.* **10**, 1686 (2018).
- 884 129. Gorelick, N. et al. Google Earth Engine: Planetary-scale geospatial analysis for everyone.
885 *Remote Sensing of Environment* **202**, 18–27 (2017).
- 886 130. Post, E. Erosion of community diversity and stability by herbivore removal under warming.
887 *Proc. R. Soc. B* **280**, 20122722 (2013).
- 888