

Changes in outdoor air pollution due to COVID-19 lockdowns differ by pollutant: evidence from Scotland

Dr Ruairaidh Dobson^a, Dr Sean Semple^a

^a Institute for Social Marketing and Health, University of Stirling, Stirling, Scotland

Corresponding author:

Dr Ruairaidh Dobson

Institute for Social Marketing & Health

G10 Pathfoot Building

University of Stirling

Stirling, Scotland

FK9 4LA

r.p.dobson@stir.ac.uk

Key words: Particulates, air pollution, indoor air

Word count: 1,404

This article has been accepted for publication in *Occupational and Environmental Medicine* following peer review. The definitive copyedited, typeset version Dobson R & Semple S (2020) Changes in outdoor air pollution due to COVID-19 lockdowns differ by pollutant: evidence from Scotland.

Occupational and Environmental Medicine, 77 (11), pp. 798-800 is available online at: <https://doi.org/10.1136/oemed-2020-106659>

© Authors 2020. Reuse of this manuscript version (excluding any databases, tables, diagrams, photographs and other images or illustrative material included where a another copyright owner is identified) is permitted strictly pursuant to the terms of the Creative Commons Attribution-

NonCommercial 4.0 International (CC-BY-NC 4.0) <http://creativecommons.org>

Abstract

Objectives: To examine the impact of COVID-19 lockdown restrictions in March/April 2020 on concentrations of nitrogen dioxide (NO₂) and ambient fine particulate (PM_{2.5}) air pollution measured at roadside monitors across Scotland by comparing data with previous years.

Methods: Publicly available data of PM_{2.5} concentrations from reference monitoring systems at sites across Scotland were extracted for the 31 day period immediately following the imposition of lockdown rules on 23rd March 2020. Similar data for 2017, 2018 and 2019 were gathered for comparison. Mean period values were calculated from the hourly data and logged values compared using pairwise t-tests. Weather effects were corrected using meteorological normalisation.

Results: NO₂ concentrations were significantly lower in the 2020 lockdown period than in the previous three years ($p < 0.001$). Mean outdoor PM_{2.5} concentrations in 2020 were much lower than during the same period in 2019 ($p < 0.001$). However, despite UK motor vehicle journeys reducing by 65%, concentrations in 2020 were within 1 µg/m³ of those measured in 2017 ($p = 0.66$) and 2018 ($p < 0.001$), suggesting that traffic-related emissions may not explain variability of PM_{2.5} in outdoor air in Scotland.

Conclusions: The impact of reductions in motor vehicle journeys during COVID-19 lockdown restrictions may not have reduced ambient PM_{2.5} concentrations in some countries. There is also a need for work to better understand how movement restrictions may have impacted personal exposure to air pollutants generated within indoor environments.

Key messages

- What is already known about this subject?
 - Road traffic has been significantly reduced in countries adopting lockdowns due to COVID-19. Research has shown that this has led to reductions in outdoor air pollution in some locations.
- What are the new findings?
 - Nitrogen dioxide concentrations declined in Scotland following the lockdown, but fine particulate matter did not despite the fall in vehicle use.
- How might this impact on policy or clinical practice in the foreseeable future?
 - Policymakers should take care not to over-estimate improvements in outdoor air quality following COVID-19 lockdowns, and should consider the impact of indoor air pollution on personal exposure during these periods.

Introduction

In the wake of the COVID-19 pandemic many countries introduced wide-ranging restrictions on individual movement and gathering, known as “lockdowns” or “stay-at-home orders”. In the UK, a lockdown was introduced at 20.30 on 23 March 2020.

These new regulations led to substantial falls in road traffic with UK data suggesting motor vehicle journeys reduced by around 65% between 16 March and 28 April 2020 [1]. The result of movement restrictions and reduced traffic volumes has been widely reported in the media (and some scientific studies) to have resulted in improved air quality and lower concentrations of common pollutants, such as fine particulate matter (PM_{2.5}) and nitrogen dioxide (NO₂) [2,3]. It has been suggested that this will result in positive health effects, due to lowered exposure to air pollution, and even that the net effect of the pandemic will be to improve health (due to the adverse health effects of exposure to air pollution, particularly PM_{2.5} [4]).

Analyses of this kind assume that road traffic-related PM_{2.5} is a significant source of personal exposure to fine particles. This may not be true in all locations. Scotland's relatively low ambient PM_{2.5} may be related more closely to natural and non-traffic sources, and may not therefore have fallen following the introductions of the lockdown measures. If PM_{2.5} in outdoor air has not declined, it is possible that net exposure to PM_{2.5} will increase, as people spend more time in their homes where generation of fine particles from activities such as cooking and smoking may produce high concentrations within enclosed and poorly ventilated spaces [5]. NO₂ is specifically associated with vehicle exhaust emissions [6] and so provides a measure of relative traffic for use in this analysis.

Methods

Scottish local authorities maintain a network of automatic monitoring stations for PM_{2.5} and other pollutants. The PM_{2.5} monitors in use comprise gravimetric monitors (TEOMs) and high-precision optical monitors (OAS). These monitors report PM_{2.5} measurements hourly and data are made publicly available on the internet.

To examine the effect of the lockdown on Scotland's air, PM_{2.5} and NO₂ data were extracted from the monitor network for the period from 24 March to 23 April in 2017, 2018, 2019 and 2020. Data from 2020 have only been provisionally validated by the Scottish Government. Data were downloaded using the openair R package.

To simulate the removal of weather effects on pollutant concentrations, meteorological normalisation using the random forest machine learning algorithm [7] was conducted using the rmweather R package. Individual models were calculated for both PM_{2.5} and NO₂ at monitoring sites around Scotland. Models were based on daily mean pollutant concentrations and incorporated wind speed, wind direction, atmospheric pressure, air temperature and relative humidity at the nearest available weather station (downloaded using the worldmet R package). Models used 64 trees and 100 samples.

Arithmetic mean concentrations were calculated for each of 70 PM_{2.5} monitoring stations and 89 NO₂ monitoring stations over this period in each year. Geometric means of these values were calculated for each local authority area where monitoring took place and for Scotland overall in each year.

To determine statistical significance in differences in 2020 PM_{2.5} and NO₂ values for this month vs each other year, both observed and normalised data were log-transformed and compared using a pairwise t-test. Statistical analysis was performed in R v4.0.2 [8].

Results

Across Scotland's air pollution monitoring network, observed and normalised NO₂ concentrations remained close to constant in 2017, 2018 & 2019 but fell substantially in 2020 (pairwise t-test $p < 0.001$ for all years) (Table 1).

By contrast, the observed geometric mean PM_{2.5} concentration over the lockdown period in 2020 was 6.6 µg/m³, very similar to the mean concentration over the same period in 2017 (6.7 µg/m³, pairwise t-test $p = 0.66$). The 2020 value showed a modest decrease (-0.8 µg/m³) in comparison with 2018 (7.4 µg/m³, $p < 0.001$) but was substantially lower than the markedly high concentrations measured in 2019 (12.8 µg/m³, $p < 0.001$). Geometric means of normalised data showed the same pattern, with the 2019 mean higher than the other three years (pairwise t-test $p < 0.001$ for all comparisons) (Table 1).

2019 was a visible outlier in observed data across all local authority areas where PM_{2.5} monitoring was conducted (Figure 1). This is likely due to a sustained meteorological event that brought fine particulate dust from the Saharan desert to the UK atmosphere beginning on 15 April 2019 and persisting through the end of the analysis period on 23 April [9]. Removing that period from the 2019 analysis reduces the mean observed value to 7.8µg/m³, similar to overall values from the three other years in this analysis.

Pollutant	2017 period geometric mean concentration (µg/m ³)	2018 period geometric mean concentration (µg/m ³)	2019 period geometric mean concentration (µg/m ³)	2020 period geometric mean concentration (µg/m ³)
PM _{2.5} (observed)	6.7	7.4	12.8	6.6
PM _{2.5} (normalised)	6.9	6.4	7.6	6.1
NO ₂ (observed)	21.9	23.7	22.4	9.9
NO ₂ (normalised)	25.8	25.4	24.4	15.1

Table 1 – Geometric mean PM_{2.5} and NO₂ in Scotland 24 March – 23 April in years 2017 – 2020, including both observed and normalised data.

Discussion

The lockdown period has provided a natural experiment to examine the potential impact of reducing car journeys on air quality in Scotland. The NO₂ data suggests that car journeys have declined substantially during the lockdown compared to the same period in the previous three years. This may lead to significant health benefits, both from reduced exposure to harmful NO₂ and in reduced rates of traffic accidents and pedestrian collisions.

However, our results suggest that the decline in vehicle-related NO₂ has not coincided with significantly reduced PM_{2.5} concentrations. The health risks of exposure to PM_{2.5} are extremely well-established, including cardiovascular disease, pulmonary illness and stroke. This research has established that reducing the number of vehicles on the road would not be an effective measure to reduce exposure to this pollutant in Scotland, and consequently would not affect incidence of these illnesses.

Our analysis is limited by the data available from the monitoring network. Seven Scottish local authority areas have no NO₂ monitors while nine have no PM_{2.5} monitors, so these data do not cover the entirety of Scotland. Data from 2020 have been provisionally validated by the Scottish Government – while they have undergone screening to identify faulty or suspect data, they have not been ratified following detailed manual review. The later discovery of a fault or error associated with a monitor could change these results retroactively (if, for instance, a new calibration factor were applied). This is unlikely – in summer 2018, three faults were identified in particle monitors across the Scotland-wide network [10]. The use of data from a wide range of sources (70 PM_{2.5} monitors and 90 NO₂ monitors) would limit the impact of a change to an individual monitor.

We have attributed the fall in normalised NO₂ concentrations in 2020 to the lockdown, but underlying effects, including a move towards less-polluting fuels and vehicles, could have contributed to this decline (though likely gradually over a period of years).


We believe these results have important policy and health implications in terms of the use of lockdowns to control future epidemics of infectious disease, and in considering how best to tackle

outdoor air pollution in different countries in the future. Lockdowns are intended to result in people spending more time in their homes. This could increase population exposure to indoor air pollution such as cooking fumes and second-hand tobacco smoke (a particular concern given the high concentrations of PM_{2.5} that can be generated by smoking indoors). Previous work suggests that living with a smoker can increase a person's daily dose of PM_{2.5} by over 80% [11].

In countries like Scotland where it appears that the lockdown has not led to reductions in outdoor fine particulate matter pollution, it is possible that personal exposure to PM_{2.5} may actually have increased rather than declined due to higher concentrations from indoor sources of particulate within the home setting. This could increase adverse health effects overall and also health inequalities - lower income people are more likely to smoke and to smoke indoors [12], and are likely to have smaller homes leading to higher PM_{2.5} concentrations from individual sources, due to smaller room volumes. If the severity of COVID-19 is related to air pollution exposure (as has been suggested [13]), increased exposure to PM_{2.5} could potentially increase the death toll of that disease. Careful and balanced consideration of both outdoor and indoor sources of PM_{2.5} is essential to tackling the health harm of air pollution effectively and equitably.

Figures

Figure 1 - Observed geometric mean PM_{2.5} and NO₂ by local government divisions (council areas) in Scotland 24 March – 23 April in years 2017 – 2020. Note that some local authorities have NO₂ monitors but not PM_{2.5} monitors.



Contributorship statement

RD and SS conceived of the idea for the study. Both authors designed the study. RD conducted data analysis and drafted the manuscript, which SS critically reviewed. SS supervised the project.

Funding

There is no funding to report for this study.

Competing interests

The authors report no competing interests.

Data Sharing/Data availability

Data used in this study are available from the Scottish Government's air quality repository (www.scottishairquality.scot).

References

- 1 Prime Minister's Office. Slides and datasets to accompany coronavirus press conference. UK Gov. 2020.<https://www.gov.uk/government/publications/slides-and-datasets-to-accompany-coronavirus-press-conference-28-april-2020> (accessed 29 Apr 2020).
- 2 Dutheil F, Baker JS, Navel V. COVID-19 as a factor influencing air pollution? *Environ. Pollut.* 2020;**263**:114466. doi:10.1016/j.envpol.2020.114466
- 3 Cadotte M. Early evidence that COVID-19 government policies reduce urban air pollution. doi:10.31223/OSF.IO/NHGX3
- 4 Polichetti G, Cocco S, Spinali A, *et al.* Effects of particulate matter (PM₁₀, PM_{2.5} and PM₁) on the cardiovascular system. *Toxicology* 2009;**261**:1–8. doi:10.1016/j.tox.2009.04.035
- 5 Klepeis NE, Bellettiere J, Hughes SC, *et al.* Fine particles in homes of predominantly low-income families with children and smokers: Key physical and behavioral determinants to inform indoor-air-quality interventions. *PLoS One* 2017;**12**:e0177718. doi:10.1371/journal.pone.0177718
- 6 Costa S, Ferreira J, Silveira C, *et al.* Integrating Health on Air Quality Assessment—Review Report on Health Risks of Two Major European Outdoor Air Pollutants: PM and NO₂. *J Toxicol Environ Heal Part B* 2014;**17**:307–40. doi:10.1080/10937404.2014.946164
- 7 Grange SK, Carslaw DC. Using meteorological normalisation to detect interventions in air quality time series. *Sci Total Environ* 2019;**653**:578–88. doi:10.1016/j.scitotenv.2018.10.344
- 8 R Core Team. R: A language and environment for statistical computing. 2019.<https://www.r-project.org/>
- 9 BBC News. Saharan dust: Why your car is covered in red dirt. 2019.<https://www.bbc.co.uk/news/uk-scotland-48037362> (accessed 29 Apr 2020).
- 10 Ricardo Energy and Environment. Scottish Air Quality Database Annual Report 2018. Didcot: 2019.
- 11 Semple S, Apsley A, Azmina Ibrahim T, *et al.* Fine particulate matter concentrations in smoking households: just how much secondhand smoke do you breathe in if you live with a smoker who smokes indoors? *Tob Control* 2015;**24**:e205-11. doi:10.1136/tobaccocontrol-2014-051635
- 12 Scottish Government Community Health & Social Care Directorate. Scottish health survey 2018: key findings. 2019.<https://www.gov.scot/publications/scottish-health-survey-2018-summary-key-findings/> (accessed 29 Apr 2020).
- 13 Conticini E, Frediani B, Caro D. Can atmospheric pollution be considered a co-factor in extremely high level of SARS-CoV-2 lethality in Northern Italy? *Environ. Pollut.* 2020;**114465**. doi:10.1016/j.envpol.2020.114465