


Journal Pre-proof

Physiological and perceptual responses to sprint interval exercise using arm versus leg cycling ergometry

Todd A. Astorino, Shealin Pierce, Madisen B. Piva, Richard S. Metcalfe, Niels B.J. Vollaard



PII: S2666-3376(24)00007-6

DOI: <https://doi.org/10.1016/j.smhs.2024.01.007>

Reference: SMHS 197

To appear in: *Sports Medicine and Health Science*

Received Date: 31 October 2023

Revised Date: 17 December 2023

Accepted Date: 26 January 2024

Please cite this article as: Astorino TA, Pierce S, Piva MB, Metcalfe RS, Vollaard NBJ, Physiological and perceptual responses to sprint interval exercise using arm versus leg cycling ergometry, *Sports Medicine and Health Science* (2024), doi: <https://doi.org/10.1016/j.smhs.2024.01.007>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Chengdu Sport University. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd.

Physiological and Perceptual Responses to Sprint Interval Exercise Using Arm versus Leg
Cycling Ergometry

Todd A. Astorino,¹ Shealin Pierce,¹ Madisen B. Piva,¹ Richard S. Metcalfe,² Niels B. J.
Vollaard³

¹Department of Kinesiology; CSU—San Marcos, San Marcos, CA;²Applied Sports, Technology,
Exercise and Medicine (A-STEM) Research Center. Swansea University, Swansea UK;³Faculty
of Health Sciences and Sport, University of Stirling, Stirling UK

Todd A. Astorino: astorino@csusm.edu

Shealin Pierce pierc056@cougars.edu

Madisen Piva piva001@cougars.csusm.edu

Richard Metcalfe r.s.metcalfe@swansea.ac.uk

Niels Vollaard n.vollaard@stir.ac.uk

Corresponding author: Todd A. Astorino Ph.D

Professor, Department of Kinesiology
California State University, San Marcos
333. S. Twin Oaks Valley Road, UNIV 320
San Marcos, CA 92096-0001
Phone: (760) 750-7351
Fax: (760) 750-3237
Email: astorino@csusm.edu

Physiological and perceptual responses to sprint interval exercise using arm versus leg cycling
ergometry

19 List of Abbreviations

20	ACE	arm cycling ergometry
21	ATP	adenosine triphosphate
22	b/min	beats per minute
23	B-HAD	3-hydroxyacyl-CoA dehydrogenase
24	BLa	blood lactate concentration
25	BMI	body mass index
26	CO _{max}	maximal cardiac output
27	FT	fast twitch
28	h	hours
29	HIIT	high intensity interval training
30	HR	heart rate
31	HR _{max}	maximal heart rate
32	h/wk	hours per week
33	kg·m ⁻²	kilograms per meter squared
34	LCE	leg cycling ergometry
35	min	minutes

36	PACES	physical activity enjoyment scale
37	rev/min	revolutions per minute
38	RPE	rating of perceived exertion
39	REHIT	reduced exertion high intensity training
40	RER	respiratory exchange ratio
41	s	seconds
42	SIE	sprint interval exercise
43	ST	slow twitch
44	$\dot{V}CO_2$	carbon dioxide production
45	\dot{V}_E	ventilation
46	$\dot{V}O_{2max}$	maximal oxygen consumption
47	$\dot{V}O_2$	oxygen uptake
48	W	watts
49	W _{max}	maximal workload
50	W/min	watts per minute
51	y	years

52

Abstract

Increases in power output and maximal oxygen consumption ($\dot{V}O_{2\max}$) occur in response to sprint interval exercise (SIE), but common use of “all-out” intensities presents a barrier for many adults. Furthermore, lower-body SIE is not feasible for all adults. We compared physiological and perceptual responses to supramaximal, but “non-all-out” SIE between leg and arm cycling exercise. Twenty-four active adults (mean \pm SD age: $[25 \pm 7]$ y; cycling $\dot{V}O_{2\max}$: $[39 \pm 7]$ mL \cdot kg $^{-1}\cdot$ min $^{-1}$) performed incremental exercise using leg (LCE) and arm cycle ergometry (ACE) to determine $\dot{V}O_{2\max}$ and maximal work capacity (W_{\max}). Subsequently, they performed four 20 second (s) bouts of SIE at 130% W_{\max} on the LCE or ACE at cadence = 120-130 rev/min, with 2 minutes (min) recovery between intervals. Gas exchange data, heart rate (HR), blood lactate concentration (BLa), rating of perceived exertion (RPE), and affective valence were acquired. Data showed significantly lower ($p < 0.001$) absolute mean ($[1.24 \pm 0.31]$ L \cdot min $^{-1}$ vs. $[1.59 \pm 0.34]$ L \cdot min $^{-1}$; $d = 1.08$) and peak $\dot{V}O_2$ ($[1.79 \pm 0.48]$ L \cdot min $^{-1}$ vs. $[2.10 \pm 0.44]$ L \cdot min $^{-1}$; $d = 0.70$) with ACE versus LCE. However, ACE elicited significantly higher ($p < 0.001$) relative mean ($[62\% \pm 9\%]$ $\dot{V}O_{2\max}$ vs. $[57\% \pm 7\%]$ $\dot{V}O_{2\max}$, $d = 0.63$) and peak $\dot{V}O_2$ ($[88\% \pm 10\%]$ $\dot{V}O_{2\max}$ vs. $[75\% \pm 10\%]$ $\dot{V}O_{2\max}$, $d = 1.33$). Post-exercise BLa was significantly higher ($[7.0 \pm 1.7]$ mM vs. $[5.7 \pm 1.5]$ mM, $p = 0.024$, $d = 0.83$) for LCE versus ACE. There was no significant effect of modality on RPE or affective valence ($p > 0.42$), and lowest affective valence recorded (2.0 ± 1.8) was considered “good to fairly good”. Data show that non “all-out” ACE elicits lower absolute but higher relative HR and $\dot{V}O_2$ compared to LCE. Less aversive perceptual responses could make this non-all-out modality feasible for inactive adults.

Key words: high intensity interval training; upper body exercise; peak power output; oxygen uptake; blood lactate concentration

1. Introduction

Sprint interval exercise (SIE) consists of brief (5–30 s), repeated, and exhaustive efforts at intensities greater than that associated with maximal oxygen consumption ($\dot{V}O_{2\max}$) or maximal work capacity (W_{\max}) separated by low intensity or resting recovery.¹ Although there are various iterations of SIE, the most widely used protocols require completion of 4–6 Wingate tests,²⁻³ 10 s cycling sprints at 170% W_{\max} ,⁴ or two or three 20 s sprints within a 10-minute (min) session (reduced-exertion high-intensity training; REHIT).⁵⁻⁷ One unique attribute of SIE is the low training volume (1–3 min) compared to high intensity interval exercise (10–16 min) or moderate intensity continuous exercise (20–60 min). It is likely that generation of extremely high work rates characteristic of SIE is critical to resultant training-induced increases in $\dot{V}O_{2\max}$,^{2-3,8} insulin sensitivity,⁹ fat oxidation,²⁻³ and oxidative capacity^{3,10} despite the extremely low training volume.

Nevertheless, SIE requires “all-out” efforts characterized by attainment of maximal cadence and in turn, power outputs higher than that associated with $\dot{V}O_{2\max}$, which may be undesirable in inactive adults. In some cases, SIE can elicit extreme fatigue, hyperventilation, nausea, and dizziness² which may reduce its widespread application in this population. In fact, Hardcastle et al.¹¹ stated that SIE is inappropriate for the typical inactive adult as it may be perceived as too arduous which would lead to feelings of displeasure and in turn, low adherence.¹² Nevertheless, data show that pleasure: displeasure remains positive (average affective valence ~ 1.0 – 1.5) in less fit adults who engage in relatively low-volume SIE.^{5,13} In a recent systematic review and meta-analysis, Hu et al.¹⁴ revealed that low-volume SIE protocols using shorter sprints and lower

number of efforts induced more positive affective responses, suggesting the feasibility of SIE in adults.

Several approaches exist to reduce the metabolic perturbation of vigorous exercise including SIE. One option is to reduce sprint duration. In young adults, Islam et al.¹⁵ compared physiological responses to work-matched bouts of treadmill-based SIE requiring durations of 5 s, 15 s, and 30 s using a 1:8 ratio of work:recovery. Compared to the longer durations, $\dot{V}O_2$ and energy expenditure were significantly higher with the 5 s sprints which was attendant with greater intention to engage in this protocol and more positive affective valence,¹⁶ emphasizing the importance of brief sprint durations to augment the tolerability of SIE. In addition, Vollaard and Metcalfe¹⁷ revealed that fewer number (2–3) and shorter intervals (10 s or 20 s) provide similar health benefits as the traditional 30 s Wingate-based SIE regimen.

An additional element that can be modified to reduce the physiological response to SIE is to not require “all-out” efforts which should attenuate the level of fatigue experienced by participants. Although all SIE is characterized by supramaximal sprints, this includes exercise intensities ranging from just above W_{max} to several-fold higher intensities achieved in all-out sprints (e.g., ~350% of $\dot{V}O_{2max}$).¹⁸ Bayati et al.¹⁹ revealed similar increases in $\dot{V}O_{2max}$ and W_{max} in response to 12 sessions of “all-out” SIT (30 s Wingate tests) compared to a higher volume of 30 s efforts at 125% W_{max} , which would suggest that the level of effort maintained during supramaximal sprints does not affect the chronic response. To our knowledge, no study has examined acute physiological and perceptual responses to SIE characterized by intervals which are supramaximal, but not all-out.

The majority of studies employing SIE used leg cycle ergometry (LCE),^{2-3,10,19} although some have employed treadmill sprinting.^{16,20-21} One disadvantage of cycling-based SIE is that it leads to lightheadedness, leg pain, and nausea and in turn, displeasure.¹¹ Furthermore, LCE is not feasible for all individuals e.g. most people with spinal cord injury.²² An alternative modality to LCE is arm cycle ergometry (ACE) which has been widely implemented in persons with heart disease²³ and spinal cord injury²⁴ to improve physical fitness and function. Price et al.²⁵ reported higher peak and mean power output, yet no difference in heart rate or respiratory exchange ratio, between the Wingate test performed using LCE versus ACE. In adults, Zinner et al.²⁶ reported that six sessions of SIE using ACE and LCE increased upper-body $\dot{V}O_{2\max}$ slightly more than that of the legs despite less work being performed during ACE. However, little is known about the acute physiological response to non “all-out” SIE performed using ACE and how this may compare to LCE. At a given submaximal or maximal absolute work rate, ACE elicits higher HR and $\dot{V}O_2$ versus LCE due to use of a smaller exercising muscle mass and the lower efficiency of arm cycling.²⁷

The aim of the present study was to compare physiological and perceptual responses to SIE between LCE and ACE characterized by supramaximal, but non-all-out efforts. Reducing the effort attendant with SIE may attenuate blood lactate accumulation, enhance perceptual responses, and in turn, make it more feasible for the majority of adults who are insufficiently active and likely intolerant of “all-out” efforts. We hypothesize that supramaximal, but non-all-out ACE will be associated with less aversive perceptual responses, but considering the lower active muscle mass and higher contribution of type 2 muscle fibers, will present a lower absolute and relative cardiopulmonary response compared to LCE.

2. Material and methods

2.1. Experimental design and subjects

This repeated measures, crossover study examined differences in various outcomes between brief bouts of SIE characterized by different active muscle mass. Participants initially underwent incremental exercise to exhaustion to determine W_{\max} and $\dot{V}O_{2\max}$ on both the leg and arm cycle ergometer. On the second visit, they completed a familiarization trial comprising two bouts of SIE on both exercise modes. For the final two sessions, order of assignment to ACE or LCE was randomized, and a minimum of 48 h separated each visit, which were held at the same time of day (08:00 to 13:00) within participants. Physiological and perceptual responses were acquired during the sessions. All participants were asked to be well-rested, hydrated, and refrain from intense exercise for 36 h prior to all sessions. A study flow diagram is shown in Figure 1.

Recreationally-active men ($n = 15$) and women ($n = 9$) were recruited by word-of-mouth. Inclusion criteria included age 18–50 y, healthy, non-obese, non-smoker, participation in 150 min/wk of moderate or 75 min/wk of vigorous exercise, and no joint issues which would be worsened by upper- or lower-body sprint cycling.

2.2. Ethical approval

Participants provided written informed consent, and study experimental procedures were reviewed and approved by the Institutional Review Board at CSU—San Marcos (Protocol 1876593-1). The study was conducted in accordance with the Declaration of Helsinki.

2.3. Testing of maximal oxygen uptake

Initially, height and body mass were determined and used to calculate body mass index (BMI). Subsequently, skinfold measurements were performed at chest, abdomen, and thigh for men and triceps, suprailiac, and thigh for women.²⁸⁻²⁹ to determine percent body fat from body density.³⁰ Then, participants completed incremental exercise to volitional exhaustion on both an electrically-braked cycle ergometer (Velotron RacerMate, Quark, SD) and arm cycle ergometer (Lode Angio, Groningen, Netherlands). Order of assignment to LCE versus ACE incremental test was randomized and separated by a 30 min recovery period.³¹ Our preliminary data in four active men and women show similar values of W_{max} (difference ≤ 4 W) and $\dot{V}O_{2max}$ (difference $\leq 2.5\%$) when these tests are performed on separate days or separated by 30 min as performed in the present study. Graded exercise on the ACE began with a 2 min warm up at 7 W after which power output was increased in a ramp-like manner by 8, 15, or 20 Watt/min (W/min) until volitional exhaustion which occurred when pedal cadence was below 50 rev/min.³² The pedal crank was aligned to the height of the shoulder joint and there was a small degree of elbow flexion. Participants were seated, required to keep their feet shoulder width apart, and encouraged to use their lower body, since lower body restriction reduces $\dot{V}O_{2max}$ and power output during ACE.³³

Incremental exercise using LCE began with a 2 min warm up at 40-60 W. Power output was subsequently increased in a ramp-like manner by 20-35 W/min until volitional exhaustion which was determined by pedal cadence below 50 rev/min. Different work rate increments were used across participants to account for differences in sex, body size, and fitness level and to ensure duration of incremental exercise between 8–12 min. Throughout exercise, heart rate (HR) was assessed continuously via telemetry (Polar, Woodbury, NY), and gas exchange data ($\dot{V}O_2$, $\dot{V}CO_2$, $\dot{V}E$, and respiratory exchange ratio [RER]) were acquired at 10 s increments using a

metabolic cart (ParvoMedics True One, Sandy, UT), which was calibrated prior to testing according to manufacturer guidelines.

$\dot{V}O_{2\max}$ was identified as the mean of the two highest 10 s values at exercise termination. Workload (in Watts) at volitional fatigue was identified as W_{\max} and used to determine the exercise intensities of subsequent SIE bouts. To verify attainment of $\dot{V}O_{2\max}$, the following criteria were used: change in $\dot{V}O_2 < 0.15 \text{ L} \cdot \text{min}^{-1}$ at $\dot{V}O_{2\max}$ and $\text{RER} > 1.10$.³⁴⁻³⁵

2.4. Familiarization session

Most participants had no experience with SIE, so a familiarization session on both ergometers was performed. After a 3 min warm-up at 20% W_{\max} , participants completed two 20 s bouts of SIE at the required cadence separated by 2 min of active recovery at 20% W_{\max} . They completed a 5 min passive recovery, then performed two SIE bouts on the other modality, whose order was randomized across participants. Perceptual responses and HR were acquired pre-exercise, immediately after each sprint, and halfway into recovery between sprints.

2.5. Completion of sprint interval exercise

SIE sessions began with a 3 min warm up at 20% W_{\max} succeeded by four 20 s sprints at 130% W_{\max} at a cadence between 120-130 rev/min, which was closely monitored during each interval. This cadence was selected for two reasons. First, pilot testing revealed that young adults can attain peak cadences during ACE exceeding 150 rev/min. In addition, prior work from our lab employing SIE on the cycle ergometer shows that men and women can achieve peak cadences $\geq 180 \text{ rev/min}$.^{2,5} Approximately 5 s before each sprint, participants were required to increase pedal cadence so by the start of the interval, they were pedaling at the desired cadence which is when resistance was applied to the ergometer. Intervals were interspersed by 2 min

recovery at 20% Wmax. This protocol was chosen as “all-out” SIE protocols comprising fewer sprint repetitions and shorter durations^{5,9} generate significant improvements in cardiorespiratory fitness⁶ yet elicit more positive affective responses.¹³⁻¹⁴ This power output is appropriate for nonathletic adults, elicits significant BLa (~12 mM),³⁶ and the 1:6 work:rest ratio is adequate to promote recovery. Gas exchange data and HR were acquired every 10 s throughout exercise. Values from each interval were determined as the two 10 s values during exercise and first value in recovery, due to the lag in HR and $\dot{V}O_2$ during SIE.³⁷ Recovery values were calculated from the last 60 s of recovery (6 values). Mean $\dot{V}O_2$, \dot{V}_E , RER, and HR were identified as the average value from the session (9 min and 20 s), not including the warm-up. Peak values were determined as the average of any three consecutive 10 s values recorded during the session.

2.6. Assessment of perceptual responses and blood lactate concentration

Prior to exercise, participants were seated and read specific instructions pertaining to what each scale represented. The Borg 6-20 RPE scale was used to measure perceived exertion in response to exercise.³⁸ To communicate the meaning of the RPE scale, participants were instructed to report their exertion according to their level of fatigue, breathing, and HR.³⁸ Affective valence (assessed using the 11-point Feeling Scale, rating from +5 very good to -5 very bad including 0)³⁹ was described by reciting the following text: *While participating in exercise, it is common to experience changes in mood. Some individuals find exercise pleasurable; whereas, others find it to be unpleasant. Additionally, feeling may fluctuate across time. That is, one might feel good and bad a number of times during exercise.* Participants were instructed to respond to each scale according to their perception at that moment, and their score was repeated back to them before being recorded. These measures were acquired pre-exercise, at the end of the warm-up, immediately on completion of each interval and 1 min into each

recovery period. Five min post-exercise, participants were administered the 18-item Physical Activity Enjoyment Scale⁴⁰ (PACES) to assess their enjoyment of each session. This scale is widely employed in similar studies analyzing how acute exercise mediates enjoyment measured post-exercise.^{32,36, 41-42} Blood samples were acquired pre-, midway (after interval 2), and 3 min post-exercise to assess changes in blood lactate concentration (BLa). Participants remained seated and after the fingertip was cleaned with a damp towel, dried, and then the first drop of blood wiped away, a 0.7 µl blood sample was taken using a lancet (Owen Mumford Inc., Marietta, GA) and portable monitor (Lactate Plus, Sports Research Group, New Rochelle, NY).

2.7. Consideration of dietary intake

To reduce the potential effects of dietary changes on study outcomes, participants were asked to complete a 36 h food diary before their first SIE session. This was submitted to the investigators who advised participants to replicate this pattern before the final SIE session, which was done in all participants.

2.8. Data analysis

Data are reported as means \pm standard deviation (*SD*) and were analyzed using SPSS Version 27 (IBM, NY). We determined the normality of data distributions using the Shapiro-Wilks test. To identify differences in our outcome measures between modalities, two-way repeated measures ANOVA was used, with two levels for modality, and three (BLa) or eight levels (gas exchange data, HR, RPE, and affective valence) for time. If a significant *F* ratio was obtained, Tukey's post hoc test was used to identify differences between means. The Greenhouse-Geisser correction was used if the sphericity assumption was violated. Paired *t*-test was used to assess differences in enjoyment and mean/peak and maximal variables between arm

and leg cycling. Cohen's d was used as a measure of effect size, with a small, medium, and large effect equal to 0.2, 0.5, and 0.8, respectively.⁴³ G Power⁴⁴ was used to confirm that a sample size of nine per condition is adequate to detect a change in $\dot{V}O_2$ equal to $0.20 \text{ L}\cdot\text{min}^{-1}$ across modalities, a difference shown in a prior study comparing these modalities.³¹ Although our study was not adequately powered to detect differences between men and women, sex was used as a between-subjects variable in these analyses. Independent t -test was used to identify significant differences in peak and mean outcomes between men and women. Statistical significance was set at $p < 0.05$.

3. Results

3.1. Comparison of maximal data between LCE and ACE

Our participants' demographic data (mean \pm SD) were as follows: age (25 ± 7) y; body fat, ($16\% \pm 6\%$); body mass index: (25 ± 4) $\text{kg}\cdot\text{m}^{-2}$; physical activity: (6 ± 3) h/wk; LCE $\dot{V}O_{2\text{max}}$: (39 ± 7) $\text{mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$. As expected, $\dot{V}O_{2\text{max}}$, W_{max} , and V_E were significantly higher in response to LCE, as was maximal RER, BLa, and HR (Table 1). The relative $\dot{V}O_{2\text{max}}$ values obtained from LCE classify our participants as having average cardiorespiratory fitness ($\dot{V}O_{2\text{max}} = [31\text{--}42] \text{ mL}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$) according to Kaminsky et al.⁴⁵ ACE-derived $\dot{V}O_{2\text{max}}$ was 69% of the mean value from LCE, which supports prior data,^{32,46} although this varied from 52%-91% across participants.

3.2. Familiarization session

This session elicited peak HR equal to ($85.7\% \pm 5.6\%$) HR_{max} and ($85.8\% \pm 6.0\%$) HR_{max} for LCE and ACE, and peak RPE and affective valence equal to (10.4 ± 2.3) vs. (10.3 ± 2.7) and (2.9 ± 1.3) vs. (3.2 ± 1.3), respectively.

3.3. Comparison of gas exchange data and heart rate between sprint interval exercise using LCE and ACE

Results showed no significant mode \times time interaction ($p = 0.51$) for $\dot{V}O_2$ although there was a main effect of time and mode ($p < 0.001$). Compared to rest, $\dot{V}O_2$ increased six-fold during bout 4 in LCE ($[0.33 \pm 0.10] \text{ L} \cdot \text{min}^{-1}$ vs. $[1.87 \pm 0.34] \text{ L} \cdot \text{min}^{-1}$, $d = 6.1$) and five-fold in response to ACE ($[0.30 \pm 0.05] \text{ L} \cdot \text{min}^{-1}$ vs. $[1.58 \pm 0.45] \text{ L} \cdot \text{min}^{-1}$, $d = 4.6$) (Figure 2). At all timepoints, LCE exhibited higher $\dot{V}O_2$ than ACE ($d = 0.74\text{-}1.30$). Ventilation showed a main effect of time and mode ($p < 0.001$) but no mode \times time interaction ($p = 0.83$). With exception of \dot{V}_E obtained in recovery after bouts 3 and 4, all exercise values were different from each other ($p < 0.05$, $d = 0.34\text{-}1.36$). Post hoc analyses showed that \dot{V}_E was higher in response to LCE versus ACE at all time points ($d = 0.44\text{-}1.02$). Resting RER was equal to (0.88 ± 0.08) and (0.89 ± 0.08) prior to LCE and ACE and significantly increased during the session ($p < 0.001$), yet there was no mode \times time interaction ($p = 0.11$) or effect of mode ($p = 0.24$). Recovery RER values were significantly higher ($p < 0.001$, $d = 0.44\text{-}3.74$) than those recorded in response to exercise and peaked after bout 2 ($[1.33 \pm 0.15]$ and $[1.40 \pm 0.14]$ for LCE and ACE). Results showed that HR increased during SIE ($p < 0.001$) and there was a significant effect of mode ($p < 0.001$) and mode \times time interaction ($p = 0.007$). All values recorded during exercise were different ($p < 0.05$) from each other with exception of the value from bout 1 and HR recorded during recovery from bouts 3 and 4.

3.4. Comparison of mean and peak responses between LCE and ACE

Mean $\dot{V}O_2$ ($\text{L} \cdot \text{min}^{-1}$) and HR (b/min) were significantly higher ($p < 0.001$) in response to LCE versus ACE, as were peak $\dot{V}O_2$ ($p < 0.001$) and HR ($p = 0.01$). Data also revealed higher

mean \dot{V}_E ($p < 0.001$) and peak \dot{V}_E ($L \cdot min^{-1}$) ($p = 0.008$) on the LCE. Results showed that LCE elicited higher relative mean ($p = 0.04$), but not peak %HRmax ($p = 0.71$) compared to ACE. Other data revealed that ACE elicited higher mean ($p = 0.02$) and peak % $\dot{V}O_{2max}$ ($p < 0.001$) as well as higher relative mean ($p = 0.048$) and peak % \dot{V}_{Emax} ($p = 0.017$). Table 2 reveals differences in these outcomes between modalities.

3.5. Comparison of blood lactate concentration between LCE and ACE

Blood lactate concentration increased during SIE (main effect, $p < 0.001$) and there was a mode \times time interaction ($p = 0.047$) (Figure 3a). Post hoc analyses revealed that BL_a after bout 2 was higher in response to LCE compared to ACE ($d = 0.83$).

3.6. Psychological responses to LCE and ACE

Figure 3b-c documents changes in RPE and affective valence in response to SIE across modalities. There was a main effect of time as RPE increased during SIE ($p < 0.001$) and peaked at values nearing 13 for both modalities, representing a “somewhat hard” level of exertion. RPE increased by approximately one unit with each successive interval and then declined by the same magnitude in recovery. Results showed no effect of mode ($p = 0.64$) or mode \times time interaction ($p = 0.46$). Similar data were shown for affective valence, which significantly declined ($p < 0.001$) in response to SIE yet there was no main effect of mode ($p = 0.84$) or mode \times time interaction ($p = 0.89$). The lowest value of affective valence was equal to (2.0 ± 1.8) and (1.8 ± 1.9) for LCE and ACE, respectively, which lies between “fairly good” and “good.” There was no difference ($p = 0.97$, $d = 0.08$) in enjoyment between modalities ($[102 \pm 15]$ and $[101 \pm 18]$ for LCE and ACE, respectively).

3.7. Exploratory sex-based analyses

Data from baseline $\dot{V}O_{2\max}$ testing showed no difference in relative $\dot{V}O_{2\max}$ ($p = 0.34$ and 0.63 for LCE and ACE), HR_{\max} ($p = 0.30$ and 0.36 for LCE and ACE), RER_{\max} ($p = 0.51$ and 0.11 for LCE and ACE), or maximal BLa ($p = 0.14$ and 0.36 for LCE and ACE) between men and women, yet significant differences occurred in $\dot{V}_{E\max}$ for LCE ($[132 \pm 32] \text{ L} \cdot \text{min}^{-1}$ vs. $[100 \pm 18] \text{ L} \cdot \text{min}^{-1}$, $p = 0.02$, $d = 1.2$) and ACE ($[99 \pm 20] \text{ L} \cdot \text{min}^{-1}$ vs. $[73 \pm 21] \text{ L} \cdot \text{min}^{-1}$, $p = 0.01$, $d = 1.3$) for men versus women.

During LCE, significant differences were shown in mean HR ($[142 \pm 12] \text{ b/min}$ vs. $[159 \pm 11] \text{ b/min}$, $p = 0.003$, $d = 1.5$; $[78\% \pm 5\%]$ vs. $[85\% \pm 6\%]$ HR_{\max} , $p = 0.004$, $d = 1.6$) and peak HR ($[163 \pm 12] \text{ b/min}$ vs. $[174 \pm 12] \text{ b/min}$, $p = 0.04$, $d = 1.0$; $[88\% \pm 5\%]$ vs. $[93\% \pm 4\%]$ HR_{\max} , $p = 0.02$, $d = 1.1$), with significantly higher values recorded in women. There was no sex difference in mean ($[56\% \pm 8\%]$ $\dot{V}O_{2\max}$ vs. $[58\% \pm 6\%]$ $\dot{V}O_{2\max}$, $p = 0.61$, $d = 0.3$) or peak $\dot{V}O_2$ ($[73 \pm 9] \text{ L} \cdot \text{min}^{-1}$ vs. $[80\% \pm 10\%]$ $\dot{V}O_{2\max}$, $p = 0.10$, $d = 0.80$) expressed according to % $\dot{V}O_{2\max}$, although men displayed higher absolute $\dot{V}O_2$ ($p < 0.002$, $d = 1.6\text{--}2.2$) which is attributed to their greater body mass. As far as \dot{V}_E , there was no difference in any outcome between men and women ($p = 0.11\text{--}0.43$) other than mean \dot{V}_E which was significantly higher in men compared to women ($[60 \pm 9] \text{ L} \cdot \text{min}^{-1}$ vs. $[51 \pm 11] \text{ L} \cdot \text{min}^{-1}$, $p = 0.03$, $d = 1.0$). There was no difference in mean ($p = 0.61$) or peak RER ($p = 0.11$) between men and women.

In response to ACE, there was no difference in mean HR when expressed in absolute ($[133 \pm 13] \text{ b/min}$ vs. $[144 \pm 22] \text{ b/min}$, $p = 0.15$, $d = 0.70$) or relative terms ($[76\% \pm 5\%]$ vs. $[79\% \pm 8\%]$ HR_{\max} , $p = 0.35$, $d = 0.5$). Similar lack of differences was shown for peak HR expressed in b/min ($[156 \pm 16] \text{ b/min}$ vs. $[169 \pm 17] \text{ b/min}$, $p = 0.10$, $d = 0.8$) and % HR_{\max} ($[89\% \pm 8\%]$ vs. $[91\% \pm 5\%]$ HR_{\max} , $p = 0.41$, $d = 0.3$). Despite no difference in relative peak $\dot{V}O_2$ between men and women ($[90\% \pm 11\%]$ vs. $[83\% \pm 9\%]$, $p = 0.12$, $d = 0.7$), mean relative $\dot{V}O_2$ was higher in

men compared to women ($[64\% \pm 9\%] \dot{V}O_{2\max}$ vs. $[56\% \pm 4\%] \dot{V}O_{2\max}$, $p = 0.03$, $d = 0.7$) as was absolute $\dot{V}O_2$ ($[1.38 \pm 0.24] \text{ L} \cdot \text{min}^{-1}$ vs. $[0.92 \pm 0.22] \text{ L} \cdot \text{min}^{-1}$, $p < 0.001$, $d = 2.0$ and $[1.98 \pm 0.41] \text{ L} \cdot \text{min}^{-1}$ vs. $[1.38 \pm 0.32] \text{ L} \cdot \text{min}^{-1}$, $p = 0.002$, $d = 1.6$). Relative \dot{V}_E was not different between men and women ($[56\% \pm 12\%] \dot{V}_{E\max}$ vs. $[58\% \pm 18\%] \dot{V}_{E\max}$, $p = 0.80$, $d = 0.2$; $[80\% \pm 20\%] \dot{V}_{E\max}$ vs. $[82\% \pm 22\%] \dot{V}_{E\max}$, $p = 0.80$, $d = 0.20$), although absolute \dot{V}_E was higher in men compared to women ($[52 \pm 11] \text{ L} \cdot \text{min}^{-1}$ vs. $[40 \pm 11] \text{ L} \cdot \text{min}^{-1}$, $p = 0.02$, $d = 1.1$; $[75 \pm 20] \text{ L} \cdot \text{min}^{-1}$ vs. $[57 \pm 17] \text{ L} \cdot \text{min}^{-1}$, $p = 0.04$, $d = 1.0$). There was no difference in mean ($p = 0.44$) or peak RER ($p = 0.44$) between men and women. No interaction ($p = 0.43$) or main effect ($p = 0.07$) was shown for BLA or PACES ($p = -0.30$ and 0.59 for LCE and ACE) between men and women.

4. Discussion

This study compared physiological and perceptual responses to SIE performed using ACE and LCE. The results oppose our hypothesis since ACE elicits a lower absolute, but a higher relative cardiovascular response versus LCE, alongside a lower BLA response. No differences in RPE, affective valence, or post-exercise enjoyment were shown between modalities. In addition, our results support our hypothesis as affective valence remained positive on average and enjoyment was relatively high, suggesting that LCE and ACE involving four 20 s supramaximal, but not “all-out” sprints, do not elicit an aversive perceptual response in recreationally-active adults. Secondary analyses suggest unique responses to SIE between men and women, which merits additional study to determine if sex impacts the chronic adaptation to sprint interval training.

Although the exercise intensity used in the SIE protocols in this study was supramaximal, the brief nature of the sprints resulted in relative peak $\dot{V}O_2$ values of 75% (LCE) and 88% (ACE) of $\dot{V}O_{2max}$, and peak HR of 90% of HRmax for both LCE and ACE. These values are similar to the cardiovascular stress associated with “vigorous exercise” according to the American College of Sports Medicine.⁴⁷ These HR values are also comparable to prior studies using low-volume “all-out” SIE, despite a much lower intensity.^{5,9} Nevertheless, contrary to our hypothesis, ACE exhibited significantly higher mean and peak % $\dot{V}O_{2max}$ than LCE. Prior data³² showed no difference in mean/peak $\dot{V}O_2$ or peak HR expressed as percentages of maximal values between HIIE (10 × 1 min at 75% PPO) performed using LCE and ACE, although mean HR was higher in response to LCE ([81% ± 5%] HRmax vs. [75% ± 7%] HRmax), which is similar to our data (Table 2).

One explanation of higher relative $\dot{V}O_2$ in response to ACE SIE is activation of accessory muscles, including the core and lower body, to assist the upper extremity in moving the pedal crank at high work rates. A secondary explanation of greater $\dot{V}O_2$ attendant with ACE SIE is incidence of a substantial $\dot{V}O_2$ slow component.⁴⁸ Compared to LCE, ACE is characterized by the use of a smaller muscle mass with a greater ratio of fast to slow twitch muscle fibers which leads to lower metabolic efficiency and higher $\dot{V}O_2$ at a given power output.^{27,49} When performing ACE in the severe intensity domain characteristic of SIE, it is possible that this slow component is augmented due to marked recruitment of fast twitch (FT) fibers, greater ventilation (Table 2), and greater disturbance of acid-base balance, all leading to a greater $\dot{V}O_2$ cost and in turn, propensity for fatigue. In addition, adults with greater FT ratio in the vastus lateralis exhibit a greater slow component than those with a preponderance of slow twitch (ST) fibers,⁴⁸ which would suggest that any muscle group having a higher ratio of FT fibers such as the upper

extremity should reveal a larger slow component during vigorous exercise. Finally, the greater relative $\dot{V}O_2$ with ACE could partly be related to differences between LCE versus ACE in the ramp test rather than the SIE sessions. However, as the $\dot{V}O_{2max}$ values obtained in the ACE ramp test are in effect 'submaximal', probably not limited by central factors, and closer to the arm muscles' 'true' maximal ability to take up oxygen, it is even more remarkable that relative $\dot{V}O_2$ during SIE is higher compared to LCE.

Our data suggest that SIE completed on the ACE imposes a greater cardiorespiratory demand expressed as % $\dot{V}O_{2max}$ than LCE (Table 2). Harvey et al.⁵⁰ required active men ($\dot{V}O_{2max}$ not measured) to perform the 30 s Wingate test using LCE and ACE. Results showed a greater aerobic contribution towards ATP supply for LCE versus ACE (17% vs. 11%), which had a significantly higher glycolytic (60% vs. 47%) contribution. In contrast, Price et al.²⁵ in active men with $\dot{V}O_{2max}$ equal to 34 mL·kg⁻¹·min⁻¹ and 48 mL·kg⁻¹·min⁻¹ on ACE and LCE showed a significantly higher aerobic contribution in response to the Wingate test performed with ACE compared to LCE (43% vs. 29%) which was consequent with a lower glycolytic contribution (39% vs. 68%). Nevertheless, in the latter study, a lower resistance was used (4% body mass vs. 5% body mass) which led to a significantly lower peak power output attained (6.9 W/kg vs. 9.8 W/kg) compared to the Harvey et al.⁵⁰ study. These methodological differences accompanied by discrepancies in calculation of the aerobic contribution likely mediate the different conclusions across studies.

Our data showing higher mean/peak % $\dot{V}O_{2max}$ and % \dot{V}_{Emax} in response to ACE corroborate prior work. Calbet et al.⁵¹ demonstrated more substantial cardiovascular strain during incremental ACE versus LCE, potentially because the $\dot{V}O_2$ attained is closer to the true maximal amount that can be taken up by the upper body. However, LCE is limited by the ability of the

cardiovascular system to deliver oxygen, which is not the case for ACE that is limited by peripheral factors. Zinner et al.²⁶ exhibited that six sessions of upper-body SIE in untrained men led to no difference in the increase in $\dot{V}O_{2\max}$ (9.8% vs. 6.1%, $p = 0.18$) than lower-body training, which was attendant with significant increases in Wingate-derived mean and peak power output and time trial performance. Their data also showed greater capillaries per fiber and a reduced oxygen deficit with upper-body training, suggesting that aerobic adaptations result from upper-body SIE, as repeatedly shown with LCE.^{3,8,10} However, there was no significant change in muscle fiber type ratio or B-HAD activity in the upper body despite a significant increase in citrate synthase activity,²⁷ so the specific adaptations mediating the increase in $\dot{V}O_{2\max}$ with upper-body SIE remain elusive. Further study is needed to elucidate if adaptations exhibited with upper-body training are associated with improved health status, and if higher relative mean and peak $\dot{V}O_2$ attendant with upper- versus lower-body SIE leads to a different chronic response, as it is possible that these localized, peripheral adaptations do not extend to better whole-body cardiometabolic health.

Our results show that peak RPE was equal to approximately 13 for both modalities, representing a ‘hard’ level of exertion. This value is lower than shown in studies using the REHIT protocol,⁷ the Tabata protocol,²⁰ and higher volume all-out SIE.^{20,36-37} The Wood et al.³⁷ study required active adults to perform eight 30 s intervals of LCE at 130% W_{\max} with 90 s recovery. Their peak HR (91%) is similar to that reported in the present study (90%, Table 2), so differences in HR do not explain the different RPE across studies. Nevertheless, participants underwent supramaximal “all-out” exercise with slightly longer interval durations and greater volume, likely augmenting the contribution of glycolysis contributing to higher BLa (~14 mM) and perceptions of fatigue, leading to a higher RPE. In the present study, the lack of difference

in peak values of HR as well as BLA, two known mediators of RPE, likely led to similar RPE between LCE and ACE.

Similar to RPE, our results showed no effect of exercise modality on affective valence. RPE, enjoyment, and pre-exercise affective valence are associated with the change in affective valence during acute interval exercise,⁵² so the lack of differences in these outcomes between modalities may partially explain this result. Our overall reduction in affective valence of ~ -2 units (i.e. ~ -0.5 units per sprint) is lower than what would be predicted based on results from a recent systematic review and meta-analysis showing that each additional “all-out” sprint in a SIE protocol elicits a ~ 1 -unit decrease in affective valence.¹³ This comparison supports our hypothesis that supramaximal but non-all-out SIE is perceived as less aversive compared with studies using all-out sprint protocols.^{20,37} Furthermore, our end-exercise value represented “fairly good” affective valence, showing that a low-volume SIE protocol requiring non-all-out sprints does not elicit aversive responses. In addition, despite our bouts requiring intensities above that associated with W_{max} , enjoyment was high and similar across modalities. Similar values (90–100) for enjoyment were shown in a recent study employing REHIT in adults with above and below average $\dot{V}O_{2max}$,⁵ although our values are higher than those revealed in inactive adults performing SIE (PACES = 83).⁴¹ Further study is merited to determine perceptual responses to similar SIE protocols in inactive adults and those with chronic disease to ascertain their feasibility as an alternative to aerobic exercise.

Our exploratory analysis demonstrates significantly higher HR in response to SIE on the cycle ergometer in women versus men. Although our study cannot identify the precise mechanism explaining this result, it may be related to the lower blood volume and left ventricular mass characteristic of women.⁵³ Hottenrott et al.⁵⁴ reported slower HR recovery to repeated Wingate

tests in women versus men, and since our protocol involves 8 min of recovery and only 80 s of work, this may explain some of our results. Nevertheless, recent data showed no sex difference in the hemodynamic and cardiovascular response (expressed as % maximal cardiac output [CO_{max}] and % $\dot{V}O_{2max}$) to three unique interval protocols performed on the cycle ergometer.⁵⁵ On the ACE, the only sex difference reported was mean % $\dot{V}O_{2max}$, which was higher in men compared to women. Potential explanations for this could be the greater upper-body muscle mass in men as well as their slower metabolic recovery to interval exercise versus women.⁵⁴ Additional investigations are needed which are adequately powered to discern potential sex differences in the physiological response to upper- and lower-body SIE.

This study has a few limitations. First, the participants included active, young, and non-obese adults naïve to SIE, so our data cannot be applied to inactive/obese populations or individuals who regularly perform these modalities. Second, our SIE protocol differed from those used in prior studies (e.g. multiple Wingate tests and Tabata), so our results are not entirely generalizable to studies using different SIE paradigms which have infinite permutations. Third, muscle fiber type differs between the upper and lower body⁵⁶ thus altering the $\dot{V}O_2$ and metabolic response to exercise, but this ratio was not determined in the present study. Fourth, despite preliminary data showing no difference in $\dot{V}O_{2max}$ between ACE and LCE when performed on the same day versus separate days, it is possible that $\dot{V}O_{2max}$ and W_{max} may have been slightly underestimated in our participants. Fifth, additional study is needed to compare responses between “all-out” and non-all-out SIE using these modalities. Lastly, no consideration of menstrual phase was made, and it is possible that hormone fluctuations may slightly impact our results. Due to known differences in body composition between men and women, additional work is needed to elucidate potential discrepancies in the cardiometabolic response to upper-

versus lower-body exercise between men and women. However, our study is strengthened by use of a large and heterogeneous sample, precise allocation of power output for all sessions, and use of a familiarization protocol which likely reduces learning effects and in turn augments the reliability of our data.

5. Conclusions

When performed regularly, SIE improves body composition and aerobic fitness yet requires a large degree of effort which can be unpleasant for many individuals. Our results show that supramaximal but non-all-out sprint interval exercise using the upper body is associated with lower absolute but greater relative cardiovascular demand versus lower body sprint interval exercise. In addition, affective valence was positive and post-exercise enjoyment was high. Clinicians may want to use low-volume SIE consisting of brief 20 s bouts that require non-all-out efforts to elicit more positive psychological responses than protocols requiring all-out sprints. In addition, upper-body sprint exercise leads to greater relative HR and $\dot{V}O_2$ versus leg cycling which may elicit a unique adaptive response.

Acknowledgements

The Authors acknowledge the participants for taking part in this study as well as Mr. Robert Sturdy for assistance with data collection.

Submission statement

All Authors have read and agree to this manuscript content. In addition, while this manuscript is being reviewed by this Journal, the manuscript will not be submitted elsewhere for review and publication.

Ethical approval

Participants provided written informed consent, and study experimental procedures were reviewed and approved by the Institutional Review Board at CSU—San Marcos (Protocol 1876593-1). The study was conducted in accordance with the Declaration of Helsinki.

Authors' contributions

TAA conceived the study, acquired ethical approval, recruited participants, collected data, analyzed data, and created the first and final draft of the manuscript. SP and MP recruited participants, collected data, assisted in data analysis, and edited the first and final draft of the manuscript. RM and NV edited the first and final draft of the manuscript.

Conflict of interest

The Authors report that no individual has any direct or indirect interest that is in direct conflict with conduction of this study.

References

1. Gillen JB, Gibala MJ. Is high-intensity interval training a time efficient exercise strategy to improve health and fitness? *Appl Physiol Nutr Metab*. 2014;39(3):409–412.
<https://doi.org/10.1139/apnm-2013-0187>
2. Astorino TA, Allen RP, Roberson DW, et al. Adaptations to high-intensity training are independent of gender. *Eur J Appl Physiol*. 2011;111(7):1279-1286.
<https://doi.org/10.1007/s00421-010-1741-y>

3. Burgomaster KA, Howarth KR, Phillips SM, et al. Similar metabolic adaptations during exercise after low volume sprint interval and traditional endurance training in humans. *J Physiol.* 2008;586(1):151–160. <https://doi.org/10.1113/jphysiol.2007.142109>.
4. Tabata I, Nishimura K, Kouzaki M, et al. Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and $\dot{V}O_{2\max}$. *Med Sci Sports Exerc.* 1996;28(10):1327-1330. <https://doi.org/10.1097/00005768-199610000-00018>
5. Astorino TA, Clausen RD, Marroquin J, Arthur B, Stiles K. Similar perceptual responses to reduced exertion high intensity interval training (REHIT) in adults differing in cardiorespiratory fitness. *Physiol Behav.* 2020;213:112687. <https://doi.org/10.1016/j.physbeh.2019.112687>
6. Cuddy TF, Ramos JS, Dalleck LC. Reduced exertion high-intensity interval training is more effective at improving cardiorespiratory fitness and cardiometabolic health than traditional moderate-intensity continuous training. *Int J Environ Res Publ Health.* 2019;16(3):483. <https://doi.org/10.3390/ijerph16030483>
7. Songsorn P, Brick N, Fitzpatrick B, et al. Affective and perceptual responses during reduced-exertion high-intensity interval training (REHIT). *Inter J Sport Exerc Psychol.* 2020;18(6):717-732. <https://doi.org/10.1080/1612197X.2019.1593217>
8. Hazell TJ, Macpherson RE, Gravelle BM, Lemon PW. 10 or 30-s sprint interval training bouts enhance both aerobic and anaerobic performance. *Eur J Appl Physiol.* 2010;110(1):153-160. <https://doi.org/10.1007/s00421-010-1474-y>
9. Metcalfe RS, Fitzpatrick B, Fitzpatrick S, et al. Extremely short duration interval exercise improves 24-h glycaemia in men with type 2 diabetes. *Eur J Appl Physiol.* 2018;118(12):2551-2562. <https://doi.org/10.1007/s00421-018-3980-2>

- 539 10. Gibala MJ, Little JP, Van Essen M, et al. Short-term sprint interval versus traditional
540 endurance training: similar initial adaptations in human skeletal muscle and exercise
541 performance. *J Physiol.* 2006;575(Pt3):901-911. <https://doi.org/10.1113/jphysiol.2006.112094>
- 542 11. Hardcastle SJ, Ray H, Beale L, Hagger MS. Why sprint interval training is inappropriate for
543 a largely sedentary population. *Front Psychol.* 2014;5:1505.
544 <https://doi.org/10.3389/fpsyg.2014.01505>
- 545 12. Biddle SJH, Batterham AM. High-intensity interval exercise training for public health: a big
546 HIT or shall we HIT it on the head? *Int J Behav Nutr Phys Act.* 2015;12:95.
547 <https://doi.org/10.1186/s12966-015-0254-9>
- 548 13. Metcalfe RS, Williams S, Fernandes G, et al. Affecting effects on affect: the impact of
549 protocol permutations on affective responses to sprint interval exercise; a systematic review and
550 meta-analysis of pooled individual participant data. *Front Sports Act Living.* 2022;4:815555.
551 <https://doi.org/10.3389/fspor.2022.815555>
- 552 14. Hu M, Jung ME, Nie J, Kong Z. Affective and enjoyment responses to sprint interval training
553 in healthy individuals: A systematic review and meta-analysis. *Front Psychol.* 2022;13:820228.
554 <https://doi.org/10.3389/fpsyg.2022.820228>
- 555 15. Islam H, Townsend LK, Hazell TJ. Modified sprint interval training protocols. Part I.
556 Physiological responses. *Appl Physiol Nutr Metab.* 2017;42(4):339-346.
557 <https://doi.org/10.1139/apnm-2016-0478>
- 558 16. Townsend LK, Islam H, Dunn E, Eys M, Robertson-Wilson J, Hazell TJ. Modified sprint
559 interval training protocols. Part II. Psychological responses. *Appl Physiol Nutr Metab.*
560 2017;42(4):347-353. <https://doi.org/10.1139/apnm-2016-0479>

17. Vollaard NBJ, Metcalfe RS. Into the health benefits of sprint interval training should focus on protocols with fewer and shorter sprints. *Sports Med.* 2017;47(12):2443-2451. <https://doi.org/10.1007/s40279-017-0727-x>
18. Ruffino JS, Songsorn P, Haggett M, et al. A comparison of the health benefits of reduced-exertion high-intensity interval training (REHIT) and moderate-intensity walking in type 2 diabetes patients. *Appl Physiol Nutr Metab.* 2017;42(2):202-208. <https://doi.org/10.1139/apnm-2016-0497>
19. Bayati M, Barzad B, Gharakhanlou R, Agha-Alinejad H. A practical model of low-volume high-intensity interval training induces performance and metabolic adaptations that resemble ‘all-out’ sprint interval training. *J Sports Sci Med.* 2011;10(3):571-576.
20. Follador L, Alves RC, Ferreira SDS, et al. Physiological, perceptual, and affective responses to six high-intensity interval training protocols. *Percept Mot Skills.* 2018;125(2):329-350. <https://doi.org/10.1177/0031512518754584>
21. Macpherson RE, Hazell TJ, Olver TD, Paterson DH, Lemon PW. Run sprint interval training improves aerobic performance but not maximal cardiac output. *Med Sci Sports Exerc.* 2011;43(1):115-122. <https://doi.org/10.1249/MSS.0b013e3181e5eacd>
22. Nightingale TE, Metcalfe RS, Vollaard NB, Bilzon JL. Exercise guidelines to promote cardiometabolic health in spinal cord injured humans: Time to raise the intensity? *Arch Phys Med Rehabil.* 2017;98(8):1693-1704. <https://doi.org/10.1016/j.apmr.2016.12.008>
23. Goodman LS, McKenzie DC, Nath CR, Schamberger W, Taunton JE, Ammann WC. Central adaptations in aerobic circuit versus walking/jogging trained cardiac patients. *Can J Appl Physiol.* 1995;20(2):178-197. <https://doi.org/10.1139/h95-013>

24. Brurok B, Helgerud J, Karlsen T, Leivseth G, Hoff J. Effect of aerobic high-intensity hybrid training on stroke volume and peak oxygen consumption in men with spinal cord injury. *Amer J Phys Med Rehabil*. 2011;90(5):407-414. <https://doi.org/10.1097/PHM.0b013e31820f960f>
25. Price M, Beckford C, Dorricott A, et al. Oxygen uptake during upper body and lower body Wingate anaerobic tests. *Appl Physiol Nutr Metab*. 2014;39(1):1345-1351. <https://doi.org/10.1139/apnm-2013-0405>
26. Zinner C, Morales-Alamo D, Ørtenblad N, et al. The physiological mechanisms of performance enhancement with sprint interval training differ between the upper and lower extremities in humans. *Front Physiol*. 2016;7:426. <https://doi.org/10.3389/fphys.2016.00426>
27. Sawka MN. Physiology of upper body exercise. *Exerc Sport Sci Rev*. 1986;14:175-211.
28. Jackson AS, Pollock ML Generalized equations for predicting body density of men. *Br J Nutr*. 1978;40(3):497-504. <https://doi.org/10.1079/bjn19780152>
29. Jackson AS, Pollock ML, Ward A. Generalized equations for predicting body density of women. *Med Sci Sports Exerc*. 1980;12(3):175-181.
30. Siri WE. Body composition from fluid space and density. In: J. Brozek J, Hanschel A Eds. *Techniques for measuring body composition*. National Academy of Science;1961:223-244.
31. Astorino TA, Robson T, McMillan DW. Classifying intensity domains from arm cycle ergometry differs versus leg cycling ergometry. *J Str Cond Res*. 2023;37(11):2192-2199. <https://doi.org/10.1519/JSC.00000000000004549>
32. Astorino TA, Emma D. Differences in physiological and perceptual responses to high intensity interval exercise between arm and leg cycling. *Front Physiol*. 2021;12:700294. <https://doi.org/10.3389/fphys.2021.700294>

- 605 33. Van Sumeren AL, Bye TK, Kilgas MA, Hartvigsen MM, Elmer SJ. Influence of the lower
606 body on seated arm cranking performance. *Int J Sports Med.* 2018;39(10):757-763.
607 <https://doi.org/10.1055/a-0633-9188>
- 608 34. Astorino TA, White AC, Dalleck LC. Supramaximal testing to confirm attainment of
609 $\dot{V}O_2$ max in sedentary men and women. *Int J Sports Med.* 2009;30(4):279-284.
610 <https://doi.org/10.1055/s-0028-1104588>
- 611 35. Astorino TA, Bediaamol N, Cotoia S, et al. Verification testing to confirm $\dot{V}O_2$ max attainment
612 in persons with spinal cord injury. *J Spinal Cord Med.* 2019;42(4):494-501.
613 <https://doi.org/10.1080/10790268.2017.1422890>
- 614 36. Olney N, Wertz T, LaPorta Z, Mora A, Serbas J, Astorino TA. Comparison of acute
615 physiological and psychological responses between moderate intensity continuous exercise and
616 three regimes of high intensity training. *J Str Cond Res.* 2018;32(8):2130-2138.
617 <https://doi.org/10.1519/JSC.0000000000002154>
- 618 37. Wood KM, Olive B, LaValle K, Thompson H, Greer K, Astorino TA. Dissimilar
619 physiological and perceptual responses between sprint interval training and high-intensity
620 interval training. *J Str Cond Res.* 2016;30(1):244-250.
621 <https://doi.org/10.1519/JSC.0000000000001042>
- 622 38. Borg G. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc.* 1982;14(5):377-
623 381.
- 624 39. Hardy CJ, Rejeski WJ. Not what, but how one feels: the measurement of affect during
625 exercise. *J Sport Exerc Psychol.* 1989;11(3):304-317. <https://doi.org/10.1123/JSEP.11.3.304>
- 626 40. Kendzierski D, DeCarlo KJ. Physical activity enjoyment scale: Two validation studies. *J*
627 *Sport Exerc Psychol.* 1991;13(1):50-64. <https://doi.org/10.1123/JSEP.13.1.50>

- 628 41. Stork MJ, Gibala MJ, Martin Ginis KA. Psychological and behavioral responses to interval
629 and continuous exercise. *Med Sci Sports Exerc.* 2018;50(10):2110-2121.
630 <https://doi.org/10.1249/MSS.0000000000001671>
- 631 42. Thum JS, Parsons G, Whittle T, Astorino TA. High-intensity interval training elicits higher
632 enjoyment than moderate intensity continuous exercise. *PLoS One.* 2017;12(1):e0166299.
633 <https://doi.org/10.1371/journal.pone.0166299>
- 634 43. Cohen, J. *Statistical Power Analysis for the Behavioral Sciences.* 2nd ed. Lawrence Erlbaum
635 Associates;1988.
- 636 44. Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: A flexible statistical power analysis
637 program for the social, behavioral, and biomedical sciences. *Behav Res Methods.*
638 2007;39(2):175-191. <https://doi.org/10.3758/bf03193146>
- 639 45. Kaminsky LA, Imboden MT, Arena R, Myers J. Reference standards for cardiorespiratory
640 fitness measured with cardiopulmonary exercise testing using cycle ergometry: data from the
641 fitness registry and the importance of exercise national database (FRIEND) registry. *Mayo Clin*
642 *Proc.* 2017;92(2):228-233. <http://doi.org/10.1016/j.mayocp.2016.10.003>
- 643 46. Pivarnik JM, Grafner TR, Elkins EB. Metabolic, thermoregulatory, and psychophysiological
644 responses during arm and leg exercise. *Med Sci Sports Exerc.* 1988;20(1):1-5.
645 <https://doi.org/10.1249/00005768-198802000-00001>
- 646 47. Garber CE, Blissmer B, Deschenes MR et al. Quantity and quality of exercise for developing
647 and maintaining cardio-respiratory, musculoskeletal, and neuromotor fitness in apparently
648 healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc.* 2011;43(7):1334-1359.
649 <https://doi.org/10.1249/MSS.0b013e318213fefb>

- 650 48. Barstow TJ, Jones AM, Nguyen PH, Casaburi R. Influence of muscle fiber type and pedal
651 frequency on oxygen uptake kinetics of heavy exercise. *J Appl Physiol*. 1996;81(4):1642–1650.
652 <https://doi.org/10.1152/jappl.1996.81.4.1642>
- 653 49. Toner MM, Sawka MN, Levine L, Pandolf KB. Cardiorespiratory responses to exercise
654 distributed between the upper and lower body. *J Appl Physiol*. 1983;14(5):1403-1407.
655 <https://doi.org/10.1152/jappl.1983.54.5.1403>
- 656 50. Harvey L, Wiegand A, Solomon C, McClellan C, Lovell DI. A comparison of upper and
657 lower-body energetics during high intensity exercise. *J Sports Med Phys Fit*. 1995;55(7-8):708-
658 713.
- 659 51. Calbet JAL, Gonzalez Alonso J, Helge J, et al. Central and peripheral hemodynamics in
660 exercising humans: leg vs. arm exercise. *Scand J Med Sci Sports*. 2015;25(S4):144-157.
661 <https://doi.org/10.1111/sms.12604>
- 662 52. Astorino TA, Vella CA. Predictors of change in affect in response to high intensity interval
663 exercise (HIIE) and sprint interval exercise (SIE). *Physiol Behav*. 2018;196:211–217.
664 <https://doi.org/10.1016/j.physbeh.2018.08.017>
- 665 53. de Simone G, Devereux RB, Daniels SR, Meyer RA. Gender differences in left ventricular
666 growth. *Hypertension*. 1995;26(6 Pt 1):979-983. <https://doi.org/10.1161/01.hyp.26.6.979>
- 667 54. Hottenrott L, Mohle M, Ide A, Ketelhut S, Stoll O, Hottenrott K. Recovery from different
668 high-intensity interval training protocols: comparing well-trained women and men. *Sports*
669 2021;9(3):34. <https://doi.org/10.3390/sports9030034>
- 670 55. Coe LN, Astorino TA. Sex differences in hemodynamic response to high-intensity interval
671 exercise. *Scand J Med Sci Sports*. 2023 (in press). <https://doi.org/10.1111/sms.14495>

56. Gollnick PD, Armstrong RB, Saubert CW, Piehl K, Saltin B. Enzyme activity and fiber composition in skeletal muscle of untrained and trained men. *J Appl Physiol.* 1972;33(3):312-319. <https://doi.org/10.1152/jappl.1972.33.3.312>

Figure Legends

1. Study Flow Diagram. LCE = leg cycling ergometry; ACE = arm cycling ergometry; SIE = sprint interval exercise

2. Changes in a) oxygen uptake, b) heart rate, c) ventilation, and d) respiratory exchange ratio in response to sprint interval exercise (SIE) performed using leg cycling ergometry (LCE) and arm cycling ergometry (ACE). Data are mean \pm SD; * = $p < 0.05$ between LCE and ACE

3. Change in a) blood lactate concentration, b) rating of perceived exertion, and c) affective valence in response to sprint interval exercise (SIE) performed using leg cycling ergometry (LCE) and arm cycling ergometry (ACE). Data are mean \pm SD; * = $p < 0.05$ between LCE and ACE

Table 1: Comparison of data from $\dot{V}O_{2\max}$ testing between leg and arm cycling ergometry (mean \pm SD)

Parameter	LCE	ACE	<i>p</i> value	Cohen's <i>d</i>
$\dot{V}O_{2\max}$ (mL·kg ⁻¹ ·min ⁻¹)	39.4 \pm 7.4	27.1 \pm 4.7	< 0.001	2.0
$\dot{V}O_{2\max}$ (L·min ⁻¹)	2.9 \pm 0.7	2.0 \pm 0.5	< 0.001	1.5
PPO (W)	272.1 \pm 57.4	132.5 \pm 36.7	< 0.001	3.0
HR _{max} (b/min)	184.4 \pm 10.0	178.0 \pm 14.4	0.001	0.6
RER _{max}	1.26 \pm 0.09	1.23 \pm 0.09	0.027	0.3
$\dot{V}_{E\max}$ (L·min ⁻¹)	121.6 \pm 31.7	90.7 \pm 24.3	< 0.001	1.1
BLa (mM)	11.0 \pm 1.9	9.7 \pm 2.3	0.002	0.7
Duration (min)	8.9 \pm 1.2	8.8 \pm 1.5	0.76	0.1
RPE (6 - 20)	16.5 \pm 2.2	15.3 \pm 3.4	0.10	0.4
Affect (+5 to -5)	0.7 \pm 2.7	0.2 \pm 2.4	0.42	0.2

$\dot{V}O_{2\max}$ = maximal oxygen uptake; PPO = peak power output; HR = heart rate; RER = respiratory exchange ratio; \dot{V}_E = ventilation; BLa = blood lactate concentration; RPE = rating of perceived exertion

Table 2: Comparison of mean and peak physiological responses during sprint interval exercise performed using leg and arm cycle ergometry (mean \pm SD).





Parameter	LCE	Range	ACE	Range	<i>p</i> value	Cohen's <i>d</i>
Mean $\dot{V}O_2$ (L·min ⁻¹)	1.59 \pm 0.34*	1.02 – 2.18	1.24 \pm 0.31	0.64 – 1.83	< 0.001	1.08
Mean $\dot{V}O_2$ (% $\dot{V}O_{2max}$)	56.9 \pm 7.2*	42 - 69	61.7 \pm 8.6	50 - 84	0.015	0.63
Peak $\dot{V}O_2$ (L·min ⁻¹)	2.10 \pm 0.44*	1.60 – 2.90	1.79 \pm 0.48	0.94 – 2.78	< 0.001	0.70
Peak $\dot{V}O_2$ (% $\dot{V}O_{2max}$)	75.4 \pm 9.7*	60 - 96	88.2 \pm 10.4	68 - 110	< 0.001	1.33
Mean HR (b/min)	148 \pm 15*	123 - 172	136 \pm 16	112 - 173	< 0.001	0.81
Mean HR (%HR _{max})	80.0 \pm 6.4*	69 - 89	76.6 \pm 6.1	66 – 90	0.038	0.58
Peak HR (b/min)	167 \pm 13*	142 – 188	160 \pm 17	133 - 188	0.014	0.49
Peak HR (%HR _{max})	90.2 \pm 5.0	81 - 98	90.0 \pm 7.0	76 - 99	0.71	0
Mean \dot{V}_E (L·min ⁻¹)	57 \pm 10*	35 - 79	48 \pm 12	20 - 78	< 0.001	0.83
Mean \dot{V}_E (% \dot{V}_{Emax})	51 \pm 10*	26 - 56	57 \pm 14	40 - 90	0.047	0.50
Peak \dot{V}_E (L·min ⁻¹)	80 \pm 13*	46 - 111	67 \pm 20	25 - 130	0.008	0.79
Peak \dot{V}_E (% \dot{V}_{Emax})	71 \pm 15*	48 - 102	81 \pm 20	55 - 120	0.017	0.64
Mean RER	1.20 \pm 0.11	1.04 – 1.38	1.21 \pm 0.08	1.06 – 1.35	0.81	0.12
Peak RER	1.45 \pm 0.16	1.21 – 1.84	1.47 \pm 0.17	1.24 – 1.71	0.65	0.11




LCE – leg cycle ergometry; ACE = arm cycle ergometry; $\dot{V}O_2$ = oxygen uptake; HR = heart rate; \dot{V}_E = ventilation; RER = respiratory exchange ratio; * = $p < 0.05$ versus ACE

25 participants provide consent and complete incremental exercise on LCE and ACE

Familiarization session of SIE using LCE and ACE

25 participants complete SIE using LCE and ACE on separate days





Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

NA