

Research

Improving the mental health of farmers: what types of remote support are acceptable, feasible, and improve outcomes? A feasibility RCT

Kate Lamont¹ · Hugo C. van Woerden² · Emma King³ · Charlotte Wendelboe-Nelson³ · Roger W. Humphry¹ · Cameron Stark² · Chris Williams⁴ · Margaret Maxwell³

Received: 22 July 2023 / Accepted: 6 December 2023

Published online: 04 January 2024

© The Author(s) 2023 [OPEN](#)

Abstract

Background The farming community have high rates of poor mental health, and are relatively ‘hard to reach’ with mental health services. The aim of this study was therefore to undertake a feasibility RCT, based on two mental health interventions. These were (1) CBT based ‘Living Life to the Full for Farming Communities’ (LLTTF-F; www.lltff.com), and (2) a holistic social and emotional support service delivered by the Royal Scottish Agricultural Benevolent Institution (RSABI). The feasibility was supplemented by process evaluation.

Methods This feasibility study aimed to recruit 40 individuals from the farming community who were experiencing a common health problem defined as a score of ≥ 8 on PHQ-9. A snowball approach was used to recruit interested individuals who had an association with farming. An initial telephone call screened for eligibility and obtained consent to randomisation to the two specified interventions, or to a thirdly group receiving a combination of both LLTTF-F and ‘Social and emotional support’. Participants were permitted to override the randomised option if they expressed a strong preference before the interventions began.

Results Thirty-two participants provided baseline and three-month data. All three interventions showed positive improvements on PHQ-9 scores as follows: the ‘combined intervention’ mean baseline score was 18.1 compared to 12.0 at 3-month follow-up (mean change 6.1). ‘Social and emotional support’ mean baseline score was 11.3 compared to 6.7 at 3-month follow-up (mean change 4.6). ‘LLTTF-F CBT-based intervention only’ mean baseline score was 11.8 compared to 4.5 at 3-month follow-up (mean change 7.3). The retention rate was 81% at three months.

In a sub-group of the LLTTF-F CBT-based intervention online materials were supplemented by telephone guided support. This approach received very positive feedback.

Conclusions Recruitment from the farming community required intense effort, and good engagement can then be retained for at least three months. There is evidence that the interventions used were feasible, and tentative evidence that they had a demonstrable effect on mental wellbeing, with the LLTTF providing the largest effect on PHQ-9 scores.

Trial Registration Number ISRCTN27173711, submitted 25/08/2023, confirmed 22/09/2023.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s44192-023-00054-1>.

✉ Hugo C. van Woerden, hugo.vanwoerden@uhi.ac.uk; ✉ Margaret Maxwell, margaret.maxwell@stir.ac.uk; Kate Lamont, Kate.Lamont@sruc.ac.uk; Emma King, emma.king@stir.ac.uk; Charlotte Wendelboe-Nelson, charlotte.wendelboe-nelson2@stir.ac.uk; Roger W. Humphry, Roger.Humphry@sruc.ac.uk; Cameron Stark, Cameron.stark@uhi.ac.uk; Chris Williams, Chris.Williams@glasgow.ac.uk | ¹Scotland’s Rural College (SRUC), Scotland, UK. ²University of the Highlands and Islands, Scotland, UK. ³University of Stirling, Scotland, UK. ⁴Glasgow University, Scotland, UK.



Abbreviations

LLTTF-F Living Life to the Full for Farming Communities

RSABI Royal Scottish Agricultural Benevolent Institution

1 Introduction

Recent systematic reviews indicate that farmers are at particular risk of mental health disorders [1, 2]. Depression in the farming community is increasing and suicide rates are among the highest in any occupational group in the UK [3]. Depression in farmers increases with age, and is higher in males, and symptoms of depression are associated with a 13% increase in sex-adjusted and age-adjusted mortality [4–7]. Farmers are also more likely to report thinking that ‘life is not worth living’. It is currently estimated that one agricultural worker per week takes their own life in the UK [3]. Risk of suicide is also higher amongst those working in specific agricultural roles such as harvesting crops and rearing animals (almost twice the national UK average) [3].

Throughout this study, we refer to farmers as including all farm workers and those who are part of the wider agricultural community, including unpaid workers and family members. This includes those who are part time farmers who simultaneously have some paid employment.

Social factors and community characteristics have an important mediating role in the mental health of the farming population [8, 9]. Farmers are more likely to turn to their own communities for support than to health or social work authorities, with many preferring to engage with advice from respected members within their communities, such as veterinary surgeons (Vets), or use anonymous sources of support such as the internet or self-help booklets [10, 11].

CBT based interventions have a strong evidence base for the management of common mental health problems [12] and are often well accepted in rural settings [13, 14]. CBT has the advantage that it can be completed in a relatively short period of time compared with other talking therapies. Its structured nature means it can be provided in different formats (including remote and self-directed formats), and it teaches practical strategies that can be used in everyday life. These qualities may appeal to farmers.

Psychological and social interventions provide different types of support, and combined approaches may be optimal [15, 16]. Our two candidate interventions therefore included the CBT based ‘Living Life to the Full for Farming Communities’ (LLTTF-F) and the Royal Scottish Agricultural Benevolent Institution’s (RSABI) social and emotional support service (see below). These interventions were delivered separately or in combination providing three arms to our RCT.

1.1 Assessing feasibility

Before undertaking a full-sized RCT, the preferences of farmers for interventions, feasibility, and uptake, need to be established. This project therefore sought to establish a ‘best-candidate’ intervention to incorporate in a larger RCT by: identifying a preferred candidate intervention for addressing mental health problems in farmers; assessing the potential to recruit farmers; and testing a potential study outcome measure (PHQ-9).

The study approach was informed by focus groups with farmers, Vets, and other farm advisors. The qualitative aspects of this study have been published elsewhere [17].

The holistic social and emotional support service (including home visits and/or telephone support) was provided by RSABI. The self-help CBT based intervention (LLTTF-F), was based on previous research [18, 19]. LLTTF-F can be delivered on-line or in booklet format, and can be provided with or without telephone guided support. The ‘counselling’ aspect of the social support intervention, was primarily a ‘listening ear’, whereas the LLTTF-F CBT, was a structured and theoretically (cognitive behavioural therapy) based approach with support to engage with the intervention through problem solving techniques.

2 Methods

A mixed-methods approach was used to assess the preferences of farmers. This was considered important to establish feasible parameters to include in larger ‘best-candidate’ RCT.

2.1 Theoretical framework

The study's theoretical framework aligns with 'pragmatism' as a research philosophy, where the research question is the important determinant of methods. This combines both positivist and interpretivist positions within the scope of a single study, depending on the nature of the research questions [20, 21]. The study also adopted an approach that was consistent with the sentiments of 'participatory action research', with a focus was on transforming the lives of socially marginalised populations. Participatory action research was particularly used to engage an advisory/reference group using appreciative enquiry [22]. More detail on the advisory group and qualitative aspects of the study are provided elsewhere [17].

2.2 Sample size

A sample size calculation, was conducted on the basis of an unpaired t-test, with the required number of observations in each arm of the study, based on an effect size from a published paper reporting a Cohen's difference, d , of 0.66 on the EQ-VAS scale [23]. A statistical power of 80%, and a statistical significance threshold of 5% was assessed within the statistical software R, based on the `pwr.t.test()` function within the package "pwr" [24]. The sample size calculation recommended 42 observations in each arm of the hypothetical study considered. However, given the practical logistics of a feasibility study, a decision was made to recruit a smaller sample of 40 individuals spread across all three arms of the study. This took into account advice from the study's statistician (RH). Participants were drawn from Scotland and northern England.

2.3 Inclusion and exclusion criteria

The Inclusion criteria were: 18 years and over; a member of the farming community; and experiencing mental health problems based on the completion over the phone of a PHQ-9 questionnaire and a score of ≥ 8 [25, 26].

Exclusion criteria were: those considered at baseline to be at significant risk of suicide based on a phone interview; currently undertaking or having engaged with CBT or other psychotherapy within the past 6 months; unable to communicate in English; or unable to give informed consent.

2.4 Recruitment

Recruitment was conducted via signposting by local Vets, farming social media, farmer support organisations (but excluding RSABI, who were leading on the provision of the alternate intervention, so as not to engender bias in participant preferences), and farming and mental health charities and support organisations. Flyers and posters were also used, for example, for recruitment at farmers marts. Interested participants were contacted by a researcher directly (phone or e-mail), according to their preference, and provided with a study information pack. If individuals consented to participate, an initial telephone screening call was undertaken to assess eligibility and to obtain informed consent for inclusion. Baseline measures were collected (see Table 1) alongside demographic characteristics (age, sex, ethnicity, educational attainment, employment status/job role). The number of data points assessed were kept to a minimum on advice from the focus groups and farming advisory group.

2.5 Randomisation

Randomisation was undertaken by full randomisation of the order of subjects, without any blocking, using the rand function within MS Excel. The randomisation options were: (1) to receive the on-line LLTTF-F CBT intervention; or (2)

Table 1 Study data collection points

	Eligibility	Baseline	3 months	6 months
Demographic characteristics	X			
PHQ-9 and suicidal ideation	X	X	X	X
3 item sense of coherence		X	X	X
EQ-5D		X	X	X
Engagement with the interventions			X	X

the RSABI social and emotional support intervention); or (3) to receive both interventions combined. Participants were allowed to switch intervention, if they indicated that they had a strong preference to do so before commencement of the intervention to which they had been randomised. All participants were offered this choice. The participants that changed intervention are shown in the study flowchart (Fig. 1). We have undertaken 'per protocol' analysis rather than intention to treat.




Fig. 1 Flow chart of participants

2.6 Living life to the full for farming communities intervention (\pm telephone support)

This intervention was based on the Living Life to the Full CBT on-line intervention, which was adapted for the farming community. The resource was provided by Five Areas Ltd, which is associated with one of the authors (CW). The programme focuses on educational online life-skills resources for people experiencing stress, anxiety and depression. It was available to participants as an online resource or as a booklet. The intervention used accessible language and included self-guided behavioural change tools.

Participants worked at their own pace but were encouraged to complete the intervention within eight weeks. The on-line version tracked module participation and completion, and additional data such as pages accessed, frequency of access, and for how long etc.

LLTTF participants were given the option of telephone support, consisting of an initial 30-min session, followed by three additional support sessions. Telephone support sessions followed a standard support algorithm and were provided by trained and supervised coaches.

2.7 The social and emotional support intervention

RSABI had an established intervention, which supported people from the Scottish agriculture community emotionally, practically and financially in times of need, providing a holistic service for individuals from the farming community to help them move forward in difficult times, for example, by supporting business reviews, providing access to counselling and providing help with essential living expenses. RSABI offered a home visit service, and a helpline staffed by mental health first aid trained staff and volunteers from 7 am to 11 pm, 365 days of the year, and a 'call out' service. Those receiving the RSABI holistic social and emotional support service, had the method of both initial and further contact recorded (telephone, on-line, home visit), along with the number of contacts, and, the types of help accessed (e.g., financial), although information on the results of this contact is not reported in detail in this paper.

2.8 Consent

To ensure support for vulnerable participants, we obtained prior consent from all participants to contact their GP if participants were considered at risk of suicide or showed evidence of deteriorating mental health as identified by their PHQ-9 scores. All participants were also made aware of helplines (such as the Samaritans and Breathing Space).

2.9 Duration of follow up

Baseline measurements were undertaken by telephone (or face to face if this was requested and a visit was feasible), as were follow up measurements at 3 months, and where possible at 6 months. A proportion of participants could not be followed up at 6 months because the study ran out of time before those who had been recruited late in the project reached their 6 month milestone. This was in large measure due to the impact of the COVID-19 pandemic.

2.10 Outcome measures

The pre-agreed primary outcome for the RCT was the change in the measure of depression and anxiety assessed by a difference of mean scores within groups and between groups. Secondary outcomes, related to Sense of Coherence, and EQ-5D, will be reported elsewhere.

Process evaluation outcomes were: recruitment, retention, and engagement with the different interventions. Three engagement questions were asked: Q1. *Did you find the intervention you received via this study helpful?* (Not at all useful, Of small benefit, Not sure, Quite helpful, Very helpful). Q2. *Did you follow the course/advice as instructed?* (No I did not, To a small degree, To a reasonable degree/mostly, Always). Q3. *Would you say you completed the course/sessions that were offered?* (Not at all/not much, Only partly/to a small degree, Mostly, Yes as instructed/offered).

2.11 Ethical approval

Ethical approval was granted by Stirling University General University Ethics Panel Reference: GUEP (19 20) 901. This study was performed in line with the principles of the Declaration of Helsinki. Informed consent was obtained from all participants. The intervention is reported in accordance with CONSORT guidelines (See Supplementary Material for assessment of the study against the guideline's criteria).

3 Results

3.1 Changes to PHQ-9 scores

All three interventions showed a positive change in PHQ-9 scores with the LLTTF-F CBT-based intervention showing most improvement as follows: Combined intervention mean baseline score was 18.1 compared to 12.0 at 3-month follow-up (mean change 6.1); Telephone accessed social and emotional support mean baseline score was 11.3 compared to 6.7 at 3-month follow-up (mean change 4.6); CBT-based intervention only mean baseline score was 11.8 compared to 4.5 at 3-month follow-up (mean change 7.3).

A summary of baseline demographic features is provided in Table 2. Ethnicity is not shown in the table as all participants were White Scottish or British.

3.2 Recruitment

We recruited and subsequently randomised 32 out of our target of 40 participants (80% of target). Face to face recruitment by someone with recognised farming or rural links was considered as the most successful mechanism. Engaging farmers in research was possible but required significant, focused effort.

Twenty nine of the 42 participants screened for eligibility met our initial inclusion criteria (eligibility of 69%). Based on researcher discussion, three participants initially classed as 'ineligible' were subsequently included as 'borderline eligible' and included in the study. Allocation details are shown in Fig. 1.

3.3 Retention

Retention rates were good, as 26 of the 32 participants (81%) provided follow-up data at three months, and 16 out of the 22 participants who were contacted at six months (73%). The main reason that only 22 out of 32 participants completed six-month questionnaires was that ten participants (31%) had been participating in the study for less than six months when the study ended, as the project data collection stage had run out of time. The late recruitment of these 10 participants to the study was related to project delays due to COVID-19 pandemic restrictions that were nationally applied to research activity.

3.4 Allocation

Initial random allocation was fairly evenly distributed: emotional/social support ($n = 10$); on-line/booklet LLTTF-F CBT \pm telephone support ($n = 10$); and combined ($n = 12$). However, allowing for a preference to be expressed resulted in final allocations of: emotional/social support ($n = 7$); on-line/booklet LLTTF-F \pm telephone support ($n = 17$); and combined intervention ($n = 8$). In summary, more farmers indicated a preference to switch to the LLTTF-F CBT-based intervention than the emotional/social support intervention, or both interventions combined.

3.5 Engagement

Assessment of engagement indicated that all the interventions were acceptable, but suggested that engagement could be enhanced further in future studies. Maximal engagement with the support materials was primarily prevented by

Table 2 Demographic characteristics by intervention (after personal preferences were incorporated)

Intervention category	Number of participants	Mean age (yrs)	Sex	What proportion of your income is derived from farming?				Which of these options best describes your formal education?			
				Female	> 10%	10–50%	> 50%	Primary/secondary school	Secondary advanced / vocational / further education	University/ postgraduate / professional	
LLTTF-F CBT	8	51.6	6 (75%)	2 (25%)	2 (25%)	1 (12.5%)	5 (62.5%)	1 (12.5%)	5 (62.5%)	2 (25%)	
Social and emotional support	7	53.6	6 (100%)	0 (0%)	0 (0%)	1 (17%)	5 (83%)	3 (43%)	2 (28.5%)	2 (28.5%)	
Both interventions	17	52.2	13 (76.5%)	4 (23.5%)	4 (22%)	1 (5.5%)	13 (72.5%)	1 (6%)	7 (41%)	9 (53%)	

participant's perceived 'lack of time', with service users stating that this was 'too much on top of everything else' in their lives. Guided, personal telephone support was very valued, especially in the early phase of engagement with the support material.

3.6 Engagement with the interventions

Of those receiving the combined intervention: 3/5 reported following advice/course to a small degree, and 2/5 mostly/always followed advice/course. All completed the course at least 'to a small degree'. On-line usage data showed that an average of nine modules were accessed.

Of those receiving telephone accessed social and emotional support, 4/5 reported following advice/course mostly as instructed, and 1/5 to 'a small degree'. In answer to the question "Would you say you completed the course/sessions that were offered?" 2/5 'mostly' completed the sessions that were offered, 2/5 to 'a small degree' and 1/5 'not at all/not much'. The mean time provided by the telephone accessed social and emotional support service was 4 h 17 min (median 2 h 15 min) and the average number of contacts overall was 29 contacts per participant (median 23). The majority (4/6 responders) felt they had followed the advice received.

Of those receiving the LLTTF-F—intervention only, 3/13 reported following advice/course to a small degree, 5/13 mostly/always followed advice/course, and 5 reported they did not follow advice/course. Four completed the course 'to a small degree', 5 completed 'mostly as instructed' and 4 followed the course 'not at all/not much'. On-line usage data from 9/13 participants showed an average of 6 modules accessed.

3.7 Effect of telephone support

More positive support was expressed for the LLTTF-F CBT based intervention when it was accompanied with telephone support, as 23/25 (92%) of those who received the LLTTF-F intervention opted for telephone support. In addition, 5/25 (25%) requested booklets to supplement, or to provide an alternative, to the online LLTTF-F.

3.8 Summary of process evaluation

A key finding is that the two interventions appeared to meet different needs. The CBT intervention had the largest effect on PQ-9 scores (in this small sample), whilst the practical support offered within the holistic social support intervention was particularly valued for other aspects, such as helping to make difficult business decisions.

4 Discussion

All the interventions resulted in reductions in PHQ-9 scores, providing some indication that the various components of the two interventions were of benefit. This was particularly the case for the CBT intervention, where there was a mean fall in PHQ-9 scores of 7.3 (from 11.8 to 4.5). This suggests that CBT may be best at addressing immediate the specific mental health issues. However, such an approach would miss out on aspects of the social support that were very highly valued, and which it is possible to speculate may help to prevent later mental health problems.

Perhaps an optimised intervention for the farming community, based on the study data on engagement, and published elsewhere [17], would be a combination of supported (and possible face-to-face) start-up session(s) alongside a CBT-based intervention. There was limited engagement with written material by some participants, and it is possible to speculate that supplementary video and audio content (such as podcasts) might be helpful, and might reflect farmers lifestyle/ Listening to material whilst working might address a perceived 'lack of time'. Practical support for farm related problems is valued but resource intensive.

4.1 Strengths and weaknesses

A number of lessons were learned from process measures to inform a larger RCT. We recruited > 70% of target (80% achieved, or 73% when removing the three who were initially 'borderline eligible' who were still entered into the study. However, recruitment was labour intensive. We met our target of $\geq 40\%$ of contacts being eligible (62% eligible), and

our target of < 30% attrition (only 4 participants withdrew (12.5%), with a further 2 (6%) classed as 'unable to contact'), leading to overall attrition of 16% at 3 months and 27% at 6 months.

The sample size was small, and we have therefore not undertaken statistical analysis beyond the difference in baseline and three-month PHQ-9 scores. Given the sample size, predominant male gender and white Scottish or British ethnicity, caution should be applied in terms of generalisability to other settings. For logistical and ethical reasons, it was not possible to include a control and this is a weakness that should be considered in further studies.

We recognise that a feasibility study does not require a power calculation, and that our approach to sample size is open to critique. However, given the emphasis on the CONSORT checklist on a power calculation, this was undertaken.

The study would have been strengthened by using a formal structure to assess the components of the low intensity support that was provided by RSABI volunteers, in line with the taxonomy developed by Glasgow and Rosen[27].

There was significant variation in baseline PHQ-9 baseline scores. This may simply be as a result of the small sample size, but it is possible to speculate that it may also indicate some self-selection by participants with different types of needs. The PHQ-9 tool has the advantage of being shorter than HADS, which was initially considered as a measure of mental health. Discussions with our project reference group (including farmers and farmer representatives) concluded that the PHQ-9 was considered to be more acceptable to individuals in the farming and crofting community, in terms of its comprehension and wording, and therefore also for telephone completion. The PHQ-9 does include a question on fatigue, and as farmers work very long hours, this question may have picked up fatigue that was not associated with mental health.

The approach that we took did not include a control group. A waiting list control design would have been an alternative approach, which would have provided a control arm to the study. However, this study was a feasibility study which also focused on establishing 'preferences' and therefore studying preferences and engagement with the interventions was prioritised.

4.2 Comparison with other literature

Interactive computerised cognitive behaviour therapy has been found to be acceptable in U.S. rural communities in relation to privacy, accessibility, user-friendliness and cultural appropriateness [13]. However, there is little current knowledge concerning preferences and acceptability or up-take of remote interventions in a UK context, and how these can best be signposted to farmers. This study indicates that people in the farming community in Scotland experiencing mental health issues or personal stresses may be willing to engage with online material, but there is still a significant appetite for supplementary paper-based resources.

There has been a recent encouraging growth in studies addressing mental health and suicide prevention in farming communities, with the publication of a range of qualitative studies in different parts of the world [14, 28–31], and a number of intervention studies [32–34]. These studies suggest the benefits of multi-modal approaches that are tailored to the farming community, as the aetiology of problems in this community is diverse. Supportive interventions clearly need to be incorporated into national strategy on farming and agricultural community development [35–37].

5 Conclusion

This study has provided evidence that those from the farming community can be recruited, although recruitment requires intense effort, and can be retained in an RCT for at least three months. There is evidence that the interventions used were feasible, and tentative evidence that these had a demonstrable benefit to mental wellbeing, using standard available measurement tools.

Acknowledgements The authors are grateful for advice and input from Jo Baughan, Chole McCulloch, Jim Hume, Chris McVey, Kate Richards and the Scottish Rural Mental Health Forum.

Author contributions The study was designed by KL and MM; RH undertook a power calculation; KL, MM, HvW, CW and CS helped develop the methods; KL undertook the interviews. HvW wrote the first draft of the paper, and all authors read and approved the final manuscript.

Funding This research was funded by the Chief Scientists Office, Scottish Government (HIPS 19/51).

Data availability The datasets generated and/or analysed during the current study are available from the authors on reasonable request.

Code availability Not applicable.

Declarations

Ethics approval and consent to participate The study was approved by the University of Stirling General University Ethics Panel Reference: GUEP (19 20) 901. The study was undertaken in line with the Declaration of Helsinki. Written informed consent was obtained from all participants in the study.

Consent for publication Not applicable.

Competing interests Chris Williams is the Director and shareholder of Five Areas Limited, and is author of the LLTTF-F course and linked booklets (www.lltff.com; <https://lltff.com/resources/lltff-farming/>), one of the interventions used by this project. All the other authors state that they have no competing interests to declare.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Daghighi Yazd S, Wheeler SA, Zuo A. Key risk factors affecting farmers' mental health: a systematic review. *Int J Environ Res Public Health*. 2019;16(23):4849.
2. Younker T, Radunovich HL. Farmer mental health interventions: a systematic review. *Int J Environ Res Public Health*. 2021;19(1):244.
3. Meltzer H, Griffiths C, Brock A, Rooney C, Jenkins R. Patterns of suicide by occupation in England and Wales: 2001–2005. *Br J Psychiatry*. 2008;193(1):73–6.
4. Thomas H, Lewis G, Thomas DR, Salmon R, Chalmers R, Coleman T, Kench S, Morgan-Capner P, Meadows D, Sillis M. Mental health of British farmers. *Occup Environ Med*. 2003;60(3):181–6.
5. Torske MO, Hilt B, Glasscock D, Lundqvist P, Krokstad S. Anxiety and depression symptoms among farmers: the HUNT study, Norway. *J Agromed*. 2016;21(1):24–33.
6. Hounsborne B, Edwards RT, Hounsborne N, Edwards-Jones G. Psychological morbidity of farmers and non-farming population: results from a UK survey. *Commun Ment Health J*. 2012;48:503–10.
7. Letnes JM, Torske MO, Hilt B, Bjørngaard JH, Krokstad S. Symptoms of depression and all-cause mortality in farmers, a cohort study: the HUNT study, Norway. *BMJ Open*. 2016;6(5): e010783.
8. Stain HJ, Kelly B, Lewin TJ, Higginbotham N, Beard JR, Hourihan F. Social networks and mental health among a farming population. *Soc Psychiatry Psychiatr Epidemiol*. 2008;43(10):843–9.
9. Roy P, Tremblay G, Robertson S, Houle J. "Do it all by myself": a salutogenic approach of masculine health practice among farming men coping with stress. *Am J Mens Health*. 2017;11(5):1536–46.
10. Nye C, Winter M, Lobley M. Farmers supporting farmers: Livestock auctions as spaces to reconstruct occupational community and counter mental health issues. *J Agromed*. 2023;28:401–14.
11. Annibal I, Price L, Sellick J. Health and Wellbeing in Farming. In. Lincoln: Lincoln International Business School; 2019.
12. Health Nif, Guideline CE: National clinical practice guideline 90: depression the NICE guidelines on the treatment and management of depression in adults-the updated edition. In. London: The British Psychological Society and The Royal College of Psychiatrists; 2010.
13. Schure MB, Howard M, Bailey SJ, Bryan B, Greist J. Exploring perceptions of a computerized cognitive behavior therapy program in a U.S. Rural Western State. *Rural Mental Health*. 2018;42(3):174–83.
14. Freund J, Buntrock C, Braun L, Thielecke J, Baumeister H, Berking M, Ebert DD, Titzler I. Digital prevention of depression for farmers? A qualitative study on participants' experiences regarding determinants of acceptance and satisfaction with a tailored guided internet intervention program. *Internet Interv*. 2022;29:100566.
15. Nason EE, Blankenship AS, Benevides E, Stump K. The role of social work in confronting the farmer suicide crisis: best practice recommendations and a call to action. *Social Work in Public Health*. 2023;38(1):21–32.
16. House A, Owens D, Storer D. Psycho-social intervention following attempted suicide: is there a case for better services? *Int Rev Psychiatry*. 1992;4(1):15–22.
17. King E, Lamont K, Wendelboe-Nelson C, Williams C, Stark C, van Woerden H, Maxwell M. Engaging the agricultural community in the development of mental health interventions: a qualitative research study. *BMC Psychiatry*. 2023. <https://doi.org/10.1186/s12888-023-04806-9>.
18. Williams C. Living life to the full. Glasgow: Media Innovations; 2008.
19. Living Life to the Full for Farming Communities [<https://lltff.com/home/living-life-to-the-full-series/lltff-for-farming-communities/>]
20. Creswell JW, Clark VLP. Designing and conducting mixed methods research. London: Sage publications; 2017.
21. Smith J, Bekker H, Cheater F. Theoretical versus pragmatic design in qualitative research. *Nurse Res*. 2011;18(2):39–51.

22. Baum F, MacDougall C, Smith D. Participatory action research. *J Epidemiol Commun Health.* 2006;60(10):854–7.
23. Johansson P, Westas M, Andersson G, Alehagen U, Broström A, Jaarsma T, Mourad G, Lundgren J. An internet-based cognitive behavioral therapy program adapted to patients with cardiovascular disease and depression: randomized controlled trial. *JMIR Mental Health.* 2019;6(10): e14648.
24. R Core Team. R: a language and environment for statistical computing. Vienna, Austria: Foundation for Statistical Computing; 2013.
25. Hansson M, Chotai J, Nordström A, Bodlund O. Comparison of two self-rating scales to detect depression: HADS and PHQ-9. *Br J Gen Pract.* 2009;59(566):e283–8.
26. Cameron IM, Crawford JR, Lawton K, Reid IC. Psychometric comparison of PHQ-9 and HADS for measuring depression severity in primary care. *Br J Gen Pract.* 2008;58(546):32–6.
27. Glasgow RE, Rosen GM. Behavioral bibliotherapy: a review of self-help behavior therapy manuals. *Psychol Bull.* 1978;85(1):1–23.
28. Kohlbeck SA, Quinn K, deRoos-Cassini T, Hargarten S, Nelson D, Cassidy L. “I’ve given up”: biopsychosocial factors preceding farmer suicide in Wisconsin. *Am J Orthopsychiatry.* 2023;93(2):131.
29. Kohlbeck S, Schramm A, deRoos-Cassini T, Hargarten S, Quinn K. Farmer suicide in Wisconsin: a qualitative analysis. *J Rural Health.* 2022;38(3):546–53.
30. Thomson AE, Waddell-Henowitch C, Herron R, Epp D, Rauch K, Ryan KD, Mullins S. Rural perspectives on suicide prevention in Canada. *Can J Commun Ment Health.* 2023;42(1):1–14.
31. Grattidge L, Hoang H, Mond J, Lees D, Visentin D, Auckland S. Exploring community-based suicide prevention in the context of rural Australia: a qualitative study. *Int J Environ Res Public Health.* 2023;20(3):2644.
32. Kaur H, Singh A, Singh S: Efficacy of psychological first aid (PFA) by peer support volunteers for suicide prevention in farmers of Punjab. In: *Understanding psychology in the context of relationship, community, workplace and culture.* Springer; 2022. p. 91–105.
33. Gunn KM, Skaczkowski G, Dollman J, Vincent AD, Brumby S, Short CE, Turnbull D. A self-help online intervention is associated with reduced distress and improved mental wellbeing in Australian farmers: the evaluation and key mechanisms of www.lfarmwell.com.au. *J Agromed.* 2022;28:378–92.
34. Thielecke J, Buntrock C, Titzler I, Braun L, Freund J, Berking M, Baumeister H, Ebert DD. Telephone coaching for the prevention of depression in farmers: results from a pragmatic randomized controlled trial. *J Telemed Telecare.* 2022. <https://doi.org/10.1177/1357633X221106027>.
35. Kandlur R, Sardana S, Richardson-Vejlgaard R. The Agrarian distress: factors explaining the will to live among rural and distressed family farmers. *Psychiatry Res Commun.* 2022;2(1): 100019.
36. Balcombe L, De Leo D. The potential impact of adjunct digital tools and technology to help distressed and suicidal men: an integrative review. *Front Psychol.* 2022;12:796371.
37. Dabkowski E, Porter JE, Barbagallo MS, Prokopiv V, Jackson MR. A scoping review of community-based adult suicide prevention initiatives in rural and regional Australia. *Int J Environ Res Public Health.* 2022;19(12):7007.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.