

This is a pre-copyedited, author-produced version of an article accepted for publication in *ICES Journal of Marine Science* following peer review. The version of record A J Brooker, R Skern-Mauritzen, J E Bron, Handling editor: David Fields; Production, mortality, and infectivity of planktonic larval sea lice, *Lepeophtheirus salmonis* (Krøyer, 1837): current knowledge and implications for epidemiological modelling, *ICES Journal of Marine Science*, Volume 75, Issue 4, 1 July 2018, Pages 1214–1234 is available online at: <https://doi.org/10.1093/icesjms/fsy015>.

**Production, mortality and infectivity of planktonic larval sea lice, *Lepeophtheirus salmonis* (Krøyer, 1837): current knowledge and implications for epidemiological modelling**

Brooker, A.J.<sup>1\*</sup>, Skern-Mauritzen, R.<sup>2</sup> & Bron, J.E.<sup>1</sup>

<sup>1</sup>Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling, FK9 4LA

<sup>2</sup>Department of Aquatic Pathogens and Diseases, Institute of Marine Research, P.O. Box 1870, Nordnes 5817, Bergen, Norway

24 \* Corresponding author: a.j.brooker@stir.ac.uk, Tel: +44 (0)1786 467881, Fax: +44 (0)1786  
25 472133

## Abstract

Current sea louse models attempt to estimate louse burdens on wild and cultured salmon by predicting the production and distribution of lice larvae and estimating the risk of transmission. While physical characteristics of water bodies and weather can be accurately modelled, many aspects of sea lice biology require further parameterisation. The aims of this review are (a) to describe current knowledge regarding the production, mortality, and infectivity of planktonic sea lice larvae and (b) to identify gaps in knowledge and suggest research approaches to filling them. Several major gaps are identified, and those likely to have the greatest impact on infection levels are (a) egg production, viability and hatching success, (b) predation in plankton and (c) copepodid infectivity profiles. A key problem identified in current parameter estimates is that they originate from a number of sources and have been determined using a variety of experimental approaches. This is a barrier to the provision of ‘best’ or consensus estimates for use in modelling. Additional and more consistent data collection and experimentation will help to fill these gaps. Furthermore, coordinated international efforts are required to generate a more complete picture of sea louse infections across all regions experiencing problems with sea lice.

Keywords: sea lice, *Lepeophtheirus salmonis*, epidemiology, modelling, Atlantic salmon

## 1 Introduction

The parasitic copepods known as sea lice remain a key constraint to the continued growth of salmonid aquaculture industries worldwide. In the North Atlantic, *Lepeophtheirus salmonis* (Krøyer, 1837) is the primary species infecting cultured Atlantic salmon (*Salmo salar* L.), whereas in the North Pacific, *Lepeophtheirus salmonis oncorhynchii* (Johnson and Albright, 1991a) is prevalent in cultured salmon, although *Caligus elongatus* von Nordmann, 1832 also has some impact. In the southern hemisphere *Caligus rogercresseyi* Boxshall and Bravo 2000 is the principal pathogenic species affecting the Chilean salmon aquaculture industry. For the Norwegian salmon industry, where costs are best characterised, the economic impact of sea lice was estimated to be in excess of 3.4 billion NOK per annum (>£300M) in 2014 for 1,272,358 tonnes production (Iversen *et al.* 2015) with costs estimated to exceed 5 billion NOK (>£390M) in 2015 for 1,303,346 tonnes = 3836.28 NOK tonne<sup>-1</sup> (Iversen pers. comm.). Higher estimates of 7–8 billion NOK per annum (>£540M) have also been presented (Rødseth, 2016). Using FAO statistics for global cultured Atlantic salmon production in 2015 (<http://www.fao.org/fishery/statistics/global-aquaculture-production/en>) for all countries that experience sea lice problems (2,332,290 tonnes) and Iversen's estimate of cost per tonne for 2015 ( $3836.28 \times 2,332,290$ ) provides a rough cost estimate of ~9 billion NOK globally for 2015 (~£700M), with costs likely to have continued to rise since then.

Current integrated pest management (IPM) strategies for sea lice control rely on a small number of licensed pesticides, few of which are effective against all stages of the parasite's life cycle, combined with effective husbandry management tools, such as single-cohort stocking, optimised stocking densities, the use of cleaner fish in polyculture and fallow periods (Skiftesvik *et al.* 2013; Leclercq, *et al.* 2013). Physical techniques to exclude lice, such as the use of barrier nets and snorkel cages, coupled with mechanical tools, including thermal and

turbulent de-licers and laser removal, also constitute an increasing component of current IPM strategies. The adoption of such an increasingly multimodal approach means that the timing of management decisions is critical to the successful control of the parasite. A central element required for the prediction of fluxes in lice populations is an understanding of the production, survival, dispersal, development and infectivity profile of the free-swimming non-infective nauplii and infective copepodid larval stages. However, despite more than three decades of research, knowledge in this area remains extremely poor.

Within the past ten years, several models have been developed that attempt to estimate lice burdens on wild and cultured salmon by predicting the production and distribution of lice larvae from salmon farms and the subsequent risk of transmission. Although complex physical coastal processes can now be reasonably accurately modelled, aspects of larval behaviour and mortality often appear oversimplified. This knowledge gap has serious consequences as it confounds the realistic estimation of the number of lice capable of infecting wild and cultured salmonid populations.

In ecological terms, sea lice can be considered r-strategists, which are characterised by small body sizes, high fecundities and short generation times. Although offspring of r-strategists are dispersed widely, they have a low probability of survival (Cavaleiro & Santos, 2014). However, sea lice differ from many other r-strategists in that they are attached to a host, which provides a permanent food source and allows anomalies, such as a larger body size, and raises the question of whether they have a high fecundity because they experience heavy losses during the larval stages or because they have a nominally unlimited food source. The high fecundity and wide larval dispersal are key aspects of the sea louse's life cycle that determine its overall survival and success. As a result, fecundity and larval biology should be the focus of efforts to predict lice burdens on fish. In the life cycle of the sea louse, however, the free-

swimming stages are essentially a ‘black box’ that cannot be easily observed directly from field studies. Once a copepodid has attached to a host, development is more predictable as development after infection is unaffected by copepodid age at infection (Tucker *et al.* 2000a; Pedersen, 2009), although at this point host factors such as host species / genotype, immunity and site of infection intervene to affect success. Transmission is still a contentious issue with disagreement over whether lice (despite water currents) are accumulated at their source (*e.g.* Krkošek *et al.* 2005 and implied by Jansen *et al.* 2012) or hydrodynamically spread over large distances (*e.g.* Brooks, 2005; Asplin *et al.*, 2014). Therefore, accurate data are urgently needed to inform and validate increasingly realistic models of larval dispersion and infectivity that combine physical processes with key aspects of lice biology to successfully predict larval dispersion and infection risk.

Early models for predicting lice burdens rely on the relationship between gravid female lice and infective larval stages, based on factors such as fecundity, mortality and moult timings, to predict future cohorts of lice available to infect fish (*e.g.* Heuch & Mo, 2001; Murray, 2002; Tucker *et al.* 2002). Although these models can predict louse numbers within a simple closed system, they cannot be applied to large, open systems, such as fjordic sea lochs where salmon are commonly farmed, as they do not take into account larval dispersion and exogenous sources of mortality, such as predation.

Particle tracking models predict the dispersal over time of particles generated at a point source using hydrodynamic models (*e.g.* Corner *et al.*, 2006), which calculate local current velocities based on local topography, fluid dynamics and external forcing from tidal elevation, freshwater inputs and wind-generated currents. Early attempts to predict the dispersal of sea lice larvae using a particle tracking model were made by Asplin *et al.* 2004 who estimated the dispersal of lice from a salmon farm in Sognefjord, Norway. Detailed currents, hydrography

and wind forcing are calculated using high-resolution, three-dimensional ocean and atmospheric models, and although a temperature-dependant larval growth model is included, there is no estimation of larval mortality or behaviour. It assumes that lice are immortal with passive behaviour, and consequently, the dispersal of lice is overestimated with larvae being spread over a distance of 100km in just a few days (Asplin *et al.* 2004).

In order to accurately estimate infection risk, it is clear that certain aspects of louse biology, such as survival, mortality and development times, need to be incorporated into these types of models, and more recent models have attempted to do this. Murray and Amundrud (2007) and Amundrud and Murray (2009) present a coupled biophysical and particle tracking model of Loch Torridon, Scotland that incorporates development times as a function of temperature and a fixed mortality rate based on laboratory observations.

More recent models have become increasingly complex, and Asplin *et al.* (2011 and 2014) present a model of a Norwegian fjord comprising a number of sub-models: a coastal ocean model, an atmospheric model, a fjord model, and a salmon louse growth and advection model. While the salmon louse sub-model includes relevant parameters regarding stage timings, it only includes a few simple behavioural parameters, *i.e.* a diel vertical migration, limited to depths above 10 m and avoidance of salinities below 20 ‰; however, it does not calculate louse mortality. A further model by Stucchi *et al.* (2011), which models the hydrographically complex Broughton Archipelago in British Columbia, Canada, includes a comprehensive sub-model of egg production, larval development, mortality and behaviour using data from the literature, including the effects of temperature and salinity on these parameters. In addition, a recent model similar to the one utilised by Asplin *et al.* (2014), which uses a mortality rate of 17 %, predicts that larval behaviour potentially has significant effects on advection (Johnsen *et al.* 2014).

Aldrin *et al.* (2013) and Kristoffersen *et al.* (2014) present a model based on a statistical network of Norwegian salmon farms. Monthly external and internal infection pressure and the risk of infection between neighbouring farms are predicted based on lice burden estimates from the previous month and the distances between neighbouring farms. The model is fitted to actual lice counts from Norwegian farms between 2003 and 2011. It uses temperature-dependent fecundity and larval demographics, although mortality rates for free-swimming larvae and chalimus stages are fixed.

While these models have made significant progress in predicting larval dispersal in semi-enclosed water bodies, model validation with field data is difficult, and there are always discrepancies between the model output and field observations. For example, Salama *et al.* (2011) and Adams *et al.* (2012) found very few larval sea lice in plankton tows in areas where models had predicted high numbers. However, correlation between predicted and observed infections appear to be more accurate for the model developed by Sandvik and colleagues (Sandvik *et al.*, 2014). Model variables are based on the best available data, and while accurate topography and hydrography data can easily be obtained, detailed information regarding the life history of sea lice is often lacking, despite over three decades of research in this area. Where models incorporate larval mortality, for instance, they use a constant mortality at each larval stage, which may be kept constant (*e.g.* Aldrin *et al.*, 2013; Johnsen *et al.*, 2014) or vary according to salinity (*e.g.* Amundrud and Murray, 2009; Adams, 2012). In reality, however, larval mortality is extremely variable according to temperature, salinity, season, moult stage and predation in the plankton, *etc.* While some data are available regarding these different parameters, others are distinctly lacking, and more research is required in these areas. Acquiring experimental data on these variables will allow the more realistic parameterisation of key elements relating to abundance and infectivity of free-swimming larval sea louse stages



for incorporation into models that may more accurately predict the risk of infection under various environmental conditions.

Some models are now considered sufficiently developed to warrant their use as components of an integrated sea louse management strategy. For example, Norwegian salmon farming will from 2017 be regulated regionally through an operational management system comprising the application of predictive models that predict louse infection intensities along the entire coastline (Asplin, 2014), combined with a process of continuous model validation and calibration against real-world data (Bjørn *et al.* 2014, Sandvik *et al.*, 2016).

The aims of this review and analysis were as follows:

1. To analyse the available literature to determine current knowledge regarding the recruitment and survival of free-swimming nauplii and copepodid larvae and factors that affect the longevity and infectivity of copepodids. Where no specific data regarding sea lice were available, the wider literature was consulted, *e.g.* predator and prey selection in plankton, to inform questions regarding the fate of sea lice larvae in the ocean.
2. To assess the remaining knowledge gaps that might be filled by experimental or field sampling studies.

Additional considerations:

- While this review focuses primarily on *Lepeophtheirus salmonis* spp., observations from other species that are problematic in salmonid aquaculture are also noted where appropriate.
- This review also focuses principally on knowledge concerning louse larvae deriving from farmed fish due to both their greater accessibility and the fact that environmental

parameters can only be sufficiently controlled or consistently measured in defined water masses.

- Hitherto, there has been some conflation of data arising from Atlantic and Pacific sea louse studies. Evidence for clear genomic and phenotypic differences between these subspecies has made it evident that the origin of data regarding these subspecies should be considered when interpreting the results.

## **2 Larval recruitment and survival**

In order to accurately predict when and how many infective copepodids are available for infection, it is necessary to quantify the rate of larval production, which is based on female fecundity, and the subsequent development and survival rates of the larvae. These are influenced by a range of biotic and abiotic factors that fluctuate seasonally and can have an impact on adult lice during mating and egg production, on eggs during development and upon larvae once they have hatched.

### **2.1 Fecundity**

The fecundity of sea lice varies considerably, and early observations showed that a single egg string can contain <100–700 eggs (Wootten *et al.* 1982). Many studies have shown that exogenous factors, such as temperature, photoperiod, salinity and food availability, interact with endogenous factors to determine fecundity in crustaceans (*e.g.* Koop & Field, 1980; Williams, 1985; Johnston & Dykeman, 1987, Maranhão & Marques, 2003). Similarly, variations in the levels of sea lice infection between seasons and under different environmental conditions suggest alterations in reproductive output in response to fluctuating environmental parameters (Ritchie *et al.* 1993).

It is clear that temperature has a strong influence on fecundity (Tully, 1989), and the number of eggs per string is positively correlated with female body size (Tully & Whelan, 1993). Heuch *et al.* (2000) found that adult female lice of wild origin in Norway were significantly larger than adult female lice of farm origin. Despite seasonal variations, lice of wild origin in Ireland were similarly found to be significantly larger and carried approximately twice as many eggs as lice of farm origin (Tully & Whelan, 1993). A similar pattern was reported by Pike and Wadsworth (1999), who noted that female lice of wild origin produced  $965 \pm 30.1$  eggs per egg string pair compared to  $758 \pm 39.4$  and  $297 \pm 19.1$  for lice originating from untreated and treated farmed salmon, respectively, on the West Coast of Ireland. At 7.2 °C, females were observed to produce a new pair of egg strings on average 11 days after the first pair were removed, while at 12.2 °C this period was reduced to 5 days, and this continued for the reproductive life of the female, with an average of 4.95 pairs of egg strings per female under experimental conditions (Heuch *et al.* 2000). In this experiment, the first pair of egg strings was always significantly shorter with the mean number of eggs increasing from 152 eggs per string to 285 eggs per string for subsequent egg strings, whereas Johnson and Albright (1991b) recorded a mean number of eggs per string of  $344.6 \pm 79.8$  in lice cultured at 10 °C and 30 ‰ originating from wild and farmed chinook salmon (*Oncorhynchus tshawytscha*) and farmed Atlantic salmon. Similarly, Gravid (1996) recorded a mean of  $141.09 \pm 22.19$  eggs per string for the first pair of egg strings,  $216.4 \pm 67.59$  eggs per string for the second pair of egg strings and  $208.2 \pm 50.97$  eggs per string for the third pair of egg strings. It appears that there may be a difference in the batch size in Atlantic *L. salmonis salmonis* (Heuch *et al.* 2000 and Gravid 1996) and the Pacific *L. salmonis oncorhynchi* (Johnson and Albright 1991a), which highlights the importance of discriminating between the two subspecies (Skern-Mauritzen *et al.*, 2014). Fecundity was found to be lower in *C. elongatus* with the number of

eggs per string being  $52.62 \pm 17.08$  in *C. elongatus* compared to  $206.2 \pm 74.09$  in *L. salmonis* at 14 °C (Gravil, 1996). Key values for fecundity are shown in Table 1.

Ritchie *et al.* (1993) and Gravil (1996) investigated the reproductive output of *L. salmonis* from salmon farms on the West Coast of Scotland and found that the number of eggs per string was negatively correlated with temperature, with significantly more eggs being produced in winter and early spring than in summer and autumn (Figure 1). In Ritchie *et al.* (1993), the mean number of eggs per string increased significantly from 147 to 246 between October and March (temperature range 12–5 °C) before decreasing to 175 eggs per string in August (13 °C). A similar pattern was seen by Gravil (1996), who found that the number of eggs per string ranged from  $194.1 \pm 66.8$  in October to  $286.9 \pm 64$  in March. There appears to be a period of lag of egg string length in response to temperature as the lowest temperature was recorded in February whereas the longest egg strings were found in March, and this lag may reflect the time required for egg strings to develop before being extruded at low temperatures. Samsing *et al.* (2016) found a similar trend in lice acclimatised in the laboratory at different temperatures with the number of eggs per string increasing from  $\sim 135 \pm 5$  at 20 °C to  $\sim 295 \pm 10$  at 5 °C. In the same experiment, it was found that the number of eggs per string produced at 3 °C was lower ( $\sim 153 \pm 10$ ) than at the higher temperatures tested. This decrease corresponded to a decreased body size and coincided with a failure in larval development, and it was speculated that this temperature could be close to the limit of their biological tolerance, at least for the tested lice (Samsing *et al.*, 2016).

A variety of other factors may also affect lice fecundity. As an example, host condition and the use of chemotherapeutants have been proposed as possible influences on egg string length and the viability of larvae (Tully & Whelan, 1993). Likewise, fecundity may vary with host species, either as a result of diet, the physiological status of the fish or genetic variation

(Johnson & Albright, 1992; MacKinnon *et al.* 1995; Mackinnon, 1998). This follows from the intimate metabolic associations between hosts and parasites, which are often reflected in the evolution of their genomes (e.g. Zarowiecki & Berrimann, 2015). It has also been suggested that host immune responses may modify lice fecundity. For instance, Grayson *et al.* (1995) found that gravid female lice on Atlantic salmon injected with extracts derived from adult *L. salmonis* had a significantly lower fecundity than control fish. Similarly, Nilsen (2016) has presented work suggesting that use of a recombinant vaccine to the salmon louse Ls4D8 protein, a homologue to subolesin in ticks and my32 in *C. rogercresseyi*, gave rise to reduction in egg strings. Host-related and abiotic conditions may not be the only factors governing salmon louse fecundity. As an example, intraspecific competition between lice on a given host is suggested to result in reduced fecundity with increasing salmon louse infection densities (Ugelvik *et al.*, 2017). Louse fecundity is clearly the product of a number of biotic and abiotic factors, most of which remain to be fully characterised.

## **2.2 Hatching**

Egg strings with non-viable eggs are sometimes extruded, and Heuch *et al.* (2000) found that this happened most frequently in the second and third batches of egg strings. Gravid (1996) reported that 2.1 % of egg strings consisted entirely of non-viable eggs. According to Heuch *et al.* (2000), the number of viable eggs per string varied according to temperature, with a median of 13.3 % of eggs being non-viable at 7.2 °C and 7.5 % being non-viable at 12.2 °C. Similarly, Samsing *et al.* (2016) found that hatching success was strongly influenced by water temperature, with 100 % success at 20 °C and 15 °C decreasing to  $28 \pm 4$  % success at 3 °C. Conversely, Gravid (1996) found no correlation between egg viability and temperature in *L. salmonis* on farmed salmon on the West Coast of Scotland with a mean of  $17.66 \pm 23.01$  %

non-viable eggs over one year. In comparison, the mean number of non-viable eggs per string in *C. elongatus* was  $28.19 \pm 24.81$  %, with 18.33 % of egg strings entirely consisting of non-viable eggs (ibid.).

Salinity has a considerable effect on hatching, and egg strings maintained at 10 °C and 10 ‰ salinity failed to develop in Johnson and Albright's (1991b) experiments. At salinities of 15 ‰ and 20 ‰, hatching success was 70 % and 78 %, respectively, but only at 20 ‰ were any active nauplii produced (19.8 %). At salinities of 25 ‰ and above, hatching success was 100 %, but at 25 ‰ only 51.1 % of nauplii were active, whereas at 30 ‰ this figure was 65.9 %. Gravid (1996) reports a similar pattern with hatching success ranging from 3.27 % in freshwater to 86.36 % at 30 ‰ salinity. The effect of photoperiod was investigated by Gravid (1996), but it had no effect on hatching period or success. Key values for hatching are shown in Table 2.

The hatching period is variable, and Johnson and Albright (1991b) report that it ranged from 18 to 65 h, with a mean of  $31.7 \pm 13$  h for egg strings incubated at 10 °C and 30 ‰ salinity. The authors of the current review consider these to be at the extreme end of hatching periods observed based on personal observations, although this may represent a difference between Atlantic and Pacific *L. salmonis*.

### **2.3 Stage timings**

Development times are highly dependent on temperature and have been addressed in various studies summarised in Table 3. Overall, the egg development time varies between 1.8–45.1 days for temperatures ranging between 2–20 °C (Johnson & Albright, 1991b, Boxaspen & Næss, 2000, Samsing *et al.*, 2016). The duration of the first nauplius stage varies between 9.2–52 h at temperatures ranging between 5–15 °C, while the corresponding duration for the

second nauplius stage varies between 33–170.3 h for temperatures ranging between 5–19 °C (Johannessen, 1977, Wootten *et al.*, 1982, Johnson & Albright, 1991b, Gravid, 1996). Durations of the stages seem to be comparable for Pacific and Atlantic lice, and reported ranges agree with the ranges found in publications where developmental times were reported for both naupliar stages combined (Gravid, 1996, Boxaspen & Næss, 2000, Samsing *et al.*, 2016). While temperature has a considerable effect on egg production and larval development, photoperiod does not appear to have any significant effect (Ritchie *et al.* 1993; Gravid, 1996). The time required for physically moulting (exuviation) from nauplius I to nauplius II and nauplius II to copepodid are reported as  $10.53 \pm 4.34$  mins and  $12.21 \pm 3.87$  mins, respectively, and during the moult the larvae are inactive and sink through the water column (Gravid, 1996). It appears that the temperature of acclimation of adult female lice is important in determining the temperature tolerance of their eggs and larvae. Johannessen (1975) reports that in adult lice cultured at 9 °C, nauplius development occurred only between 8–11 °C, whereas acclimation at 11.5 °C allowed larval development up to 22 °C. In adult lice maintained at 3 °C, however, nauplii failed to develop to copepodids (Samsing *et al.*, 2016).

## **2.4 Survival**

Nauplii that hatch successfully are planktonic. At this stage they do not feed, but are lecithotrophic (yolk feeding) and rely on their energy reserves until they moult to infective copepodids and find a suitable host (Johnson & Albright, 1991b). The survival of sea lice and the rate at which they deplete their energy reserves are strongly influenced by temperature and salinity. The size of larvae and their lipid stores is also dependant on season, and Gravid (1996) reports that nauplius I larvae were largest in August with a mean body width of 214.05 µm and a mean lipid reserve width of 135.84 µm compared to 197.76 µm and 112.98 µm in May

for mean body width and mean lipid reserve width, respectively. It is likely that increased energy reserves will increase the longevity or compensate for a higher temperature-dependent metabolism of the non-feeding larval stages, although no data are available comparing survival at different times of year.

Johnson & Albright (1991b) report that active copepodids were only obtained at salinities above 30 ‰ at 10 °C (35.2 % active), although survival was extremely variable ranging from 0–80.6 % per egg string. Similarly, Gravid (1996) found that copepodids were only obtained at salinities greater than 25 ‰, and at 10 °C and 35 ‰, 18.33 % reached the infective copepodid stage with nearly 50 % mortality being seen in the nauplius I stage. Samsing *et al.* (2016) found that sea lice larvae from Scotland did not proceed past the nauplius II stage at 5 °C and 3 °C, respectively, but died before moulting to copepodids, and at 7.5 °C, very few copepodids were obtained (Gravid, 1996). In sea lice adapted to low temperatures, however, copepodids were obtained from 25 % of egg strings reared at 2 °C, 42 % at 3 °C, 100 % at 4 °C and 75 % at 5 °C (Boxaspen and Næss, 2000). In *C. elongatus*, Pike *et al.* (1993) report 90 % survival from the nauplius stage to the copepodid stage at 15 °C with this figure decreasing to 60 % at 5 °C.

As with all copepods, sea lice have preferred environmental conditions, which are determined by their physiological tolerances. Copepodids that were transferred from full-strength seawater to 5 ‰ salinity survived for just 3 h at 10 °C, and those transferred to 10 ‰ salinity survived for less than one day (Johnson & Albright, 1991b). A similar experiment by Gravid (1996) found that the median survival time was 14.87 h at 0–10 ‰. While copepodids can osmoregulate above 16 ‰, their haemolymph becomes rapidly diluted below 12 ‰, and they are unable to regulate cell volume and die within a few hours (Hahnenkamp & Fyhn, 1985; Pike & Wadsworth, 1999).



Once nauplii moult to copepodids, they need to find a suitable host before their lipid reserves are depleted, and the rate at which this occurs is also influenced by temperature and salinity. Hyperosmotic regulation is energetically costly, and an increased energy demand significantly reduces the survival time of copepodids due to their limited energy reserves (Torres *et al.* 2002). Johnson and Albright (1991b) report that survival was prolonged at salinities of 15–30 ‰ and temperatures of 5–15 °C, and that mean survival times were between two and eight days. Similarly, Wootten *et al.* (1982) report that the mean survival time of copepodids at 12 °C was 4 days at an unspecified salinity. In Gravid (1996), the median survival time of copepodids was 54 h at 15 ‰, 67 h at 20 ‰, 68 h at 25 ‰, 55 h at 30 ‰ and 64 h at 35 ‰, which reflects the increased energy required for hyperosmotic regulation at lower salinities. Conversely, Bricknell *et al.* (2006) report the median survival time of *L. salmonis* copepodids to be 4 h at 16 ‰, 6 h at 19 ‰, 8 h at 23 ‰, 11 h at 26 ‰, 24 h at 29 ‰, 22 h at 33 ‰ and 25 h at 36 ‰. The reason for the differences in survival times reported in Gravid (1996) and Bricknell *et al.* (2006) is unknown, although Bricknell *et al.* used copepodids that were a few days old and cultured them with aeration whereas Gravid used unaerated containers.

According to Johnson and Albright (1991b), the maximum survival time was 17 days at 10 °C and 25 ‰ salinity, and copepodids in lower salinities (15–20 ‰) were generally less active than those maintained at higher salinities (25–30 ‰). In full strength seawater (35 ‰), the maximum survival time of copepodids at 10 °C was 18 days (Gravid, 1996). Due to the reduced hatching success and subsequent low survival of *L. salmonis* in low salinities, it is likely that they may be excluded from salinities less than 15 ‰ (Johnson & Albright, 1991b), and survival is severely compromised at salinities below 29 ‰ (Tucker *et al.* 2000b). Although survival is reduced at lower salinities, short-term exposure to reduced salinities does not have a long-term impact on the development of surviving copepodids (Bricknell *et al.* 2006). Attachment to a

host was not observed to improve survival at reduced salinities (Hahnenkamp & Fyhn, 1985) and these authors suggest that, unlike adult lice, the copepodid and chalimus stages are unable to use ions obtained from their host to replace those lost to a hypo-osmotic environment. However, it appears likely that due to their small size, attached larvae will receive at least some protection from reduced salinities through boundary layer effects coupled with close contact with the host/host mucus, and it is also clear that as these are feeding stages, some protection would be received from ingested host tissue.

The survival time of copepodids is inversely related to temperature, and Samsing *et al.* (2016) report that the survival time of 80 % of copepodids was 12.5 days at 7 °C, 13 days at 10 °C, 9.5 days at 15 °C and 6 days at 20 °C; at 5 °C it was reduced to 10 days. This pattern is presumably due to lower metabolism and, therefore, increased longevity of energy reserves at lower temperatures, although at very low temperatures there appear to be other factors limiting survival. Median survival times reported by Gravid (1996) were 116 h at 5 °C, 90 h at 10 °C and 82 h at 15 °C at full salinity (35 ‰), although these appear to be gross underestimations and may be due to sub-optimal culture conditions. There is, however, a seasonal investment by adult females in reproduction as nauplii are larger and have larger energy stores in summer than in winter (Gravid, 1996). At higher temperatures, metabolism is higher and larvae are more active, so their energy stores are more rapidly depleted (*ibid.*). It is possible that the increase in the size of larvae and their energy stores in summer may be a compensatory mechanism to account for their energy stores being depleted more rapidly than in winter, which ensures that their life expectancy is similar to that at colder winter temperatures. Further experimental work is required to confirm this. Key values for survival are shown in Table 4.

### **3 Behaviour**

While both of the free-swimming larval stages are planktonic, the nauplius stages of sea lice are principally dispersal stages, whereas the copepodid stage must locate, re-establish contact with and subsequently infect a suitable host. The larvae are subject to currents, which serve to disperse them over a wide area, and although the larvae have limited movement capabilities, their dispersal can be partially influenced by certain behaviours, *e.g.* aggregating at particular depths in the water column (Johnsen *et al.*, 2014). In order to maximise their chances of survival and host interception, they must be able to respond to cues present in their environment and react to them appropriately. Their behavioural responses can be categorised according to the following activities (Bron *et al.*, 1993):

1. Predator avoidance
2. Avoidance of adverse environmental conditions
3. Movement into or maintenance within host-rich environments
4. Host location
5. Host contact/settlement
6. Confirmation of host suitability

Cues that may play a role in influencing the behaviour of sea lice larvae include light, chemical, pressure, temperature and water flow/vibration.

### ***3.1 Swimming speeds / activity***

Both nauplius and copepodid stages have been observed to actively swim upwards as they are negatively buoyant, and their movements are punctuated by periods of passive sinking (Bron, 1993, Gravid, 1996). Haury and Weihs (1976) suggest that this behaviour theoretically saves energy compared to continuous swimming at a fixed depth, which is particularly important for the lecithotrophic larvae of *L. salmonis*, which must conserve their limited energy reserves wherever possible. Despite their energy considerations, copepodids must maintain their

position in the water column where their chances of encountering hosts are highest (Bron, 1993). However, Gravid (1996) found the activity of nauplii and copepodids to be dependent on temperature; at 5 °C their movements were reduced and they aggregated at the bottom of containers, whereas at 10 °C and 15 °C they spent more time actively swimming than passively sinking and aggregated at the surface. However, these results may be affected by insufficient acclimation.

Copepodids swim more rapidly than nauplii and have longer swimming periods and shorter rest periods (Bron, 1993). Gravid (1996) reports that the mean swimming speed of nauplii was  $1.25 \pm 0.16 \text{ cm s}^{-1}$ , whereas the mean swimming speed of copepodids was  $2.14 \pm 0.24 \text{ cm s}^{-1}$ . The mean sinking speeds were  $0.09 \pm 0.01 \text{ cm s}^{-1}$  and  $0.10 \pm 0.03 \text{ cm s}^{-1}$  for nauplii and copepodids, respectively. In this study, the maximum speed recorded was  $10.23 \text{ cm s}^{-1}$  when stimulated by vibration of the test chamber and gives an indication of the swimming ability of copepodids. A similar one-second burst speed of  $9 \text{ cm s}^{-1}$  was recorded by Heuch and Karlsen (1997), although speeds of  $2 \text{ cm s}^{-1}$  were sustained when stimulated. In comparison, reported salmon swimming speeds are two orders of magnitude higher (Colavecchia *et al.* 1998). Thus, while chemotaxis may be important in positioning the larvae in suitable water masses, the pursuit of a salmon host, as opposed to the interception of it at close range, is not a viable strategy.

Current speed and host swimming speed affect the ability of infecting copepodids to make initial contact with the host and to remain attached following contact. Given the respective speeds of copepodids and salmonids, the former cannot pursue the host but must intercept it by burst-swimming when detecting it in the water column. The exposure time of the copepodid to the host reduces with increasing current/host swimming speed, which in turn reduces the window of opportunity for infection. In addition, the low-flow zone (boundary layer) caused

by drag at the surface of the fish, becomes thinner with increasing current/host speed, which increases the exposure of the copepodid to the ambient water flow during attachment. This means that at higher flows, the copepodid has less opportunity to make contact and is more likely to be removed from the host by the current (Bron, 1993). The greater boundary layer thickness and, hence, shelter from the ambient current offered by fin rays held perpendicular to the direction of water flow is considered to provide some explanation of the observed greater frequency of copepodid settlement on the fins of hosts (Bron, 1993, Bron *et al.*, 1993). Similarly, the slower swimming speed of fish in tank challenges may explain the largely artefactual attachment of copepodids to the gills in such trials, an observation rarely made under field conditions (*ibid.*). While larger fish swim faster, this is offset by the provision of a larger surface area for settlement and a greater boundary layer/shelter provided by larger fins. Frenzl (2014) observed declining numbers of attaching copepodids with increasing current speeds. Following a dose of 2,500 copepodids fish<sup>-1</sup> introduced in a flume challenge, highest infection occurred at 0 cm s<sup>-1</sup> (mean 8.4 copepodids per fish) and lowest at 32.6 cm s<sup>-1</sup> (mean 0.2 copepodids per fish).

Little is known concerning the effects of competition for space and/or resources during initial copepodid settlement. However, Frenzl (2014) has demonstrated a non-linear increase of infection numbers with challenge dose in flume challenges, possibly suggesting the increasing saturation of available settlement niches with increasing numbers available for infection.

### **3.2 Light**

Copepodids of *L. salmonis* are highly photopositive and move towards the illuminated zone of the vessel in laboratory experiments even at low light intensities (Johannessen, 1975; Wootten *et al.* 1982; Bron *et al.* 1993, Gravid, 1996). The nauplius stages are also

photopositive, but the nauplius I stage only exhibits a positive response at light intensities of 200 lux or more, whereas this value is 85 lux in the nauplius II (Gravil, 1996). Whereas nauplii exhibit increasing activity with increasing light intensity, copepodids do not (ibid.). The free-swimming larval stages of *C. elongatus* are also phototactic, with the copepodids showing a contrasting greater response to light than the nauplii stages (Hogans & Trudeau, 1989). In *L. salmonis*, a peak response was seen at a wavelength 500 nm in the nauplius II stage (Gravil, 1996) and 550 nm in the copepodid stage (Bron *et al.*, 1993, Gravil, 1996), and this corresponds to the maximum transmitted light intensity at twilight, which may be a cue for vertical migration in copepodids as suggested for free-living copepods (Forward & Douglass, 1986). In flume challenges, Frenzl (2014) found maximum sensitivity of copepodids to light at 455 nm. In addition to the response to constant light, evidence for a response to changing light intensities/shadows (scototaxis) in adult sea lice (authors' qualitative observations) and copepodids (Fields *et al.* 2017) strongly indicates a behavioural response towards moving objects obstructing or reflecting light.

Heuch *et al.* (1995) found a strong diel vertical migration in *L. salmonis* copepodids where they gathered near the surface during the day and spread out into deeper layers at night. Despite the recognised photopositive behaviour of copepodid stages, a number of authors observed successful settlement or attempted settlement in darkness (Johnson & Albright, 1991b; Bron *et al.*, 1993, Heuch *et al.*, 2007, Frenzl, 2014), although settlement success was generally lower than when under illumination. As salmon remain in deeper waters during the day and rise to the surface at night, they swim through a population of sinking or rising copepodids every 12 h (Heuch *et al.*, 1995). In addition, vertically migrating hosts produce stronger currents than resting fish, and pressure waves in front of swimming fish trigger a looping behaviour allowing nearby copepodids to avoid predation and attach to a host (Bron *et al.*, 1993; Heuch & Karlsen,

1997; Heuch *et al.*, 2007). Bron *et al.* (1993) and Gravid (1996) also demonstrated that copepodids are negatively geotactic, *i.e.* they swim towards the surface, which also suggests that they tend to aggregate in surface waters. Presumably, these experiments were conducted with illumination, and therefore, it is not known whether copepodids would be negatively geotactic in the dark when they would normally spread out into deeper water. In the study by Heuch *et al.* (1995), 6 m-deep mesocosm bags were suspended in the water column, and therefore, the vertical migrations of copepodids were limited by the depth of the bags. Zooplankton appear to scale their vertical migrations according to the available water depth (Young & Watt, 1993), so the relationship of experiments with constrained depths to the natural situation is uncertain. This has implications for the dispersal of lice by water currents as current velocity and direction often vary with depth. It is clear, however, that wind forcing can be a dominant component of sea lice dispersal (Murray & Amundrud, 2007; Amundrud & Murray, 2009), and therefore, improved knowledge of the diel vertical migration of copepodids between surface and deeper waters would allow the wind forcing component of sea louse dispersal to be predicted more accurately.

### 3.3 Salinity

In salinities less than 21 ‰, the swimming ability of nauplii and copepodids is lost, although full activity is recovered if the exposure time is short (< 5 minutes) (Gravid, 1996). Bricknell *et al.* (2006) found that copepodids actively avoided salinities lower than 27 ‰ by orientating themselves in a vertical sinking position and occasionally actively swimming downwards. Given a choice, they will remain in full strength seawater. Energy is expended for osmoregulation and to maintain their position in the water column, as sinking rates increase with decreasing salinity due to water density changes (Bricknell *et al.*, 2006). It is likely that copepodids avoid areas of low salinity as they require increased energy expenditure, which

reduces survival time (Torres *et al.*, 2002). As low salinities reduce the activity levels of copepodids, their ability to respond to host cues is reduced (Bricknell *et al.*, 2006).

### **3.4 Currents**

It has been proposed, although supporting evidence is lacking, that copepodids may actively migrate to river mouths where high concentrations of salmon smolts are present at certain times of year, which would increase their probability of encountering a host (Carr & Whoriskey, 2004; Costelloe *et al.*, 2004; McKibben & Hay, 2004). Studies in estuarine areas in Ireland suggest that copepodids are not found near the mouths of rivers for the majority of the year (Costelloe *et al.*, 1998a), but high concentrations coincide with the seaward migration of salmon smolts and the freshwater migration of adult salmon (Costelloe *et al.*, 1998a; McKibben & Hay, 2004). As copepodids are capable of actively altering their position in the water column, it is possible that they may be able to use vertical positioning to compensate for lack of long distance swimming capabilities, using tidal currents to migrate towards river mouths, although no evidence has been found to support this. As copepodids have been shown to remain active in the water column (Bron *et al.*, 1993; Heuch *et al.*, 1995; Gravid, 1996), they are distributed within a water body according to the prevailing currents and are, thus, unlikely to directly influence their large-scale movement towards a particular location. It has been suggested that at some times of year, a high concentration of copepodids near river mouths could result from hatching of egg strings from lice on adult salmon, which often congregate at river mouths prior to their migration upstream, particularly during periods of low river flow (Jonsson *et al.*, 1990; Smith *et al.*, 1994). Similarly, the absence of copepodids at river mouths during periods of high rainfall might simply be due to salmon migrating rapidly upstream when river flow is high (Costelloe *et al.*, 1998a,b).

### **3.5 Host location**



The responses of sea lice copepodids to physical cues, such as light and salinity, enable them to gather in areas where host fish are likely to be found, and mechanical cues enable them to infect a host. Chemoreception also plays an important role in host location, with copepodids employing the cues provided by kairomones, specific chemicals released by host fish, to improve the probability of host encounter. Copepodids swim with a general search pattern, but once a host odour has been detected, a host-encounter search pattern is switched on, which consists of increased duration and frequency of turning during the normal sinking and swimming behaviour (Genna, 2002). A directional component is also apparent whereby activated copepodids swim towards a suitable odour source over a distance of centimetres (Bailey *et al.*, 2006), although a group of salmon might initiate a response over a scale of metres (Mordue Luntz & Birkett, 2009). Experiments have shown that *L. salmonis* copepodids are attracted to odours from salmon and sea trout, and behavioural activation and positive upstream chemotaxis occur in the presence of salmon-derived compounds (Devine *et al.*, 2000; Genna, 2002; Ingvarsdottir *et al.*, 2002; Bailey *et al.*, 2006). While both light and chemoreception elicit behavioural responses in the infective copepodids, it has been shown that the effect of light on the swimming response is stronger than that of responses elicited by olfactory cues and that the two sources of sensory cues may act in combination to give stronger and more persistent responses (Fields *et al.* 2017). Non-host odours activate copepodids, but positive chemotactic movements are not observed, indicating that *L. salmonis* can discriminate between salmonid hosts and other non-host fish from their odour (Bailey *et al.*, 2006). In comparison, *C. elongatus*, which is a generalist and infects many different species of fish, demonstrates behavioural changes to chemical cues from a wide range of fish, although physical cues may be more dominant in this species (Mordue Luntz & Birkett, 2009). Although the activity of copepodids appears to be affected by temperature, with reduced activity at lower temperatures (Tucker *et al.* 2000b), it is not known whether low temperatures

affect the switch to host-seeking behaviour and the distance over which they may be able to detect host cues.

Despite their avoidance of areas of low salinity, the use of haloclines by copepodids has been proposed as a host-finding mechanism, since host odours may accumulate in thin layers where a density gradient occurs. In this respect, 80 % of copepodids were observed to aggregate at the confluence of a 15–30 ‰ step-salinity gradient in laboratory experiments (Heuch, 1995). In addition, positioning close to a halocline may increase the chance of encountering a host, as salmon have been observed to follow salinity gradients (Lyse *et al.*, 1998; Finstad *et al.*, 2000)

#### **4 Infectivity**

While some previous models of sea louse dispersion include a mortality factor, they do not account for variations in infectivity, *i.e.* the ability of a louse encountering a fish to infect it. Infection can be considered in terms of a two-phase process comprising a reversible attachment phase following contact and an irreversible settled phase during which the copepodid becomes physiologically committed and can no longer re-enter a free-swimming state. In the salmon louse, the former phase comprises initial copepodid attachment using the antennae (Bron *et al.*, 1991) followed by manoeuvres to embed the anterior of the cephalothorax. At some point following initial attachment, the copepodid commences feeding and starts the process of metamorphosis and moulting to the chalimus I stage. Although the precise triggers and point of irreversible commitment remain to be identified, antimicrobial peptides (AMPs) have been shown to affect *C. rogercresseyi* frontal filament development *in vitro* (Núñez-Acuña *et al.* 2016). It is, therefore, incorrect to assume that, once the copepodid stage is reached, 100 % infection will occur (Gravil, 1996). Dispersion on currents and host location behaviour bring the copepodids into the same locality as potential hosts, but the

process of infection is influenced by various factors, including salinity, light, temperature, season, a range of host factors and copepodid age. A further difficulty encountered in the literature is the somewhat nebulous concept of ‘infection success’. For some authors, copepodids attaching to the fish are counted directly. However, given the reversible nature of initial attachment and difficulty of capturing fish without dislodging attached copepodids, such counts may be prove less accurate, although they provide an estimate of successful contact and attachment. As an alternative, many authors only count infection success following the moult to chalimus I, at which point larvae are hard to dislodge due to the permanent frontal filament attachment. This latter approach, however, incorporates a far greater potential for the superposition of host immunity / site selection effects upon the successful completion of the copepodid instar.

#### ***4.1 Age at infection***

As lecithotrophic larval stages are reliant on their energy reserves for swimming, moulting and host infection, the excessive depletion of these reserves prior to infection can result in the loss of infective capability. As copepodids age, a higher proportion display reduced activity due to the depletion of energy reserves or senescence (Bron, 1993). Gravid (1996) found that the mean size of lipid vesicles in the mid-gut of copepodids was significantly reduced after seven days, and Tucker *et al.* (2000a) reports a significant reduction in the calorific value of *L. salmonis* larvae over seven days with a sharp decline after five days. By measuring stored lipid volume, it is possible to determine age and viability in individual copepodids, and these can be divided into three loose categories: early copepodids with an apparent increase in lipid volume reflecting incorporation of naupliar lipids into distinct vesicles in the gut; mid-life copepodids, which show a downward trend in lipid levels and may be the most active individuals with mature infective capabilities; and late copepodids with low reserves of lipid,

which may be less capable of infection (Cook *et al.*, 2010). The depletion of energy reserves, which consist primarily of lipids, might also result in a loss of buoyancy, making swimming more energetically costly (Bron, 1993), although Gravid (1996) found no evidence to support this. Gravid (1996) observed three stages of activity: newly moulted copepodids swam in spontaneous bursts without stimulation; at eight days at 10 °C, 50 % of copepodids were only active when stimulated; after eight days, remaining copepodids only showed activity after being stimulated by a water jet from a pipette. This suggests that copepodids may adopt a strategy of energy conservation if a host is not located after a certain period of time, and that by only becoming active when stimulated, they preserve their remaining energy stores as long as possible.

This reduced activity level affects infectivity, and Gravid (1996) reports that copepodid infection success at 10 °C and 35 ‰ salinity was  $22.22 \pm 8.32$  % at one day old and  $14 \pm 8.71$  % at seven days old. At seven days old, approximately 20 % of copepodids were active without stimulation and 40 % were active with or without stimulation. Bron (1993) reports similar infection rates with 23.2 % settlement under illuminated conditions and 18.4 % settlement in the dark for 1–3-day-old copepodids, although there was no significant difference in settlement between light and dark conditions. For a cohort of copepodids hatched within 24 h, Frenzl (2014) found in flume challenges that maximal infectivity was obtained at 4 days post-moult to copepodid, with the infectivity of the cohort declining by 6 days through mortalities and lower infective capabilities. Tucker *et al.* (2000a) found that infection success (measured as the proportion of larvae used for infection that were found on the fish at day 5 after infection) was approximately 75 % at 11 °C and approximately 20 % at 6.5 °C in one-day-old and three-day-old copepodids, with infection success declining significantly in seven-day-old copepodids, although lice in this experiment were collected and cultured at 10 °C before being

used in experiments, which may have affected the results. The ability of copepodids to infect hosts past seven days old is known from experiments with *L. salmonis* (Pedersen 2009), but detailed temporal infectivity profiles have not been published. However, infection success is clearly linked to both the longevity and activity of the copepodid stage. Despite infection success being dependent on copepodid age, the survival of copepodids once attached to a host was not observed to differ between copepodids that infect at different ages (Tucker *et al.*, 2000a; Pedersen, 2009), which is likely due to the commencement of feeding once attached to a host. This suggests that key determinants of variability of larval infection levels in Atlantic salmon act prior to host settlement *i.e.* within the black box comprising egg production to host contact.

#### ***4.2 Impacts of environmental variables on infection***

Host settlement success is also reduced at lower salinities, which coincides with a decrease in their energy reserves (Tucker *et al.*, 2000a,b; Bricknell *et al.*, 2006). It is likely that the physiological stress associated with reduced salinity rapidly depletes the energy reserves of copepodids, which causes premature senescence and results in levels of settlement success similar to those found in older copepodids (Bricknell *et al.*, 2006). These authors report that infection levels were reduced by 45 % at 26 ‰ (~14 % infection), 55 % at 19 ‰ (~10 % infection) and 87.5 % at 12 ‰ (~1 % infection) compared to full-strength seawater, which was not wholly attributable to reduced survival at these salinities. At 4 ‰ no copepodids were found on the fish.

While settlement success is lower with reduced energy reserves, Samsing *et al.* (2016) used degree days to normalise copepodid energy reserves cultured at different temperatures; at 30 degree days from hatching, settlement success was  $41.6 \pm 2.0$  % at 20 °C,  $53.2 \pm 2.3$  % at 10 °C and  $2.1 \pm 0.4$  % at 5 °C. Key values for infectivity are shown in Table 5.

### 4.3 Post-attachment variables

A number of variables intervene between initial attachment of the copepodid and successful moulting to the chalimus I stage. In particular, once attached, the copepodid becomes susceptible to host defences, particularly in terms of innate host immunity, often expressed through inflammatory processes. The success of the host response in controlling infection depends upon a number of variables including the species / genotype of the host fish, its age, maturity, health and welfare / stress status and interactions of immune capabilities with environmental parameters such as temperature. The role of the host in mediating infection success will only be covered briefly here as it has been extensively reviewed and investigated by previous authors (Braden *et al.* 2017, Fast 2014, Skugor *et al.* 2008, Tadiso *et al.* 2011 *inter alia*). In Atlantic salmon, initial infection by the copepodid can elicit a detectable transcriptomic host response within 1 day post infection (dpi) (Tadiso *et al.* 2011) and some Pacific salmon species, *e.g.* juvenile coho, are able to mount a rapid and successful inflammatory response following infection (Johnson and Albright, 1992; Fast *et al.*, 2002; Jones, 2011) that is capable of killing infecting copepodids within a few days. Atlantic salmon show a less developed inflammatory response and are generally considered to show a poor capacity for removing infecting copepodids (Johnson and Albright, 1992). Despite this observation, different genetic stocks or families of Atlantic salmon can show significant differences in their capacity to resist infection, although the mechanisms underlying differential resistance are currently poorly understood. Jodaa Holm *et al.* (2015) have suggested that differential resistance may reflect the ability of the host to avoid immunosuppression by the parasite. In a comparison of salmon family susceptibility, Gharbi *et al.* (2015) demonstrated a ~60% difference in the median infection count at 7 dpi (chalimus I) for the least and most susceptible salmon families tested by copepodid infection challenge

and calculated a genetic heritability of 0.3 for this trait making it a good candidate for selective breeding. The capacity of salmon to reduce infection success may also be modified by extrinsic factors such as diet and temperature. Functional feeds containing a range of active plant or bacterial extracts have, for example, been shown to have significant effects on infection success, providing infection reductions of up to 50% (Jensen *et al.* 2014, Jodaa Holm *et al.* 2016, Sutherland *et al.* 2017).

Sea lice, like other arthropod parasites, can also suppress or redirect host immune responses by the use of a range of secretory excretory products (SEPs) including prostaglandin E-2, trypsin, peroxinectin and a range of other proteases, peroxidases and potential defensin classes (Fast, 2014; Øvergård *et al.*, 2016). The success of the parasite in immunomodulating the host depends on the individual host's innate susceptibility and its state at the time of infection. Similarly, the status of the parasite can be important such that, for example, genetic family differences may affect infection success (Ljungfeldt *et al.*, 2014) although the point at which success is mediated and the mechanisms involved remain unknown.

## **5 Mortality through predation**

Once sea lice have attached to a host, their chances of survival are increased as they have a constant food supply and external factors affecting survival are relatively few, *e.g.* adverse environmental conditions, host immune response and predation by cleaner fish. During their free-swimming planktonic stages, however, they form a part of a complex plankton food web and are subject to selective and non-selective predation by other plankton and sessile filter feeders such as bivalve molluscs. Global approximations of the partitioning of wider zooplankton mortality suggest that predation accounts for 67–75 % of total mortality in the plankton (Hirst & Kiørboe, 2002). Although predation is likely to have a significant impact on sea lice survival, there are currently no estimates of sea lice predation mortality in the

literature due to the difficulty in obtaining this kind of information. Some sea lice dispersion models do include a fixed mortality rate for the free-swimming stages, *e.g.* Amundrud and Murray (2009) used a fixed mortality rate of  $0.01 \text{ h}^{-1}$  for nauplii and copepodids. Providing an estimate of predation mortality is difficult as plankton assemblages vary considerably according to season and location (*e.g.* Daewel *et al.*, 2014), and prey selection sizes vary amongst the different actively or passively predating species represented in the zooplankton community at any time (Hansen *et al.* 1994; Wirtz, 2011, Wirtz, 2012). As a consequence of a lack of specific data, the following discussion seeks to provide guidance based on wider knowledge of zooplankton, which may be used by researchers to formulate research questions or provide initial parameters for models.

### **5.1 Plankton community structure**

In regional marine ecosystems, several processes govern the structure and dynamics of plankton communities. These processes vary according to geographical location, resulting in distinct ocean regions with their own typical plankton assemblages. Small copepods dominate inshore zooplankton with their seasonal abundance following that of the phytoplankton, and clupeid and scombrid fish are the main consumers of pelagic invertebrates (Kaiser, 2005).

These broad ocean regions may further be characterised according to ocean processes in different sub-regions, *e.g.* the North Sea, the Norwegian Sea. The abundance of different species that are predators of sea lice larvae and the abundance of other prey will affect the mortality rate of sea lice larvae. Therefore, providing data on larval predation by different plankton assemblages and characterising the plankton assemblage at a specific location represents an important step in predicting mortality rates due to predation.



## 5.2 *Predator selectivity*

The body sizes of predator and prey are fundamental in the study of aquatic food webs (Brooks & Dodson, 1965; Woodward *et al.*, 2005). A ‘feeding kernel’ represents a description of the probability of prey ingestion given as a function of feeding rate vs. prey size (Figure 2) (Visser & Fiksen, 2013; Wirtz, 2014). Selective grazing in the presence of a broad spectrum of prey size plays an important role in variable feeding relationships (Sommer & Stibor, 2002), and in the case of larval sea louse predation, the abundance of similar-sized prey must be considered as well as the abundance and size selectivity of predators.

Although the relationship between predator and prey body sizes is the primary determinant of grazing selectivity, feeding modes can also affect the size range of plankton selected. Feeding modes can be broadly classified as passive and active ambush feeding, feeding-current feeding and cruise feeding (Kiørboe, 2011), and predators may adjust their feeding behaviour in response to the density of food items, (*e.g.* Kiørboe & Saiz, 1995; Boenigk & Arndt, 2002, Frost, 1972; Visser *et al.*, 2009, Saiz & Kiørboe, 1995). This behavioural plasticity shrinks the overall spectrum of potential prey towards a specific sub-range, and Wirtz (2014) describes two feeding kernels: one for ingestion, which is based on the size range of prey that can be ingested based on biomechanical principles, and one for selection, which describes the actual size range of prey selected according to the availability of prey of various sizes (Figure 2). At high prey densities, many ambush and suspension feeders, such as copepods, typically have a high selectivity resulting in a narrow selection kernel (figure 2a), whereas many facultative, omnivorous feeders, such as jellyfish, typically have broad ingestion and selection kernels (figure 2b) (Wirtz, 2014).

## 5.3 *Prey selection*

Prey size selection is determined according to the equivalent spherical diameter (ESD), which is the longest axis of the prey, *i.e.* length for sea lice larvae. Johnson and Albright (1991a) report that the length of the nauplius I was  $0.54 \pm 0.04$  mm, the nauplius II was  $0.56 \pm 0.01$  mm and the copepodid was  $0.70 \pm 0.01$  mm in *L. salmonis oncorhynchi* collected from British Columbian waters. Schram (1993) reports similar ranges for *L. salmonis salmonis* collected in Norway.

Potential predators of sea lice larvae are likely to include obligate and facultative carnivorous zooplankton and planktivorous fish, and given their geographical distribution, predators may be represented by chaetognaths, ctenophores, scyphozoa, euphausiids, mysids and scombrid and clupeid fish. In addition, the larval stages of most fish species rely on copepods as their principal dietary component (Kaiser, 2005).

Chaetognaths, or arrow worms, are important predators of copepods and are probably major contributors to the structuring of many marine ecosystems (Steele & Frost, 1977). Chaetognaths are ambush predators, and Fulton (1984) found that active copepods, such as *Acartia tonsa*, decreased in abundance in the presence of *Sagitta hispida*, whereas inactive swimmers, such as *Oithona* spp. did not as encounter rates were lower. As sea lice larvae are active swimmers, it is likely that they will be predated by chaetognaths of a suitable size category.

Ctenophores, or comb jellies, are found throughout the world's oceans, and all are predatory, feeding on zooplankton (Fowler, 1911). If food is plentiful, they can eat ten times their own weight per day (Reeve *et al.* 1978). In laboratory experiments, copepodid I larvae of *Calanus pacificus* with a mean length of 0.74 mm and mean swimming speed of  $0.32 \text{ mm s}^{-1}$ , hence similar in size to sea lice larvae, were most susceptible to predation by *P. bachei*, and later juvenile stages, which are larger, were less susceptible to predation (Greene *et al.*, 1986).

Scyphozoa, or jellyfish, are generally larger than many other predators in the plankton, and are seasonally common in many coastal environments including those most commonly employed for marine salmonid aquaculture (Doyle *et al.*, 2007). Scyphozoa typically range from 2–40 cm, and their stinging or filter-feeding tentacles enable them to ingest various zooplankton taxa of different sizes, including copepods (Purcell, 1992; Purcell *et al.*, 1994; Suchman & Sullivan, 1998). However, research has shown that scyphozoa are highly selective, and prey size has a significant impact on feeding rates (Suchman & Sullivan, 1998, 2000). As scyphozoa are neither visual nor raptorial feeders, they select prey as a consequence of prey vulnerability, and prey with faster swimming speeds and poor escape responses are most vulnerable to predation (Suchman & Sullivan, 2000).

Euphausiid and mysid shrimps are two groups of arthropods that are ubiquitous throughout the world's oceans, and due to their high abundance and position in the food chain, they are important components of marine food chains (Båmstedt & Karlson, 1998). While most are omnivorous filter feeders and feed on phytoplankton and detritus, some are carnivorous and feed on other zooplankton (Cripps & Atkinson, 2000). In the Norwegian Sea, the copepod *Calanus finmarchicus* (which has similar-sized juvenile stages to sea lice) is a dominant prey of euphausiid shrimp (Båmstedt & Karlson, 1998).

The larval stages of many fish species rely on copepods as their principal dietary component, and although larger gadoids, such as Atlantic cod (*Gadus morhua*) switch to piscivory as adults, smaller species, such as Norway pout (*Trisopterus esmarkii*) and clupeids, such as herring (*Clupea harengus*) remain planktivorous throughout their lives (Daewel *et al.*, 2014). As larval fish are active raptorial predators and rely on sight to detect prey, active prey may be more susceptible to predation. Tiselius & Jonsson (1990) and Doall *et al.* (1998) suggest that the high turn rates of sea lice copepodids during host-seeking behaviour may make them

more attractive to predators, such as fish larvae. Some adult fish, such as scombrids and clupeids, feed on plankton throughout their lives, and switch between feeding modes depending on prey density (Janssen, 1976). Zooplankton consumption by fish in the North Sea has been estimated at 19–25 g C m<sup>-2</sup> year<sup>-1</sup> of which 28 % of overall zooplankton consumption can be attributed to early life stages of fish (Heath, 2007). In frontal zones, fish larvae could consume up to 3–4 % day<sup>-1</sup> of the fraction of preferred zooplankton sizes (Munk *et al.*, 1994).

In addition to planktonic predators, sessile feeders, particularly bivalve molluscs and cnidarians, could also have a potential impact on larval sea louse survival. Bivalve molluscs, specifically the blue mussel *Mytilus edulis*, have been suggested to provide efficient clearance of mesoplankton of the same size order as sea lice larvae (Davenport *et al.*, 2000). Only blue mussels and scallops (*Placopecten magellanicus*) have been specifically investigated in terms of their ability to clear larval sea lice (Molloy *et al.*, 2011, Bartsch *et al.*, 2013). Molloy *et al.* (2011) demonstrated that mussels were capable of removing copepodids from the water column under experimental conditions and this was also demonstrated by Bartsch *et al.* (2013) who showed that mussels and scallops could remove 18–38 % of presented copepodids per hour. While it has been suggested that mussels or other bivalves might, therefore, be employed to help control sea lice on farms (Molloy *et al.*, 2011, Bartsch *et al.*, 2013), it has been noted (Bravo, pers. comm.) that close proximity of mussel farms and salmon farms in Chile has not served to reduce apparent levels of sea lice infections.

The foregoing observations on levels of predation of zooplankton support the suggestion that the mortality of free-living sea lice stages, *i.e.* nauplii and copepodids, is likely to be high during the planktonic phase.

## **6 Research gaps identified, recommendations and conclusions**

A broad range of factors impact the levels of egg production by host-attached lice and the subsequent proportion of the initial extruded egg number that go on to successfully infect fish as copepodid larvae. Figure 3 shows the stages of the sea louse life cycle that determine the number of copepodids available for infection and their infection success and summarises the factors reviewed in this study that may affect subsequent levels of infection.

A simplified conceptual framework can be employed to summarise the findings of this review, which describes the relationships between the production and loss of free-swimming larval lice and aspects of their behaviour that together determine subsequent infection levels:

$$S = EP_hP_pP_dP_sP_eI$$

Where  $S$  is number of successfully infecting copepodids,  $E$  is the number of eggs produced,  $P_h$  is the probability of hatching,  $P_p$  is the probability of avoiding predation,  $P_d$  is the probability of successful development from nauplii to copepodids,  $P_s$  is the probability of copepodid mortality due to senescence,  $P_e$  is the probability of encountering an appropriate host, and  $I$  is the mean infectivity of the copepodid population. The operational use of this conceptualised framework requires the estimation of the components of each of these variables, which are themselves influenced by a range of biotic (*e.g.* host) and abiotic (*e.g.* water temperature) factors and each other, *i.e.* they are not independent. As each component (or loss) is multiplicative, the uncertainties in each component may result in very wide error margins in  $S$ . Therefore, it is important to define and continue to refine each component through extensive data collection and parameterisation to reduce the level of error.

By forming a table of these variables and the observable factors that may influence them (Table 6), it is clear that there are a considerable number of permutations, each requiring observational data to allow variables to be fully defined. While a number of these variables have been previously investigated, as described in this review, a lack of data for some variables

results in an incomplete dataset (Table 6). Furthermore, a lack of standardisation and consistency across different studies due to various experimental conditions and the origin of experimental lice, *e.g.* of Atlantic or Pacific origin, farmed or wild origin, cold-adapted or not, means that many data points are not directly comparable. In addition, some studies are based on laboratory experiments conducted under controlled conditions, whereas others are based on field data. Gravid (1996) recorded the widths of nauplius I larvae and the lipid reserves from field-collected lice at different times of year, and although no other studies considered seasonal variations in their experiments *per se* (Table 6), seasonal variation subsumes a number of observable/observed factors, such as temperature, photoperiod and salinity, and other factors that are not considered here, such as host condition and plankton assemblages.

## **6.1 Key gaps in knowledge identified**

There are a very great number of gaps in our knowledge concerning the variables affecting levels of sea louse infections. Some variables, however, are likely to have both a greater proportional/numerical impact and to be more tractable to parameterisation by experimental means. These are addressed below with reference to the conceptual framework defined above.

### **6.1.1 Egg production ( $E$ ), egg viability and hatching success ( $P_h$ )**

Previous estimates of egg production in the literature vary across more than an order of magnitude, are relatively inconsistent and are incomplete in their coverage of relevant factors. As this is the key input variable driving subsequent modelled infection levels, better estimates of production are an obvious priority. In addition to this, it is clear from the relatively sparse

earlier studies that have been conducted that egg viability and hatching success are rarely, if ever, 100 % and can be substantially lower than this according to a range of factors (Table 2).

Egg production level is influenced by a broad range of factors including temperature (and temperature adaptation), salinity, host state (nutrition, immunity, stress, species, genotype), egg batch and others. For this reason, it will be extremely difficult to establish realistic values through tightly controlled laboratory experiments alone. Egg production can, however, easily be established through a programme of farm sampling over a year, with counts of eggs per millimetre and the measurement of egg string lengths being conducted on-farm using a stereomicroscope or in the laboratory following sample preservation. Laboratory analysis could also employ image analysis to increase accuracy and sample throughput. During the sampling period, the recording of farm metadata, such as temperature, salinity, salmon stock, feed source, treatment regime *etc.*, would allow an accurate and informative predictive model to be produced. In order to give a better picture of total egg production, samples from wild salmonids would also be helpful as it is well-recognised that egg strings sourced from lice on wild fish tend to have higher numbers of eggs (Tully & Whelan, 1993; Pike & Wadsworth, 1999).

Laboratory experiments could investigate controllable factors, *e.g.* using a range of temperatures and salinities, ideally for lice sampled from different ambient temperatures, *e.g.* winter, spring, and summer.

The viability of eggs and hatching success are key mediators of the final number of released larvae. These parameters can be obtained by examining and hatching egg strings from challenges and/or farm samples under controlled conditions of temperature and salinity.

### 6.1.2 Predation in plankton ( $P_p$ )

The level of predation of larval sea lice in the plankton remains unknown. However, it is clear from other plankton studies that losses to predation are likely to be substantial. In addition, the level of predation will vary according to season, local weather conditions and the composition of the plankton assemblage at any given time. Knowledge of predation levels will not only facilitate more accurate modelling of infection levels but could also guide co-ordinated treatment strategies at particular times of year.

Even with good estimates of larval production, the fate of larvae in the plankton is a key mediator of numbers available to infect fish. Plankton studies are notoriously difficult and are not easily amenable to laboratory-based experiments. To achieve estimates of mortality in plankton, mesocosm studies offer the best approach, whereby in different seasons local plankton are enclosed in a mesocosm, and a known number of larval sea lice are introduced to the system. Following a period to allow for predation, the filtering of the mesocosm will allow estimations of plankton types/species present and the clearance rates of sea louse larvae. The use of molecular tools might also allow an investigation of the major predators in any given plankton sample.

Using the same system with introduced ‘sentinel’ salmonids, one could also establish the resulting infection levels, which, while not wholly realistic, would allow some estimation of both the effects of predation and also encounter rate on infection success.

### 6.1.3 Infectivity profile ( $I$ )

To date, there has been a tendency to equate the number of copepodids in the water column with the number of infecting individuals. From previous observations, however, it is apparent that there is a profile of infectivity, *i.e.* the ability of lice encountering a fish to infect it as they



age, with newly moulted individuals being less infective than those having matured for 1–2 days and a subsequent decline of infectivity towards death. Infection success requires definition as not all copepodids that attach to a host may establish a successful infection; the number of copepodids developing to the chalimus I stage and developing a permanent attachment via a frontal filament may be an appropriate measure of infection success. Even under the optimal conditions of an experimental infection challenge, the infective success of maximally infective copepodids is rarely higher than 50 % and is frequently lower. From the literature, few researchers have attempted to establish infection profiles for cohorts of copepodids under different conditions of, for example, temperature, salinity and current speed, despite clear evidence that these factors will all affect infection success. Most challenge experiments employ static tanks and long exposure times, providing a totally inaccurate reflection of probabilities for real-world infection success.

While the infectivity profile needs to be better established under laboratory conditions, these will not fully reflect field conditions but will tend to provide an overestimate of infection success rate. Using standard tank challenges it is possible to profile the infectivity of copepodids with age and under different temperature and salinity conditions. However, a more accurate reflection of infectivity can be achieved using flume experiments where fish are exposed to copepodids under current flow conditions more reflective of field conditions.

One important source of potentially valuable data concerning losses incurred between egg hatching and the reinfection of hosts is the detailed farm louse counts already conducted in many countries. Assuming knowledge of seasonal levels of egg production and viability, which may be easily obtained, the annual profile of copepodid/chalimus counts, can, at least for some more hydrographically constrained regions, provide an indication of the proportion of hatched larvae that successfully re-establish infections on fish.

952

## 953 **6.2 Co-ordinated research**

954 In order to obtain the greatest benefits from modelling studies, the gaps identified need to be  
955 filled for lice and environments in all of the regions experiencing problems with *L. salmonis*  
956 and independently for other species *e.g.* *C. rogercresseyi*. This means co-ordinating  
957 international efforts to ensure that studies are inter-comparable, and this would ideally be  
958 achieved through international agreements for matched funding by key national industry and  
959 government funders.

960

## 961 **6.3 Conclusions**

962 The estimation of lice burdens on wild and cultured fish can inform the timing of pest  
963 management decisions in salmonid aquaculture. In the life cycle of the sea louse, egg  
964 production, survival of free-swimming stages and infectivity of survivors are key determinants  
965 of the number of lice re-establishing host infection. Despite several decades of research,  
966 however, knowledge of this area of sea louse biology is lacking, which confounds the accurate  
967 estimation of lice infections using epidemiological modelling. Even where parameters have  
968 been measured by researchers, the wide variety of data sources and experimental approaches  
969 employed, limits the possibility of providing ‘best’ or consensus values for use in modelling.  
970 With further research of the key variables that affect the production and survival of free-  
971 swimming larval sea lice, it should be possible to more accurately model the production and  
972 dispersal of lice from cage aquaculture and wild fish, which will inform the optimum timing  
973 of pest management procedures. Furthermore, with an improved knowledge of larval sea louse  
974 mortality, it may be possible to incorporate natural processes into management decisions and  
975 to manage timing of treatments appropriately, *e.g.* reflecting larval predation following spring

algal blooms. While many aspects of louse biology are important in determining the numbers of lice available for infection, care should be taken to avoid the over-parameterisation of sea louse infection models. The identification of the key variables from the complex biology of sea lice that have the greatest impact on their numbers can be achieved through a sensitivity analysis of model parameters. Accurate predictions of sea lice infections are a single component of integrated pest management protocols, and when used in conjunction with the continuous monitoring of lice populations on farmed fish and effective treatment procedures, it should be possible to minimise the environmental and economic impact of these pathogens on farmed and wild salmonids.

## **7 Acknowledgements**

This study was funded by a Scottish Aquaculture Research Forum grant (SARF108). The authors would like to thank Dr Darren Green for guidance on mathematical modelling.

## 8 References

- Adams, T., Black, K., MacIntyre, C., MacIntyre, I., and Dean, R. 2012. Connectivity modelling and network analysis of sea lice infection in Loch Fyne, west coast of Scotland. *Aquaculture Environment Interactions*, 3: 51–63.
- Aldrin, M., Storvik, B., Kristoffersen, A. B., and Jansen, P. A. 2013. Space-time modelling of the spread of salmon lice between and within Norwegian marine salmon farms. *PLoS ONE*, 8:1–10.
- Amundrud, T. L., and Murray, A. G. 2009. Modelling sea lice dispersion under varying environmental forcing in a Scottish sea loch. *Journal of Fish Diseases*, 32: 27–44.
- Asplin, L., Johnsen, I. A., Sandvik, A. D., Albretsen, J., Sundfjord, V., Aure, J., and Boxaspen, K. K. 2014. Dispersion of salmon lice in the Hardangerfjord. *Marine Biology Research*, 10: 216–225.
- Asplin, L., Boxaspen, K. K., and Sandvik, A. D. 2011. Modelling the distribution and abundance of planktonic larval stages of *Lepeophtheirus salmonis* in Norway. In *Salmon lice: an integrated approach to understanding parasite abundance and distribution*, pp. 31–50 Ed. by S. Jones and R. Beamish. Oxford: Wiley-Blackwell. 336 pp.
- Asplin, L., Boxaspen, K., and Sandvik, A. D. 2004. Modelled distribution of salmon lice in a Norwegian fjord. *ICES CM Documents* (pp. 1–12).
- Bailey, R. J., Birkett, M. A, Ingvarsdóttir, A., Mordue (Luntz), A. J., Mordue, W., O’Shea, B., Pickett, J. A., and Wadhams, L. J. 2006. The role of semiochemicals in host location and non-host avoidance by salmon louse (*Lepeophtheirus salmonis*) copepodids. *Canadian Journal of Fisheries and Aquatic Science*, 63: 448–456.

- 1011 Båmstedt, U. and Karlson, K. 1998. Euphausiid predation on copepods in coastal waters of  
1012 the Northeast Atlantic. *Marine Ecology Progress Series*, 172: 149–168.
- 1013 Bartsch, A., Robinson, S. M. C., Liutkus, M., Ang, K. P., Webb, J., and Pearce, C. M. 2013,  
1014 Filtration of sea louse, *Lepeophtheirus salmonis*, copepodids by the blue mussel,  
1015 *Mytilus edulis*, and the Atlantic sea scallop, *Placopecten magellanicus*, under different  
1016 flow, light and copepodid-density regimes. *Journal of Fish Diseases*, 36: 361–370.
- 1017 Bjørn, P. A., Asplin, L., Karlsen, Ø., Finstad, B., and Taranger, G. L. 2014. Utvikler  
1018 modelsystem for få kontroll på Lusa. (English translation: Development of a system of  
1019 models to control lice). In *Fisken og Havet*, særnummer 1–2014.  
1020 Havforskningsrapporten 2014.
- 1021 Boenigk, J. and Arndt, H. 2002. Bacterivory by heterotrophic flagellates: community structure  
1022 and feeding strategies. *Antonie van Leeuwenhoek*, 81: 465–480.
- 1023 Boxaspen, K. and Næss, T. 2000. Development of eggs and the planktonic stages of salmon  
1024 lice (*Lepeophtheirus salmonis*) at low temperatures. *Contributions to Zoology*, 69: 51–  
1025 56.
- 1026 Braden L. M., Sutherland B. J., Koop B. F., and Jones S. R. 2017. Enhanced transcriptomic  
1027 responses in the Pacific salmon louse *Lepeophtheirus salmonis oncorhynchi* to the  
1028 non-native Atlantic Salmon *Salmo salar* suggests increased parasite fitness. *BMC*  
1029 *Genomics*, 18: 1.
- 1030 Bricknell, I. R., Dalesman, S. J., Shea, B. O., Pert, C. C., and Luntz, A. J. M. 2006. Effect of  
1031 environmental salinity on sea lice *Lepeophtheirus salmonis* settlement success.  
1032 *Diseases of Aquatic Organisms*, 71: 201–212.

- 1033 Bron, J. E. 1993. A study of the biology and behaviour of the copepodid larva of the salmon  
1034 louse *Lepeophtheirus salmonis* (Krøyer, 1837) (Copepoda; Caligidae). PhD Thesis,  
1035 University of Stirling.
- 1036 Bron, J. E., Sommerville, C., and Rae, G. H. 1993. Aspects of the behaviour of copepodid  
1037 larvae of the salmon louse. *In* Pathogens of wild and farmed fish: Sea lice, 2nd edn,  
1038 pp. 125–142. Ed. by G. A. Boxshall and D. Defaye. Ellis Horwood Ltd., New York.  
1039 378 pp.
- 1040 Bron, J. E., Sommerville, C., Jones, M., and Rae, G. H. (1991). The settlement and attachment  
1041 of early stages of the salmon louse, *Lepeophtheirus salmonis* (Copepoda: Caligidae)  
1042 on the salmon host, *Salmo salar*. *Journal of Zoology*, 224: 201–212.
- 1043 Brooks, K. M. 2005. The effects of water temperature, salinity, and currents on the survival  
1044 and distribution of the infective copepodid stage of sea lice (*Lepeophtheirus salmonis*)  
1045 originating on Atlantic salmon farms in the Broughton Archipelago of British  
1046 Columbia, Canada. *Reviews in Fisheries Science*, 13: 177–204.
- 1047 Brooks, J. L. and Dodson, S. I. 1965. Predation, body Size, and composition of plankton.  
1048 *Science*, 150: 28–35.
- 1049 Carr, J. and Whoriskey, F. 2004. Sea lice infestation rates on wild and escaped farmed Atlantic  
1050 salmon (*Salmo salar* L.) entering the Magaguadavic River, New Brunswick.  
1051 *Aquaculture Research*, 35: 723–729.
- 1052 Cavaleiro, F. I. and Santos, M. J. 2014. Egg number-egg size: An important trade-off in  
1053 parasite life history strategies. *International Journal for Parasitology*, 44: 173–182.

- 1054 Colavecchia, M., Katopodis, C., Goosney, R., Scruton, D. A. and McKinley, R. S. 1998.  
1055 Measurement of burst swimming performance in wild Atlantic salmon (*Salmo salar*)  
1056 using digital telemetry. *Regulated Rivers: Research and Management* 14: 41–51.
- 1057 Cook, P. F., McBeath, S. J., Bricknell, I. R., and Bron, J. E. (2010). Determining the age of  
1058 individual *Lepeophtheirus salmonis* (Krøyer, 1837) copepodids by measuring stored  
1059 lipid volume; proof of principle. *Journal of Microscopy*, 240: 83–6.
- 1060 Corner, R. A., Brooker, A. J., Telfer, T. C. and Ross, L. G. 2006. A fully integrated GIS-based  
1061 model of particulate waste distribution from marine fish-cage sites. *Aquaculture*, 258:  
1062 299–311.
- 1063 Costello, M. J., Burrridge, L., Chang, B., and Robichaud, L. 2004. Sea Lice 2003 - Proceedings  
1064 of the sixth international conference on sea lice biology and control. *Aquaculture*  
1065 *Research*, 35: 711–712.
- 1066 Costelloe, M., Costelloe, J., O'Donohoe, G., Coghlan, N. J., Oonk, M., and Van Der Heijden,  
1067 Y. 1998a. Planktonic distribution of sea lice larvae, *Lepeophtheirus salmonis*, in  
1068 Killary Harbour, West Coast of Ireland. *Journal of the Marine Biological Association*  
1069 *of the UK*, 78: 853–874.
- 1070 Costelloe, M., Costelloe, J., Coghlan, N., Donohoe, G. O., and Connor, B. O. 1998b.  
1071 Distribution of the larval stages of *Lepeophtheirus salmonis* in three bays on the west  
1072 coast of Ireland. *ICES Journal of Marine Science*, 55: 181–187.
- 1073 Cripps, G. C. and Atkinson, A. 2000. Fatty acid composition as an indicator of carnivory in  
1074 Antarctic krill, *Euphausia superba*. *Canadian Journal of Fisheries and Aquatic*  
1075 *Sciences*, 57: 31–37.

1076 Daewel, U., Hjøllø, S. S., Huret, M., Ji, R., Maar, M., Niiranen, S., and Wolfshaar, K. E. Van  
1077 De. 2014. Predation control of zooplankton dynamics: a review of observations and  
1078 models. *ICES Journal of Marine Science*, 71: 254–271.

1079 Davenport, J., Smith, R., and Packer, M., 2000. Mussels *Mytilus edulis*: significant consumers  
1080 and destroyers of mesoplankton. *Marine Ecology Progress Series*, 198: 131–137.

1081 Devine, G. J., Ottir, A. I., Mordue, W., Pike, A. W., Pickett, J., Duce, I. A. N., and Luntz, A.  
1082 J. M. 2000. Salmon lice, *Lepeophtheirus salmonis*, exhibit specific chemotactic  
1083 responses to semiochemicals originating from the salmonid, *Salmo salar*. *Journal of*  
1084 *Chemical Ecology*, 26: 1833–1847.

1085 Doall, M. H., Colin, S. P., Strickler, J. R., and Yen, J. 1998. Locating a mate in 3D: the case  
1086 of *Temora longicornis*. *Philosophical Transactions of the Royal Society B: Biological*  
1087 *Sciences*, 353: 681–689.

1088 Doyle, T. K., Houghton, J. D. R., Buckley, S. M., Hays, G. C., and Davenport, J. 2007. The  
1089 broad-scale distribution of five jellyfish species across a temperate coastal  
1090 environment. *Hydrobiologia*, 579: 29–39.

1091 Fast, M. D. 2014. Fish immune responses to parasitic copepod (namely sea lice) infection.  
1092 *Developmental and Comparative Immunology*, 43: 300–312.

1093 Fast M. D., Ross N. W., Mustafa A., Sims D. E., Johnson S. C., Conboy G. A., Speare D. J.,  
1094 Johnson G., and Burka J. F. 2002. Susceptibility of rainbow trout *Oncorhynchus*  
1095 *mykiss*, Atlantic salmon *Salmo salar* and coho salmon *Oncorhynchus kisutch* to  
1096 experimental infection with sea lice *Lepeophtheirus salmonis*. *Diseases of Aquatic*  
1097 *Organisms*, 52: 57–68.



- 1098 Fields, D. M., Skiftesvik, A. B., and Browman, H. I. 2017. Behavioural responses of infective-  
1099 stage copepodids of the salmon louse (*Lepeophtheirus salmonis*, Copepoda: Caligidae)  
1100 to host-related sensory cues. *Journal of Fish Diseases*, 00: 1–10.
- 1101 Finstad, B., Grimnes, A., and Hvidsten, N. A. 2000. Laboratory and field investigations of  
1102 salmon lice [*Lepeophtheirus salmonis* (Kroyer)] infestation on Atlantic salmon (*Salmo*  
1103 *salar* L.) post-smolts. *Aquaculture Research*, 1: 795–803.
- 1104 Forward, R. J. and Douglass, J. 1986. Crustacean larval visual sensitivity during diel vertical  
1105 migration. *In* *Proceedings of the 21st European Marine Biology Symposium, Gdansk,*  
1106 *14<sup>th</sup> September 1986*, pp. 59–66.
- 1107 Fowler, G. H. 1911. Ctenophora. *In* *Encyclopaedia Britannica*. 7, 11<sup>th</sup> edn, p. 593. Ed. By H.  
1108 Chisholm. Cambridge University Press.
- 1109 Frenzl, B. 2014. Understanding key factors associated with the infection of farmed Atlantic  
1110 Salmon by the salmon louse, *Lepeophtheirus salmonis*. PhD thesis, University of  
1111 Stirling.
- 1112 Frost, B. W. 1972. Effects of size and concentration of food particles on the feeding behaviour  
1113 of the marine planktonic copepod *Calanus pacificus*. *Limnology and Oceanography*,  
1114 17: 805–815.
- 1115 Fulton, R. S. 1984. Effects of chaetognath predation and nutrient enrichment on enclosed  
1116 estuarine copepod communities. *Oecologia*, 62: 97–101.
- 1117 Genna, R. L. 2002. The chemical ecology, physiology and infection dynamics of the sea louse  
1118 copepodid, *Lepeophtheirus salmonis* Kroyer. PhD thesis, University of Aberdeen.

- 1119 Gharbi, K., Matthews, L., Bron, J., Roberts, R., Tinch, A., and Stear, M. 2015. The control of  
1120 sea lice in Atlantic salmon by selective breeding. *Journal of the Royal Society Interface*  
1121 12, 8pp.
- 1122 Gravid, H. R. 1996. Studies on the biology and ecology of the free swimming larval stages of  
1123 *Lepeophtheirus salmonis* (Kroyer, 1838) and *Caligus elongatus* Nordmann, 1832  
1124 (Copepoda: Caligidae). PhD thesis, University of Stirling.
- 1125 Grayson, T. H., John, R. J., Wadsworth, S., Greaves, K., Cox, D., Roper, J., Wrathmell, A. B.,  
1126 Gilpin, M. L., and Harris, J. E. 1995. Immunization of Atlantic salmon against the  
1127 salmon louse: identification of antigens and effects on louse fecundity. *Journal of Fish*  
1128 *Biology*, 47: 85–94.
- 1129 Greene, C. H., Landry, M. R., and Monger, B. C. 1986. Foraging behavior and prey selection  
1130 by the ambush entangling predator *Pleurobrachia bachei*. *Ecology*, 67: 1493–1501.
- 1131 Hahnenkamp, L. and Fyhn, H. J. 1985. The osmotic response of salmon louse, *Lepeophtheirus*  
1132 *salmonis* (Copepoda: Caligidae), during the transition from sea water to fresh water.  
1133 *Journal of Comparative Physiology B*, 155: 357–365.
- 1134 Hansen, B., Bjornsen, P. K., and Hansen, P. J. 1994. The size ratio between planktonic  
1135 predators and their prey. *Limnology and Oceanography* 39: 395–403
- 1136 Haury, L. and Weihs, D. 1976. Energetically efficient swimming behaviour of negatively  
1137 buoyant zooplankton. *Limnology and Oceanography*, 21: 797–803.
- 1138 Heath, M. R. 2007. The consumption of zooplankton by early life stages of fish in the North  
1139 Sea. *ICES Journal of Marine Science*, 64: 1650–1663.

- 1140 Heuch, P. A. 1995. Experimental evidence for aggregation of salmon louse copepodids  
1141 (*Lepeophtheirus salmonis*) in step salinity gradients. Journal of the Marine Biological  
1142 Association of the UK, 75: 927–939.
- 1143 Heuch, P. A., Doall, M. H., and Yen, J. 2007. Water flow around a fish mimic attracts a  
1144 parasitic and deters a planktonic copepod. Journal of Plankton Research,  
1145 29(Supplement 1): i3–i16.
- 1146 Heuch, P. A., Parsons, A., and Boxaspen, K. 1995. Diel vertical migration: A possible host-  
1147 finding mechanism in salmon louse (*Lepeophtheirus salmonis*) copepodids? Canadian  
1148 Journal of Fisheries and Aquatic Science, 52: 681–689.
- 1149 Heuch, P. A. and Karlsen, E. 1997. Detection of infrasonic water oscillations by copepodids  
1150 of *Lepeophtheirus salmonis* (Copepoda Caligida). Journal of Plankton Research, 19:  
1151 735–747.
- 1152 Heuch, P. A. and Mo, T. A. 2001. A model of salmon louse production in Norway: effects of  
1153 increasing salmon production and public management measures. Diseases of Aquatic  
1154 Organisms, 45: 145–52.
- 1155 Heuch, P. A., Nordhagen, J. R., and Schram, T. A. 2000. Egg production in the salmon louse  
1156 (*Lepeophtheirus salmonis* (Kroyer)) in relation to origin and water temperature.  
1157 Aquaculture Research, 31: 805–814.
- 1158 Hirst, A. and Kiørboe, T. 2002. Mortality of marine planktonic copepods: global rates and  
1159 patterns. Marine Ecology Progress Series, 230: 195–209.
- 1160 Hogans, W. E., and Trudeau, D. J. 1989. Preliminary studies on the biology of sea lice, *Caligus*  
1161 *elongatus*, *Caligus curtus* and *Lepeophthierus salmonis* (Copepoda: Caligoida)

1162 parasitic on cage-cultured salmonids in the lower Bay of Fundy. Canadian Technical  
1163 Report of Fisheries and Aquatic Science, 1715: iv+14pp.

1164 Ingvarsdottir, A., Birkett, M. A., Duce, I., Genna, R. L., Mordue, W., Pickett, J. A., Wadhams,  
1165 L. J., and Mordue (Luntz), A. J. 2002. Semiochemical strategies for sea louse control:  
1166 host location cues. Pest Management Science, 58: 537–545.

1167 Iversen A., Hermansen Ø., Andreassen O., Brandvik R. K., Marthinussen A., and Nystøyl R.  
1168 2015 Kostnadsdrivere i lakseoppdrett (Cost Drivers in Salmon Farming). No. 41/2015,  
1169 1–43. Nofima [in Norwegian].

1170 Janssen, J. 1976. Feeding Modes and Prey Size Selection in the Alewife (*Alosa*  
1171 *pseudoharengus*). Journal of the Fisheries Research Board of Canada, 33: 1972–1975.

1172 Jansen , P.A., Kristoffersen, A.B., Viljugrein, H., Jimenez, D., Aldrin, M. and Stien, A. 2012.  
1173 Sea lice as a density-dependent constraint to salmonid farming. Proceedings of the  
1174 Royal Society B, 279: 2330–2338.

1175 Jensen, L., Provan, F., Larssen, E., Bron, J., and Obach, A. (2014). Reducing sea lice  
1176 (*Lepeophtheirus salmonis*) infestation of farmed Atlantic salmon (*Salmo salar* L.)  
1177 through functional feeds. Aquaculture Nutrition, 21.

1178 Jodaa Holm, H., Wadsworth, S., Bjelland, A. K., Krasnov, A., Evensen, Ø., and Skugor, S.  
1179 2016. Dietary phytochemicals modulate skin gene expression profiles and result in  
1180 reduced lice counts after experimental infection in Atlantic salmon. Parasites and  
1181 Vectors. 9: 271.

1182 Jodaa Holm, H., Santi, N., Kjølglum, S., Perisic, N., Skugor, S., and Evensen, Ø. 2015.  
1183 Difference in skin immune responses to infection with salmon louse (*Lepeophtheirus*

1184 *salmonis*) in Atlantic salmon (*Salmo salar* L.) of families selected for resistance and  
 1185 susceptibility. Fish and Shellfish Immunology, 42: 384–94.

1186 Johannessen, A. 1975. Salmon louse, *Lepeophtheirus salmonis*, Kroyer (Copepoda,  
 1187 Caligidae). Independent larval stages, growth and infection in salmon (*Salmo salar* L.)  
 1188 from breeding plants and commercial catches in west Norwegian waters 1973–1974.  
 1189 Thesis in Fish Biology, University of Bergen.

1190 Johannessen, A. 1977. Early stages of *Lepeophtheirus salmonis* (Copepoda, Caligidae). Sarsia,  
 1191 63: 169–176.

1192 Johnsen, I., Fiksen, Ø., Sandvik, A. D., and Asplin, L. 2014. Vertical salmon louse behaviour  
 1193 as a response to environmental conditions and its influence on regional dispersion in a  
 1194 fjord system. Aquaculture Environment Interactions, 5: 127–141.

1195 Johnson S. C. and Albright L. J. 1992. Comparative susceptibility and histopathology of the  
 1196 response of naïve Atlantic, Chinook and coho salmon to experimental infection with  
 1197 *Lepeophtheirus salmonis* (Copepoda: Caligidae). Diseases of Aquatic Organisms, 14:  
 1198 179–193.

1199 Johnson, S., and Albright, L. 1991a. The developmental stages of *Lepeophtheirus salmonis*  
 1200 (Kroyer, 1837) (Copepoda: Caligidae). Canadian Journal of Zoology, 69: 929–950.

1201 Johnson, S. C. and Albright, L. J. 1991b. Development, growth, and survival of  
 1202 *Lepeophtheirus salmonis* (Copepoda: Caligidae) under laboratory conditions.  
 1203 Canadian Journal of Zoology, 69: 929–950.

1204 Johnston, C. E. and Dykeman, D. 1987. Observations on body proportions and egg production  
 1205 in the female parasitic copepod (*Salmincola salmoneus*) from the gills of Atlantic

1206 salmon (*Salmo salar*) kelts exposed to different temperatures and photoperiods.  
 1207 Canadian Journal of Zoology, 65: 415–419.

1208 Jones, S. R. M. 2011. Mechanisms of Resistance among Salmon to the Parasitic Copepod  
 1209 *Lepeophtheirus salmonis*. Journal of Aquaculture Research and Development, S2:003.

1210 Jonsson, N., Jonsson, B., and Hansen, L. P. (1990). Partial segregation in the timing of  
 1211 migration of Atlantic salmon of different ages. Animal Behaviour, 40: 313–321.

1212 Kaiser, M. J. (Ed.). (2005). Marine ecology: processes, systems, and impacts. Oxford  
 1213 University Press.

1214 Kiørboe, T. 2011. How zooplankton feed: mechanisms, traits and trade-offs. Biological  
 1215 Reviews, 86: 311–39.

1216 Kiørboe, T. and Saiz, E. 1995. Planktivorous feeding in calm and turbulent environments, with  
 1217 emphasis on copepods. Marine Ecology Progress Series, 122: 135–145.

1218 Koop, K. and Field, J. G. (1980). The influence of food availability on population dynamics  
 1219 of a supralittoral isopod, *Ligia dilatata* brandt. Journal of Experimental Marine  
 1220 Biology and Ecology, 48: 61–72.

1221 Kristoffersen, A. B., Jimenez, D., Viljugrein, H., Grøntvedt, R., Stien, A., and Jansen, P. A.  
 1222 2014. Large scale modelling of salmon lice (*Lepeophtheirus salmonis*) infection  
 1223 pressure based on lice monitoring data from Norwegian salmonid farms. Epidemics,  
 1224 9: 31–39.

1225 Krkošek, M., Lewis, M.A and Volpe, J. P. 2005. Transmission dynamics of parasitic sea lice  
 1226 from farm to wild salmon. Proceedings of the Royal Society B, 272: 689–96.

- 1227 Leclercq, E., Davie, A., and Migaud, H. 2013. Delousing efficiency of farmed ballan wrasse  
1228 (*Labrus bergylta*) against *Lepeophtheirus salmonis* infecting Atlantic salmon (*Salmo*  
1229 *salar*) post-smolts. Pest Management Science, 70: 1274–1282.
- 1230 Ljungfeldt, L. E. R., Espedal, P. G., Nilsen, F., Skern-Mauritzen, M., and Glover, K. A. 2014.  
1231 A common-garden experiment to quantify evolutionary processes in copepods: The  
1232 case of emamectin benzoate resistance in the parasitic sea louse *Lepeophtheirus*  
1233 *salmonis*. BMC Evolutionary Biology, 14: 108.
- 1234 Lyse, A. A., Stefansson, S. O., and Ferno, A. 1998. Behaviour and diet of sea trout post-smolts  
1235 in a Norwegian fjord system. Journal of Fish Biology, 52: 923–936.
- 1236 MacKinnon, B. M. 1998. Host factors important in sea lice infections. ICES Journal of Marine  
1237 Science, 55: 188–192.
- 1238 MacKinnon, B. M., Bailey, J. K., O'Flynn, F. M., and Friars, G. W. 1995. Can selection  
1239 improve resistance? Northern Aquaculture, 11: 3.
- 1240 Maranhão, P. and Marques, J. C. (2003). The influence of temperature and salinity on the  
1241 duration of embryonic development, fecundity and growth of the amphipod  
1242 *Echinogammarus marinus* Leach (Gammaridae). Acta Oecologica, 24: 5–13.
- 1243 McKibben, M. A. and Hay, D. W. 2004. Distributions of planktonic sea lice larvae  
1244 *Lepeophtheirus salmonis* in the inter-tidal zone in Loch Torridon, Western Scotland in  
1245 relation to salmon farm production cycles. Aquaculture Research, 35: 742–750.
- 1246 Molloy S. D., Pietrak, M. R., Bouchard, D. A., and Bricknell, I. 2011. Ingestion of  
1247 *Lepeophtheirus salmonis* by the blue mussel *Mytilus edulis*. Aquaculture, 311: 61–64.

- 1248 Mordue Luntz, A. J. and Birkett, M. A. 2009. A review of host finding behaviour in the  
1249 parasitic sea louse, *Lepeophtheirus salmonis* (Caligidae: Copepoda). Journal of Fish  
1250 Diseases, 32: 3–13.
- 1251 Munk, P., Nielsen, T. G., and Castle, C. 1994. Trophodynamics of the plankton community at  
1252 Dogger Bank: predatory impact by larval fish. Journal of Plankton Research, 16: 1225–  
1253 1245.
- 1254 Murray, A. G. 2002. Using observed load distributions with a simple model to analyse the  
1255 epidemiology of sea lice (*Lepeophtheirus salmonis*) on sea trout (*Salmo trutta*). Pest  
1256 Management Science, 58: 585–94.
- 1257 Murray, A. G., and Amundrud, T. L. 2007. Using coupled biophysical – particle tracking  
1258 models of sea lice in Loch Torridon. In: Proceedings of the International Congress on  
1259 Modelling and Simulation, Christchurch, New Zealand, December 2007. pp. 2848–  
1260 2853.
- 1261 Nilsen, F. 2016. Sea lice vaccine. Presentation available online  
1262 [http://www.fhf.no/media/127279/10\\_-\\_vaksine\\_-\\_grunnleggende\\_kunnskap\\_-](http://www.fhf.no/media/127279/10_-_vaksine_-_grunnleggende_kunnskap_-_nilsen_-_slrc-uib.pdf)  
1263 [\\_nilsen\\_-\\_slrc-uib.pdf](http://www.fhf.no/media/127279/10_-_vaksine_-_grunnleggende_kunnskap_-_nilsen_-_slrc-uib.pdf). pp 1-20. Accessed 29/12/2017.
- 1264 Núñez-Acuña, G., Marambio, J. P., Valenzuela, T., Wadsworth, S., and Gallardo-Escárate, C.  
1265 2016. Antimicrobial peptides from *Salmon salar* skin induce frontal filament  
1266 development and olfactory/cuticle-related genes in the sea louse *Caligus*  
1267 *rogercresseyi*. Aquaculture, 464: 171-177.
- 1268 Øvergård, A., Hamre, L. A., Harasimczuk, E., Dalvin, S. Nilsen, F., and Grotmol, S. 2016.  
1269 Exocrine glands of *Lepeophtheirus salmonis* (Copepoda: Caligidae): Distribution,



- developmental appearance, and site of secretion. *Journal of Morphology*, 277: 1097–4687.
- Pedersen, T. M. 2009. Instar growth and development of *Leophtheirus salmonis* copepodids. MSc Thesis, University of Bergen.
- Pike, A. and Wadsworth, S. 1999. Sea lice on salmonids: their biology and control. *Advances in Parasitology*, 44: 233–337.
- Pike, A. W., Mordue (Luntz), A. J., and Ritchie, G. 1993. The development of *Caligus elongatus* Nordmann from hatching to copepodid in relation to temperature. *In* Pathogens of wild and farmed fish: Sea lice, 2nd edn, pp. 51–60. Ed. by G. A. Boxshall and D. Defaye. Ellis Horwood Ltd., New York. 378 pp.
- Purcell, J. E. 1992. Effects of predation by the scyphomedusan *Chrysaora quinquecirrha* on zooplankton. *Marine Ecology Progress Series*, 87: 65–76.
- Purcell, J. E., Nemazie, D. A., Dorsey, S. E., Houde, E. D., and Gamble, J. C. 1994. Predation mortality of bay anchovy *Anchoa mitchilli* eggs and larvae due to scyphomedusae and ctenophores in Chesapeake Bay. *Marine Ecology Progress Series*, 114: 47–58.
- Reeve, M. R., Walter, M. A., and Ikeda, T. 1978. Laboratory studies of ingestion and food utilization in lobate and tentaculate ctenophores. *Limnology and Oceanography*, 23: 740–751.
- Ritchie, G., Mordue, A. J., Pike, A. W., and Rae, G. H. 1993. The reproductive output of *Lepeophtheirus salmonis* adult females in relation to seasonal variability of temperature and photoperiod. *In* Pathogens of wild and farmed fish: Sea lice, 2nd edn, pp. 153–165. Ed. by G. A. Boxshall and D. Defaye. Ellis Horwood Ltd., New York. 378 pp.

- 1293 Rødseth, G. I. 2016. Beregninger på fem milliarder i lusekostnader er bare deler av sannheten.  
 1294 iLaks.no. Retrieved from [https://ilaks.no/beregninger-pa-fem-milliarder-i-](https://ilaks.no/beregninger-pa-fem-milliarder-i-lusekostnader-er-bare-deler-av-sannheten/)  
 1295 lusekostnader-er-bare-deler-av-sannheten/ Last accessed 10/06/17. [in Norwegian].
- 1296 Saiz, E. and Kiørboe, T. 1995. Predatory and suspension feeding of the copepod *Acartia tonsa*  
 1297 in turbulent environments. Marine Ecology Progress Series, 122: 147–158.
- 1298 Salama, N. K. G., Murray, A. G., and Rabe, B. 2011. Modelling dispersal of salmon lice in a  
 1299 large fjordic system: Loch Linnhe, Scotland. In 19th International Congress on  
 1300 Modelling and Simulation (pp. 12–16). Perth, Australia.
- 1301 Samsing, F., Oppendal, F., Dalvin, S., Johnsen, I., Vagseth, T., and Dempster, T. 2016. Salmon  
 1302 lice (*Lepeophtheirus salmonis*) development times, body size and reproductive outputs  
 1303 follow universal models of temperature dependence. Canadian Journal of Fisheries and  
 1304 Aquatic Science, 73:1841–1851.
- 1305 Sandvik, A., Bjørn, P., Ådlandsvik, B., Asplin, L., Skarðhamar, J., Johnsen, I., Myksvoll, M.,  
 1306 and Skogen, M. 2016. Toward a model-based prediction system for salmon lice  
 1307 infestation pressure. Aquaculture Environment Interactions, 8:527–542.
- 1308 Sandvik, A. D., Skarðhamar, J., Bjørn, P. A., Asplin, L., Llinares, S. R. M., and Johnsen, I. A.  
 1309 2014. The predictability of sea lice density in Norwegian Fjords. In Sea Lice 2014:  
 1310 Sustainable Solutions in Sea Lice Management. 31st August – 5th September,  
 1311 Portland, Maine. p. 46.
- 1312 Schram, T. A. 1993. Supplementary descriptions of the developmental stages of  
 1313 *Lepeophtheirus salmonis* (Kroyer, 1837) (Copepoda:Caligidae). In Pathogens of wild  
 1314 and farmed fish: Sea lice, 2nd edn, pp. 30–47. Ed. by G. A. Boxshall and D. Defaye.  
 1315 Ellis Horwood Ltd., New York. 378 pp.

1316 Skern-Mauritzen, R., Torrissen, O., and Glover, K. A. 2014. Pacific and Atlantic  
1317 *Lepeophtheirus salmonis* (Krøyer, 1838) are allopatric subspecies: *Lepeophtheirus*  
1318 *salmonis salmonis* and *L. salmonis oncorhynchi* subspecies novo. BMC Genetics, 15:  
1319 32.

1320 Skiftesvik, A. B., Bjelland, R., Durif, C. M. F., Johansen, I., and Browman, H. I. 2013.  
1321 Delousing of Atlantic Salmon (*Salmo salar*) by cultured vs. wild ballan wrasse (*Labrus*  
1322 *bergylta*) Aquaculture, 402: 113–118.

1323 Skugor, S., Glover, K. A., Nilsen, F., and Krasnov, A. 2008. Local and systemic gene  
1324 expression responses of Atlantic salmon (*Salmo salar* L.) to infection with the salmon  
1325 louse (*Lepeophtheirus salmonis*). BMC Genomics, 9: 1–18.

1326 Smith, G. W., Smith, I. P., and Armstrong, S. M. (1994). The relationship between river flow  
1327 and entry to the Aberdeenshire Dee by returning adult Atlantic salmon. Journal of Fish  
1328 Biology, 45: 953–960.

1329 Sommer, U. and Stibor, H. 2002. Copepoda - Cladocera - Tunicata: The role of three major  
1330 mesozooplankton groups in pelagic food webs. Ecological Research, 17: 161–174.

1331 Steele, J. H. and Frost, B. W. 1977. The structure of plankton communities. Proceedings of  
1332 the Royal Society B, 280: 485–534.

1333 Stucchi, D. J., Guo, M., Foreman, M. G. G., Czajko, P., Galbraith, M., Mackas, D. L., and  
1334 Gillibrand, P. A. 2011. Modeling sea lice production and concentrations in the  
1335 Broughton Archipelago, British Columbia. In Salmon lice: an integrated approach to  
1336 understanding parasite abundance and distribution, pp. 117–150. Ed. by S. Jones and  
1337 R. Beamish. Wiley-Blackwell, Oxford, UK. 336 pp.

1338 Suchman, C. L., and Sullivan, B. K. 2000. Effect of prey size on vulnerability of copepods to  
 1339 predation by the scyphomedusae *Aurelia aurita* and *Cyanea* sp. Journal of Plankton  
 1340 Research, 22: 2289–2306.

1341 Suchman, C. L., and Sullivan, B. K. 1998. Vulnerability of the copepod *Acartia tonsa* to  
 1342 predation by the scyphomedusa *Chrysaora quinquecirrha*: effect of prey size and  
 1343 behavior. Marine Biology, 132: 237–245.

1344 Sutherland, B. J., Covello, J. M., Friend, S. E., Poley, J. D., Koczka, K. W., Purcell, S. L.,  
 1345 MacLeod, T. L., Donovan, B. R., Pino, J., González-Vecino, J. L., Gonzalez, J.,  
 1346 Troncoso, J., Koop, B. F., Wadsworth, S. L., and Fast, M. D. 2017. Host parasite  
 1347 transcriptomics during immunostimulant-enhanced rejection of salmon lice  
 1348 (*Lepeophtheirus salmonis*) by Atlantic salmon (*Salmo salar*). FACETS, 2: 477–495.

1349 Tadiso, T. M., Krasnov, A., Skugor, S., Afanasyev, S., Hordvik, I., and Nilsen, F. 2011. Gene  
 1350 expression analyses of immune responses in Atlantic salmon during early stages of  
 1351 infection by salmon louse (*Lepeophtheirus salmonis*) revealed bi-phasic responses  
 1352 coinciding with the copepod-chalimus transition. BMC Genomics, 12: 141.

1353 Tiselius, P. and Jonsson, P. R. 1990. Foraging behaviour of six calanoid copepods:  
 1354 observations and hydrodynamic analysis. Marine Ecology Progress Series, 1–2: 23–  
 1355 33.

1356 Torres, G., Gime, L., and Anger, K. 2002. Effects of reduced salinity on the biochemical  
 1357 composition (lipid, protein) of zoea 1 decapod crustacean larvae. Journal of  
 1358 Experimental Marine Biology and Ecology, 277: 43–60.


- 1359 Tucker, C. S., Norman, R., Shinn, A. P., Bron, J. E., Sommerville, C., and Wootten, R. 2002.  
1360 A single cohort time delay model of the life-cycle of the salmon louse *Lepeophtheirus*  
1361 *salmonis* on Atlantic salmon *Salmo salar*. Fish Pathology, 37: 107–118.
- 1362 Tucker, C., Sommerville, C., and Wootten, R. 2000a. An investigation into the larval  
1363 energetics and settlement of the sea louse, *Lepeophtheirus salmonis*, an ectoparasite  
1364 copepod of Atlantic salmon, *Salmo salar*. Fish Pathology, 35: 137–143.
- 1365 Tucker, C. S., Sommerville, C., and Wootten, R. 2000b. The effect of temperature and salinity  
1366 on the settlement and survival of copepodids of *Lepeophtheirus salmonis* (Krøyer ,  
1367 1837) on Atlantic salmon, *Salmo salar* L. Journal of Fish Diseases, 23: 309–320.
- 1368 Tully, O. 1992. Predicting infestation parameters and impacts of caligid copepods in wild and  
1369 cultured fish populations. Invertebrate Reproduction and Development, 22: 91–102.
- 1370 Tully, O. 1989. The succession of generations and growth of the caligid copepods *Caligus*  
1371 *Elongatus* and *Lepeophtheirus Salmonis* parasitising farmed atlantic salmon smolts  
1372 (*Salmo Salar* L.). Journal of the Marine Biological Association UK, 69: 279–287.
- 1373 Tully, O. and Whelan, K. F. 1993. Production of nauplii of *Lepeophtheirus salmonis* (Kroyer)  
1374 (Copepoda: Caligidae) from farmed and wild salmon and its relation to the infestation  
1375 of wild sea trout (*Salmo trutta* L.) off the west coast of Ireland in 1991. Fisheries  
1376 Research, 17: 187–200.
- 1377 Ugelvik, M. S., Skorpning, A., and Mennerat, A. 2017. Parasite fecundity decreases with  
1378 increasing parasite load in the salmon louse *Lepeophtheirus salmonis* infecting  
1379 Atlantic salmon *Salmo salar*. Journal of Fish Diseases, 40: 671–678.
- 1380 Visser, A. W. and Fiksen, Ø. 2013. Optimal foraging in marine ecosystem models: selectivity,  
1381 profitability and switching. Marine Ecology Progress Series, 473: 91–101.

- 1382 Visser, A. W., Mariani, P., and Pigolotti, S. 2009. Swimming in turbulence: zooplankton  
1383 fitness in terms of foraging efficiency and predation risk. *Journal of Plankton Research*,  
1384 31: 121–133.
- 1385 Williams, J. A. 1985. The role of photoperiod in the initiation of breeding and brood  
1386 development in the amphipod *Talitrus saltator* Montagu. *Journal of Experimental*  
1387 *Marine Biology and Ecology*, 86: 59–72.
- 1388 Wirtz, K. W. 2014. A biomechanical and optimality-based derivation of prey-size  
1389 dependencies in planktonic prey selection and ingestion rates. *Marine Ecology*  
1390 *Progress Series*, 507: 81–94.
- 1391 Wirtz, K. W. 2012. Who is eating whom? Morphology and feeding type determine the size  
1392 relation between planktonic predators and their ideal prey. *Marine Ecology Progress*  
1393 *Series*, 445: 1–12.
- 1394 Wirtz, K. W. 2011. Non-uniform scaling in phytoplankton growth rate due to intracellular  
1395 light and CO<sub>2</sub> decline. *Journal of Plankton Research*, 33: 1325–1341.
- 1396 Woodward, G., Speirs, D. C., and Hildrew, A. G. 2005. Food Webs: From Connectivity to  
1397 Energetics. *Advances in Ecological Research*, 36: 85–135.
- 1398 Wootten, R., Smith, J. W., and Needham, E. A. 1982. Aspects of the biology of the parasitic  
1399 copepods *Lepeophtheirus salmonis* and *Caligus elongatus* on farmed salmonids, and  
1400 their treatment. *Proceedings of the Royal Society of Edinburgh B*, 81: 185–197.
- 1401 Young, S. and Watt, P. 1993. Behavioral mechanisms controlling vertical migration in  
1402 *Daphnia*. *Limnology and Oceanography*, 38: 70–79.

1403 Zarowiecki, M. and Berriman, M. (2015). What helminth genomes have taught us about  
1404 parasite evolution. *Parasitology*, 142(S1): S85–S97.

1405

1406    **9 List of Figures**



1407


1408    **Figure 1.** Relationship between water temperature and the number of eggs per egg string in  
1409    *Lepeophtheirus salmonis* from salmon farms on the West Coast of Scotland. Redrawn from  
1410    Ritchie *et al.*, 1993

1411


1412

1413





**Figure 2.** Typical ingestion (light grey area) and selection (dark grey area) feeding kernels for (a) narrow-range, selective feeders, *e.g.* copepods, and (b) broad-range, unselective feeders, *e.g.* jellyfish, where prey are abundant. Adapted and redrawn from Wirtz (2014)



**Figure 3.** A conceptual model of the stages of the sea louse life cycle that determine the number of copepodids available for infection and their infection success with factors that may affect survival/infectivity at each stage. Open arrows show the life cycle and black arrows show the factors that may affect each stage of the life cycle.

1425 **10 List of Tables**

1426 **Table 1.** Key values of fecundity in *L. salmonis* (means  $\pm$  SD). References: (a) Heuch *et al.*, 2000, (b) Johnson & Albright, 1991b, (c) Gravid,  
1427 1996, (d) Pike & Wadsworth, 1999, nd = no data available.

|                                   | <i>L. salmonis salmonis</i> |                   |  | <i>L. salmonis oncorhynchi</i> |                  |                               |
|-----------------------------------|-----------------------------|-------------------|--|--------------------------------|------------------|-------------------------------|
|                                   | Time (d)                    | Egg string pairs  | No. of eggs  | Time (d)                       | Egg string pairs | No. of eggs                   |
| <i>Egg string production rate</i> |                             |                   |  |                                |                  |                               |
| 7.2 °C                            | 11 <sup>a</sup>             | -                 | -  | nd                             | -                | -                             |
| 12.2 °C                           | 5 <sup>a</sup>              | -                 | -  | nd                             | -                | -                             |
| <i>Production capacity</i>        | -                           | 4.95 <sup>a</sup> | -  | -                              | nd               | -                             |
| <i>No. egg strings per string</i> |                             |                   |  |                                |                  |                               |
| 7.2 °C                            | -                           | -                 | 1st string 152<br>subsequent strings<br>285 <sup>a</sup> | -                              | -                | nd                            |
| 10 °C                             | -                           | -                 | nd   | -                              | -                | 344.6 $\pm$ 79.8 <sup>b</sup> |
| 1 <sup>st</sup> string            | -                           | -                 | 141.09 $\pm$ 22.19 <sup>c</sup>                          | -                              | -                | nd                            |
| 2 <sup>nd</sup> string            | -                           | -                 | 216.4 $\pm$ 67.59 <sup>c</sup>                           | -                              | -                | nd                            |
| 3 <sup>rd</sup> string            | -                           | -                 | 208.2 $\pm$ 50.97 <sup>c</sup>                           | -                              | -                | nd                            |
| wild lice                         | -                           | -                 | 965 $\pm$ 30.1 <sup>d</sup>                              | -                              | -                | nd                            |
| farmed untreated lice             | -                           | -                 | 758 $\pm$ 39.4 <sup>d</sup>                              | -                              | -                | nd                            |
| farmed treated lice               | -                           | -                 | 297 $\pm$ 19.1 <sup>d</sup>                              | -                              | -                | nd                            |

1428

**Table 2.** Key values of hatching in *L. salmonis*. (Means  $\pm$  SD, parentheses indicate ranges).  
References: (a) Gravid, 1996, (b) Heuch *et al.*, 2000, (c) Johnson & Albright, 1991b, (d) Samsing *et al.*, 2016, nd = no data available.

|                                   | <i>L. salmonis salmonis</i> |                  | <i>L. salmonis oncorhynchi</i> |                            |
|-----------------------------------|-----------------------------|------------------|--------------------------------|----------------------------|
|                                   | Proportion                  | Time (h)         | Proportion                     | Time (h)                   |
| <i>Non-viable egg strings</i>     | 2.41% <sup>a</sup>          | -                | nd                             | -                          |
| <i>Non-viable eggs per string</i> | 17.66% <sup>a</sup>         | -                | nd                             | -                          |
| 7.2 °C                            | 13.3% <sup>b</sup>          | -                | nd                             | -                          |
| 12.2 °C                           | 7.5% <sup>b</sup>           | -                | nd                             | -                          |
| <i>Hatching period</i>            |                             |                  |                                |                            |
| 5 °C                              | -                           | 240 <sup>a</sup> | -                              | nd                         |
| 7 °C                              | -                           | 192 <sup>a</sup> | -                              | nd                         |
| 10 °C                             | -                           | 144 <sup>a</sup> | -                              | 31.7 $\pm$ 17 <sup>c</sup> |
| <i>Hatching success at 10 °C</i>  |                             |                  |                                |                            |
| 0 ppt                             | 3.27% <sup>a</sup>          | -                | nd                             | -                          |
| 10 ppt                            | nd                          | -                | 0% <sup>c</sup>                | -                          |
| 15 ppt                            | nd                          | -                | 70% <sup>c</sup>               | -                          |
| 20 ppt                            | nd                          | -                | 78% <sup>c</sup>               | -                          |
| 25 ppt                            | nd                          | -                | 100% <sup>c</sup>              | -                          |
| 30 ppt                            | 86.36% <sup>a</sup>         | -                | nd                             | -                          |
| <i>Hatching success at 34 ppt</i> |                             |                  |                                |                            |
| 3 °C                              | 28 $\pm$ 4% <sup>d</sup>    | -                | nd                             | -                          |
| 5 °C                              | 85 $\pm$ 4% <sup>d</sup>    | -                | nd                             | -                          |
| 7 °C                              | 90 $\pm$ 4% <sup>d</sup>    | -                | nd                             | -                          |
| 10 °C                             | 87 $\pm$ 3% <sup>d</sup>    | -                | nd                             | -                          |
| 15 °C                             | 100% <sup>d</sup>           | -                | nd                             | -                          |
| 20 °C                             | 100% <sup>d</sup>           | -                | nd                             | -                          |
| <i>Viability of nauplii</i>       |                             |                  |                                |                            |
| 20 ppt                            | nd                          | -                | 19.8% (0–89.9)                 | -                          |
| 25 ppt                            | nd                          | -                | 51.1% (12–94.1)                | -                          |
| 30 ppt                            | nd                          | -                | 65.9% (9.7–95)                 | -                          |

**Table 3.** Key stage timings for *L. salmonis* (mean values). References: (a) Johnson & Albright, 1991b, (b) Johannessen, 1977, (c) Wootten *et al.*, 1982, (d) Gravid, 1996, (e) Boxaspen & Næss, 2000, (f) Samsing *et al.*, 2016, nd = no data available.

|  | <i>L. salmonis salmonis</i>                        |                    | <i>L. salmonis oncorhynchi</i> |                    |
|--|--|--------------------|--------------------------------|--------------------|
|  | Time (d)   | Time (h)           | Time (d)                       | Time (h)           |
| <i>Egg development time</i>              |  |                    |                                |                    |
| 2 °C                                     | 45.1 ± 0.5 <sup>e</sup>                            | -                  | nd                             | -                  |
| 3 °C                                     | 35.2 ± 0.4 <sup>e</sup><br>20.8 ± 1.5 <sup>f</sup> | -                  | nd                             | -                  |
| 4 °C                                     | 27.6 ± 0.2 <sup>e</sup>                            | -                  | nd                             | -                  |
| 5 °C                                     | 21.6 ± 0.1 <sup>e</sup><br>13.0 ± 7.8 <sup>f</sup> | -                  | 17.5 <sup>a</sup>              | -                  |
| 9 °C                                     | 33–39 <sup>b</sup>                                 | -                  | nd                             | -                  |
| 9.5 °C                                   | 25 <sup>b</sup>                                    | -                  | nd                             | -                  |
| 10 °C                                    | 8.7 ± 0.1 <sup>e</sup><br>4.6 ± 1.3 <sup>f</sup>   | -                  | 8.6 <sup>a</sup>               | -                  |
| 11.5 °C                                  | 10–14 <sup>b</sup>                                 | -                  | nd                             | -                  |
| 15 °C                                    | 2.88 ± 1.0 <sup>f</sup>                            | -                  | 5.5 <sup>a</sup>               | -                  |
| 20 °C                                    | 1.8 ± 0.5 <sup>f</sup>                             | -                  | nd                             | -                  |
| <i>Duration of first nauplius stage</i>  |  |                    |                                |                    |
| 5 °C                                     | -  | nd                 | -                              | 52 <sup>a</sup>    |
| 7.5 °C                                   | -  | 43.25 <sup>d</sup> | -                              | nd                 |
| 9.2 °C                                   | -  | 35 <sup>b</sup>    | -                              | nd                 |
| 10 °C                                    | -  | nd                 | -                              | 30.5 <sup>a</sup>  |
| 12 °C                                    | -  | 18 <sup>c</sup>    | -                              | nd                 |
| 15 °C                                    | -  | nd                 | -                              | 9.2 <sup>a</sup>   |
| 15.5 °C                                  | -  | 12 <sup>b</sup>    | -                              | nd                 |
| <i>Duration of second nauplius stage</i> |  |                    |                                |                    |
| 5 °C                                     | -  | nd                 | -                              | 170.3 <sup>a</sup> |
| 9.2 °C                                   | -  | 77 <sup>b</sup>    | -                              | nd                 |
| 10 °C                                    | -  | nd                 | -                              | 56.9 <sup>a</sup>  |
| 11 °C                                    | -  | 63 <sup>bc</sup>   | -                              | nd                 |
| 12 °C                                    | -  | 46 <sup>c</sup>    | -                              | nd                 |
| 15 °C                                    | -  | nd                 | -                              | 35.6 <sup>a</sup>  |
| 19 °C                                    | -  | 33 <sup>c</sup>    | -                              | nd                 |
| <i>Development time to copepodid</i>     |  |                    |                                |                    |
| 2 °C                                     | -  | 1644 <sup>e</sup>  | -                              | nd                 |
| 5 °C                                     | -  | 276 <sup>f</sup>   | -                              | nd                 |

|       |   |                        |   |    |
|-------|---|------------------------|---|----|
| 7 °C  | - | 168 <sup>f</sup>       | - | nd |
| 10 °C | - | 111–177.5 <sup>d</sup> | - | nd |
|       |   | 305 <sup>e</sup>       |   |    |
|       |   | 108 <sup>f</sup>       |   |    |
| 15 °C | - | 36 <sup>f</sup>        | - | nd |
| 20 °C | - | 48 <sup>f</sup>        | - | nd |

1436

1437 **Table 4.** Key values of survival for *L. salmonis* larvae (50 % survival times (LT<sub>50</sub>) are shown  
1438 unless specified otherwise). References: (a) Gravid, 1996, (b) Johnson & Albright, 1991b, (c)  
1439 Bricknell *et al.*, 2006, (d) Wootten *et al.*, 1982, (e) Samsing *et al.*, 2016, nd = no data available.

|   | <i>L. salmonis salmonis</i> |                     |                 | <i>L. salmonis oncorhynchi</i> |                    |                       |
|---|-----------------------------|---------------------|-----------------|--------------------------------|--------------------|-----------------------|
|   | width(μm)                   | Proportion          | Time (h)        | width (μm)                     | Proportion         | Time (h)              |
| <i>Nauplius I width</i>                 |                             |                     |                 |                                |                    |                       |
| May                                     | 187.76 <sup>a</sup>         | -                   | -               | nd                             | -                  | -                     |
| August                                  | 214.05 <sup>a</sup>         | -                   | -               | nd                             | -                  | -                     |
| <i>Nauplius I lipid reserve width</i>   |                             |                     |                 |                                |                    |                       |
| May                                     | 112.98 <sup>a</sup>         | -                   | -               | nd                             | -                  | -                     |
| August                                  | 135.84 <sup>a</sup>         | -                   | -               | nd                             | -                  | -                     |
| <i>Survival to copepodid at 10 °C</i>   |                             |                     |                 |                                |                    |                       |
| <25 ppt                                 | -                           | 0% <sup>a</sup>     | -               | -                              | nd                 | -                     |
| <30 ppt                                 | -                           | nd                  | -               | -                              | 0% <sup>b</sup>    | -                     |
| 30 ppt                                  | -                           | nd                  | -               | -                              | 35.2% <sup>b</sup> | -                     |
| 35 ppt                                  | -                           | 18.33% <sup>a</sup> | -               | -                              | nd                 | -                     |
| <i>Copepodid survival time at 10 °C</i> |                             |                     |                 |                                |                    |                       |
| 0-10 ppt                                | -                           | -                   | 15 <sup>a</sup> | -                              | -                  | nd                    |
| 5 ppt                                   | -                           | -                   | nd              | -                              | -                  | 3 <sup>b</sup>        |
| 10 ppt                                  | -                           | -                   | nd              | -                              | -                  | <24 <sup>b</sup>      |
| 15 ppt                                  | -                           | -                   | 54 <sup>a</sup> | -                              | -                  | nd                    |
| 16 ppt                                  | -                           | -                   | 4 <sup>c</sup>  | -                              | -                  | nd                    |
| 19 ppt                                  | -                           | -                   | 6 <sup>c</sup>  | -                              | -                  | nd                    |
| 20 ppt                                  | -                           | -                   | 67 <sup>a</sup> | -                              | -                  | nd                    |
| 23 ppt                                  | -                           | -                   | 8 <sup>c</sup>  | -                              | -                  | nd                    |
| 25 ppt                                  | -                           | -                   | 68 <sup>a</sup> | -                              | -                  | max. 17d <sup>b</sup> |
| 26 ppt                                  | -                           | -                   | 11 <sup>c</sup> | -                              | -                  | nd                    |
| 29 ppt                                  | -                           | -                   | 24 <sup>c</sup> | -                              | -                  | nd                    |

|  |   |   |  |   |   |    |
|--|---|---|--|---|---|----|
| 30 ppt                                   | - | - | 55 <sup>a</sup>  | - | - | nd |
| 33 ppt                                   | - | - | 22 <sup>c</sup>  | - | - | nd |
| 35 ppt                                   | - | - | 64 (max. 18d) <sup>a</sup>                               | - | - | nd |
| 36 ppt                                   | - | - | 25 <sup>c</sup>  | - | - | nd |
| <i>Copepodid survival time at 35 ppt</i> |   |   |  |   |   |    |
| 5 °C                                     | - | - | 116 <sup>a</sup><br>240 (LT <sub>80</sub> ) <sup>e</sup> | - | - | nd |
| 7 °C                                     | - | - | 300 (LT <sub>80</sub> ) <sup>c</sup>                     | - | - | nd |
| 10 °C                                    | - | - | 90 <sup>a</sup><br>312 (LT <sub>80</sub> ) <sup>c</sup>  | - | - | nd |
| 12 °C                                    | - | - | 96 <sup>d</sup>  | - | - | nd |
| 15 °C                                    | - | - | 82 <sup>a</sup><br>228 (LT <sub>80</sub> ) <sup>e</sup>  | - | - | nd |
| 20 °C                                    | - | - | 144 (LT <sub>80</sub> ) <sup>e</sup>                     | - | - | nd |

1440

1441 **Table 5.** Key variables of infectivity in *L. salmonis salmonis* larvae. References: (a) Cook *et*  
1442 *al.*, 2010, (b) Bron, 1993, (c) Gravid, 1996, (d) Tucker *et al.*, 2002, (e) Samsing *et al.*, 2016, (f)  
1443 Bricknell *et al.*, 2006. No infectivity data is available for *L. salmonis oncorhynchi*.

|  | Infectivity capability       | Lipid reserves             | Proportion                 |
|--|------------------------------|----------------------------|----------------------------|
| <i>Copepodid age</i>                         |                              |                            |                            |
| 7–10d  | Increasing <sup>abcd</sup>   | Good <sup>abcd</sup>       | -                          |
| 11–15d                                       | Mature <sup>abcd</sup>       | Decreasing <sup>abcd</sup> | -                          |
| 16–20d                                       | Less capable <sup>abcd</sup> | Low <sup>abcd</sup>        | -                          |
| <i>Infection success at 10 °C and 35 ppt</i> |                              |                            |                            |
| 1-day-old copepodids                         | -                            | -                          | 22.22 ± 8.32% <sup>c</sup> |
| 7-day-old copepodids                         | -                            | -                          | 14 ± 8.71% <sup>c</sup>    |
| <i>Infection success aged 1–3 d</i>          |                              |                            |                            |
| Illumination                                 | -                            | -                          | 23.2% <sup>b</sup>         |
| No illumination                              | -                            | -                          | 18.4% <sup>b</sup>         |
| <i>Infection success at 35 ppt</i>           |                              |                            |                            |
| 5 °C   | -                            | -                          | 2.1 ± 0.4% <sup>e</sup>    |
| 6.5 °C                                       | -                            | -                          | 20% <sup>d</sup>           |
| 10 °C  | -                            | -                          | 53.2 ± 2.3% <sup>e</sup>   |
| 11 °C  | -                            | -                          | 75% <sup>d</sup>           |
| 20 °C  | -                            | -                          | 41.6 ± 2.0% <sup>e</sup>   |

*Infection success at 12 °C*

|        |   |   |                  |
|--------|---|---|------------------|
| 12 ppt | - | - | 1% <sup>f</sup>  |
| 19 ppt | - | - | 10% <sup>f</sup> |
| 26 ppt | - | - | 14% <sup>f</sup> |
| 34 ppt | - | - | 31% <sup>f</sup> |

**Table 6.** A summary table of parameters influencing the production, timing and survival of sea lice larvae and observable biotic and abiotic factors that may influence them. Cells marked with an X represent areas where some data already exist and blank cells represent areas of data deficiency. References: (a) Heuch *et al.*, 2000, (b) Tully & Whelan, 1993, (c) Gravid, 1996, (d) Ritchie *et al.*, 1993, (e) Johnson & Albright, 1991a, (f) Tully, 1992, (g) Samsing *et al.*, 2016, (h) Johnson & Albright, 1991b, (i) Johannessen, 1977, (j) Boxaspen & Næss, 2000, (k) Wootten *et al.*, 1982.

| Parameter                         | Origin:<br>wild /<br>farmed | Variable factor |          |                        | Season | Reference        |
|-----------------------------------|-----------------------------|-----------------|----------|------------------------|--------|------------------|
|                                   |                             | Temp.           | salinity | Light /<br>photoperiod |        |                  |
| Female size                       | X                           | X               |          | X                      |        | a, b, c          |
| Egg string<br>production rate     |                             | X               |          |                        |        | a                |
| No. of eggs                       | X                           | X               |          | X                      |        | a, c, d, e, f, g |
| Egg development<br>time           |                             | X               |          |                        |        | h, i, j          |
| Egg development<br>time           |                             | X               |          |                        |        | g                |
| Hatching period                   |                             | X               | X        |                        |        | c, h             |
| Egg viability                     |                             | X               | X        | X                      |        | a, c, h          |
| Hatching success                  |                             | X               | X        | X                      |        | c, g, h          |
| Nauplius I<br>development time    |                             | X               |          |                        |        | c, g, h, i, j, k |
| Nauplius II<br>development time   |                             | X               |          |                        |        | g, h, i, j, k    |
| Nauplius I width                  |                             |                 |          |                        | X      | c                |
| Nauplius I lipid<br>reserve width |                             |                 |          |                        | X      | c                |
| Survival to<br>copepodid          |                             | X               | X        |                        |        | c, i             |



Copepodid survival  
time

X

X

c, h, i

---

1452