

1 This is the peer reviewed version of the following article: Lyons, P. P., Turnbull, J.  
2 F., Dawson, K. A. and Crumlish, M. (2017), Effects of low-level dietary  
3 microalgae supplementation on the distal intestinal microbiome of farmed  
4 rainbow trout *Oncorhynchus mykiss* (Walbaum). *Aquaculture Research*, 48: 2438–  
5 2452, which has been published in final form at <https://doi.org/10.1111/are.13080>. This  
6 article may be used for non-commercial purposes in accordance With Wiley Terms and  
7 Conditions for self-archiving.

**Effects of low level dietary microalgae supplementation on the distal intestinal microbiome of farmed rainbow trout *Oncorhynchus mykiss***

Philip P. Lyons<sup>ab\*</sup>, James F. Turnbull<sup>a</sup> Karl A. Dawson<sup>b</sup> and Margaret Crumlish<sup>a</sup>

<sup>a</sup> *Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, United Kingdom*

<sup>b</sup> *Alltech Biotechnology Inc., 3031 Catnip Hill Pike, Nicholasville KY 40356, USA*

**Abstract**

In this study, high throughput 16S rRNA sequencing was used to investigate the effect of a novel ingredient, dietary microalgae (*Schizochytrium limacinum*), on the distal intestinal microbiome of farmed rainbow trout *Oncorhynchus mykiss*. Dietary microalgae are rich in omega 3 polyunsaturated fatty acids, can be produced sustainably, and have been shown to have beneficial effects on host health. Microbial community profiles were compared between the distal intestinal contents of fish fed a control diet and a treatment diet that partially substituted dietary microalgae for the fish oil component (5% inclusion). The results of this research showed that the microbial communities of both fish populations were composed of similar microbial taxa, however the treatment group fed the microalgae supplement possessed a greater level of microbial diversity than those in the control group. A limited number of bacterial taxa were discriminatory between diets and were significantly elevated in the treatment group, notably operational taxonomic units (OTU's) assigned to the genera *Streptococcus*, *Leuconostoc*, *Lactobacillus*, *Lactococcus* and *Weissella*. However, the overall structure of the intestinal microbiome between control and treatment groups was not found to be significantly different. The treatment group displayed a heavier mean weight and condition factor at the end of the trial period. The results of this study suggest that microalgae can be used as a replacement for a proportion of fish oil in aquafeeds, with minor changes to the intestinal microbiome of farmed rainbow trout, and positive effects on growth.

**Keywords:** aquaculture, bacteria, intestine, microalgae, microbiome, rainbow trout, 16S rRNA sequencing

\*Corresponding author: P Lyons, Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, UK. E-mail: [p.p.lyons@stir.ac.uk](mailto:p.p.lyons@stir.ac.uk)

## 1. Introduction

Numerous studies have reported that diet type is a major driver in shaping the GI microflora of both terrestrial and aquatic animals (Ringo & Olsen 1999; Merrifield et al 2010; Merrifield, Dimitroglou, Foey, Davies, Baker, Bogwald, Castex & Ringo 2010; Merrifield et al 2011; Gatesoupe, Huelvan, Le Bayon, Sevre, Aaasen, Degnes, Mazurais, Panserat, Zambonino-Infante & Kaushik 2014; Kormas, Meziti, Mente. & Frentzos 2014; Miyake, Ngugi & Stingl 2015). A number of studies have investigated the impact of different diets on the intestinal microflora of farmed salmonids, mainly focusing on the impact of selected probiotics, prebiotics and immunostimulants included within the diet formulation. The vast majority of these investigations have been undertaken using culture-based and low resolution molecular microbiological techniques that provide an incomplete picture of the intestinal microbiome.

However, more recently, high throughput sequencing technologies have been used to examine the effect of diet on the intestinal microbiome of fish in far greater detail. Desai et al (2012) used 454 pyrosequencing to demonstrate reproducible effects on the intestinal microbiome of farmed rainbow trout fed soybean meal (SBM), noting changes in the ratio of Firmicutes: Proteobacteria as a result of supplementation. Ingerslev, Von Gersdorff, Jorgensen, Lenz Strube, Larsen, Dalsgaard, Boye & Madsen (2014) used the HiSeq® platform to demonstrate changes in the structure of the intestinal microflora of rainbow trout fry fed with marine versus plant-based dietary ingredients. In contrast to these results, Wong, Waldrop, Summerfelt, Davidson, Barrows, Kenney, Welch, Wiens, Snekvik, Rawls & Good (2013) reported that the intestinal microbiome of rainbow trout was largely unaffected by dietary alterations and resulted in only very minor changes to specific microbial community assemblages. Furthermore, substantial inter animal variation in the microbial community structure between individual fish has been reported (Mansfield, Desai, Nilson, Van Kessel, Drew and Hill 2010) suggesting that analysis of pooled samples are not suitable in studies of the intestinal

microbiome. The use of pyrosequencing platforms play an important role in this regard, permitting high resolution analysis of individual gut microbiomes, leading to more reliable conclusions regarding the effect of dietary alterations on the structure of the microbial communities in the fish gastrointestinal (GI) tract.

It has been widely reported that the gut microflora of aquatic animals is responsible for the digestion of algal cells, the production of both amino acid and short-chain fatty acids, in addition to secreting inhibitory compounds that protect against colonization of the gut by bacterial pathogens (Austin 2006; Nayak 2010; Ghanbari, Kneifel & Domig 2015). Research concerning the impact of microalgae on the structure of the intestinal microbiome however is limited and has hitherto primarily focused on wild herbivorous fish species that consume algal substrates in their natural habitat (Choat & Clements 1998; Ward, Steven, Penn, Methe & Detrich 2009; Smriga, Sandin & Azam 2010) with only a single study examining farmed fish species (Cerezuela, Fumanal, Tapia-Paniagua, Meseguer, Morinigo & Esteban (2012). Conflicting results have been reported in these studies, with some reporting increases and others reporting decreases in microbial diversity as a result of dietary algal consumption. This suggests that whilst diet impacts the diversity of gut microflora identified in fish, the relationship between novel dietary components such as microalgae, and the structure of the intestinal microbiome, is not clear and thus further detailed examination is undoubtedly required.

The primary objective of this study was to characterize the intestinal microbiome of farmed rainbow trout fed both a standard control diet, and a treatment diet containing a dietary microalgae supplement (5%), in order to test whether differences in diet composition lead to alterations in the structure of the microbial community. The aquafeed sector recognizes the need to provide dietary alternatives to fish oil which provide comparative health benefits to farmed fish species. Therefore, the secondary aim of this research was to test for any

differences in growth performance between the control and treatment groups and whether or not this could be correlated with the composition of the intestinal microbiome. It was hypothesized that feeding farmed rainbow trout slightly different diets would alter the structure of the intestinal microbiome in these fish.

## 2. Materials and methods

### 2.1 Dietary formulation

Two diets, one control and one treatment, were formulated at the Hellenic Centre for Marine Research (HCMR, Anavyssos Attiki, Greece). These diets were similar except that the experimental diet contained a 5% whole cell microalgae ingredient (*Schizochytrium limacinum*; Alltech Biotechnology, Nicholasville USA). The whole cell microalgae principally replaced the fish oil component in the treatment diet (3%), but also partially replaced soybean concentrate and wheat meal ingredients, each at a level of 1% (Table S1). Both diets met or exceeded the guideline nutrient requirements for rainbow trout (National Research Council 2011).

### 2.2 Experimental design and sampling protocol

Farmed rainbow trout (*O. mykiss*) were obtained from a local trout farm and transferred to the Aquatic Research Facility (ARF) at the University of Stirling Institute of Aquaculture (Stirling, UK). The average weight of the fish on arrival at the ARF was  $31.7 \pm 2.6$  g. Fish were quarantined in a communal tank for 10 days, prior to random allocation into twelve 100 L tanks ( $n=25$  tank<sup>-1</sup>) maintained on a flow through system, under a 12h light and 12h dark cycle and an ambient water temperature ( $14 \pm 1^\circ\text{C}$ ). All instructions and guidelines set by the UK Home Office under the Animal Welfare Act of 1986 were adhered to throughout this experimental trial. Each tank was randomly allocated the diets, giving four replicates per treatment, and each group was hand fed a ration of approximately 2% of their body weight twice daily.

At the end of the 15 week trial period, a total of three fish from each of four replicates per treatment were randomly removed for sampling. Fish were sacrificed with a lethal dose of anaesthetic benzocaine (Sigma Aldrich®) and swabbed with 100% ethanol before dissection through the ventral surface. The tissues surrounding the visceral fat were removed and the

distal gut contents (~150 mg) were aseptically collected by gently squeezing the tissue with a sterile forceps, and placed into a sterile 2 ml capped microtube (Alpha laboratories<sup>®</sup>) containing 1 ml of lysis buffer (Qiagen). The gut was then incised and washed with a sterile 0.85% (w/v) NaCl solution, and the intestinal mucous was carefully removed from the gut wall. This material was placed into the same tube as the gut contents. All tubes were placed on dry ice before DNA extraction later the same day, in order to ensure optimal sample integrity. In addition to the intestinal samples, three pellets from each diet and a sample of the tank biofilm were also processed as described above, to compare the microbial communities of both the diets themselves and of the tank biofilm, with the intestinal microbiome of the trout.

### *2.3 Growth performance*

The length and weight of each fish sampled at the end of the trial period was recorded to measure growth performance, thermal growth coefficient (TGC) and condition factor (K). Final fish weight was measured as the mean final weight of each group  $\pm$  standard error of the mean (SEM). TGC was calculated using the formula  $TGC = (W_2^{(1/3)} - W_1^{(1/3)})/D^{(0)} \times 1000$  where  $W_2$  and  $W_1$  are weight at the end and at the start of the trial respectively, and  $d^0$  represents degree days. K was calculated using Fulton's equation  $K = (10^5 \times \text{weight})/\text{Length}^3$ .

### *2.4 DNA extraction*

A total of 150 mg of intestinal content material from each individual fish suspended in 1 ml of buffer ASL (Qiagen) was processed for DNA extraction. A further sample containing only 1 ml of buffer ASL was processed as a negative control. Samples were firstly disrupted using a Mini bead-beater 16 (Biospec Ltd.) at maximum speed for four separate cycles of 35 s each. Samples were allowed to settle, and total genomic DNA was extracted and purified using the QIAamp DNA Stool Mini Kit (Qiagen, Hilden, Germany), with the following modifications to the manufacturer's protocol: 150 mg starting material in 1ml buffer ASL; suspension heated at



95°C for 10 min to improve lysis of Gram positive bacteria; 0.5 Inhibitex tablet per sample in 700 µl supernatant; final sample elution volume of 50 µl. After extraction, the DNA concentration of all samples was determined both spectrophotometrically (Thermo Scientific NanoDrop™ 1000, DE, USA) and fluorometrically (Qubit® Life Technologies) to ensure optimal DNA purity, and stored at -20°C for subsequent processing.

## *2.5 16S rRNA PCR and pyrosequencing*

A PCR was first carried out using universal eubacterial primers 27F (AGAGTTTGATCMTGGCTAG) and 1492R (TACGGYTACCTTGTTACGACTT) (Weisburg et al 1991) that target the full length bacterial 16S rRNA gene sequence, to confirm the presence of ample microbial community DNA and to rule out the presence of any potential inhibitory compounds. The extraction from buffer ASL was included in the PCR run to check for the presence of microbial DNA in the reagent itself. The PCR conditions for this confirmatory reaction were as follows; denaturation at 95°C for 5 min, followed by 30 cycles of denaturation at 94°C for 2 min, annealing at 50°C for 1 min and elongation at 72°C for 2 min; before final elongation at 72°C for 10 min. Products were then visualized on a 1.5% (w/v) agarose gel, run at 100V for approximately 1 h 15 min. The presence of a single strong PCR product of 1500bp was considered to be indicative of the presence of microbial community DNA.

Illumina libraries were prepared following the method described by Caporaso, Lauber, Walters, Berg-Lyons, Huntley, Fierer, Owens, Betley, Fraser, Bauer, Gormley, Gilbert, Smith, & Knight (2012) using the NEXTflex 16S Amplicon-Seq kit (Bio Scientific, Austin USA). A total of 50 ng of template DNA was used for each individual sample and the V4 hypervariable region of the bacterial 16S rRNA gene (length 292bp) was amplified using primers 515F (GTGCCAGCMGCCGCGGTAA) and 806R (GGACTACHVGGGTWTCTAAT) (GATC

Biotechnology Inc., Konstanz). The PCR conditions were as follows; initial denaturation at 95°C for 5 min ; 25 cycles of denaturation at 95°C for 30 s, annealing at 55°C for 30 s and extension at 72°C for 30 s; followed by a final extension step at 72°C for 5 min. All samples were amplified in triplicate and all products purified using Agencourt Ampure XP beads (Beckman Coulter Ltd.). The products of the first PCR served as template for a second PCR with the same conditions as the first, however the number of cycles was reduced to eight, and Illumina sequencing adapters were added to the primers in the reaction mix. Following amplification, PCR products were purified using Agencourt Ampure XP (Beckman Coulter) with a modified 1:1 volume of PCR product to Ampure XP beads. Purified amplicons were quantified with Qubit, pooled in equal concentration and the final quality of the pooled library was validated using a Bioanalyzer 2100 (Agilent Technologies, Waldbronn, Germany). The final library was sequenced using the Illumina MiSeq® NGS system at GATC Biotechnology (Konstanz, Germany).

## *2.6 Bioinformatics*

Demultiplexing was performed with Casava v. 1.8 (Illumina) and reads representing the PhiX or reads not matching indices were removed. The open-source software Mothur (Schloss 2009) was used to process sequences from the demultiplexed 16S rRNA gene libraries. Sequences were firstly merged using the make.contigs command. Reads containing ambiguous bases, homopolymer runs greater than 8 bases, and sequences of less than 150 base pairs in length were removed from the dataset. Remaining sequences were aligned against mothur's Silva reference database, after customizing the reference alignment to concentrate on the v4 region only (length = 292bp). Further denoising of the dataset was performed using mothur's pre clustering algorithm, allowing for up to two differences between sequences. This sorted sequences by abundance, ordering from most abundant to least and identified sequences within

two nucleotides of each other. If sequences met these conditions they were merged. Chimeric sequences were then removed from the dataset using the UCHIME (Edgar, Haas, Clemente, Quince & Knight 2011) algorithm in mothur as a final denoising step prior to taxonomic classification.

For taxonomic analyses, sequences were annotated using the Bayesian classifier implemented by the ribosomal database project (RDP) Release 11 (Centre for Microbial Ecology, Michigan State University, East Lansing, MI, USA). A minimum confidence bootstrap threshold of 80% was required for each assignment, thus >80% of the classifications returned the same taxonomic assignment for a given read, after one thousand iterations. Sample coverage, rarefaction curves, bias-corrected Chao 1 richness and Simpson's index of diversity were calculated based on assembled OTU's using mothur. Samples were rarefied to the sample with the lowest number of sequences (sample AF6, n=314,961) before performing these diversity analyses, to ensure that any observed differences in diversity were not caused by uneven sampling depth.

## *2.7 Statistical analyses*

A student's t-test was performed to compare the growth performance data between control and treatment groups, and differences were considered significant at  $p < 0.05$ . The similarity of the structure and membership of the microbial communities found in each of the samples was calculated by creating a distance matrix based on the thetaYC (Yue & Clayton 2005) coefficient using the dist.seqs algorithm in mothur. This distance matrix was visualized using principal coordinate analysis, which allowed the intestinal microbial community profiles from the control and treatment groups to be compared. In addition, a dendrogram was created to further describe the similarity of the samples to each other (data not shown). Parsimony (Schloss & Handelsman 2006) and UniFrac (Lozupone and Knight 2005) analyses were

performed to determine whether any observed community structure clustering between diets was statistically significant. Finally, metatstats (White, Nagarajan & Pop 2009), LEfSe (Segata, Izard, Waldron, Gevers, Miropolsky, Garrett & Huttenhower 2011) and Indicator (McCune, Grace & Urban 2002) analyses were performed within mothur, in order to determine whether there were any phylotypes that exhibited a statistically significant representation between the control and treatment samples, and results were considered as significant at two levels,  $p < 0.05$  and  $p < 0.01$ . The same statistical analyses were also used to compare feed pellet/biofilm samples with the intestinal samples from the control and treatment groups.

### 3. Results

#### 3.1 Growth performance

All fish consumed both diets readily and upon conclusion of the trial, the weighed individuals from the treatment group had a higher mean weight and condition factor than the control group. The final mean weight and condition factor ( $\pm$  SE) for the treatment group was  $136.6 \pm 12.1$ g and  $1.44 \pm 0.06$  whereas these values for the control group were  $116.5 \pm 9.3$ g and  $1.33 \pm 0.04$  respectively (Figure S1). A t-test was performed using the Minitab 15 statistical software to test for significant differences between the performance parameters for both groups, however no such differences were found ( $p = 0.107$ ).

#### 3.2 Sequence data and diversity analyses

After quality filtering of sequences, a total of 18,282,541 sequences remained for analysis, which grouped into a total of 660 OTU's. After subsampling to that of the library containing the least number of reads (sample AF6,  $n=314,961$ ), rarefaction curves generated in mothur showed a trend towards a greater level of microbial diversity in the treatment group with a greater number of overall OTU's being recorded (Figure S2). This trend was reflected in the inverse simpson and Chao1 diversity indices, with the three richest samples (AF7, AF4 and AF6) belonging to the treatment group (Table 1). A very high level of sequence coverage was achieved in the analysis, with all rarefaction curves reaching saturation and Good's coverage estimations reaching >99% for each sample, indicating that the vast majority of microbial phylotypes present were sampled in the analysis.

### 3.3 Microbial community composition and influence of diets

The overall microbial community composition was similar in both the control and treatment populations of fish. The distribution of OTU's at the phylum level of both the control and treatment libraries is illustrated below (Figure 1). The vast majority of reads were assigned to nine separate bacterial phyla, although an overall total of 13 phyla were recorded. Within these phyla, 13 microbial classes dominated (Figure 2). The mean number of OTU's classified to genus level observed in the control group was 99 (maximum of 177, minimum of 58), whereas in the treatment group the mean was 135 (maximum of 255, minimum of 77) (Table 1), reflecting the trend towards an increased level of microbial diversity in these fish. Considerable variability amongst individuals was noted.

The Tenericutes were the dominant phylum identified in the libraries recovered from both the control and treatment groups, with *Mycoplasma* being the most dominant genus observed in both groups. This suggests that the abundance of *Mycoplasma* was not affected by diet type. The remaining OTU's primarily belonged to the Firmicutes, Proteobacteria and Spirochaetes. OTU's assigned to Bacteroidetes, Actinobacteria, Deinococcus-Thermus, Candidate Division WPS-1 and Fusobacteria were detected at much lower levels of sequence abundances. Within the Firmicutes, the most frequently observed OTU's were *Acetanerobacterium*, *Weissella*, *Catelllicoccus*, *Streptococcus*, *Leuconostoc*, *Lactobacillus*, *Lactococcus*, *Ornithinibacillus* and *Sediminbaciullus*. *Acetanaerobacterium* represented the second most dominant OTU recorded overall, and was present in higher mean relative sequence abundances in the group fed the control diet. Sequences assigned to the Proteobacteria were observed more frequently in the treatment fish and the most dominant OTU's within this phylum belonged to the  $\gamma$  subclass, and in particular *Acinetobacter*, *Escherischia/Shigella*, *Enterobacter*, *Pseudomonas* and *Pantoea*. The  $\alpha$  and  $\beta$  subclasses were also represented and the dominant OTU's recorded from these classes were *Ahrensia* and *Sphingomonas* and *Delftia* and *Pelomonas* respectively. The

Spirochaetes were principally represented by the genus *Brevinema*, however *Sphaerochaeta* was also detected. This microbial class was most abundant in the treatment fish, with an overall mean sequence abundance of 3.1%, versus 0.7% in the control fish. Members of the class Bacteroidetes were infrequently recorded, and the dominant OTU's assigned to this class recorded in this study were *Flavobacterium* and *Cloacibacterium*. Similarly, OTU's assigned to the Fusobacteria were poorly represented within all libraries analysed, with *Fusobacterium* and *Cetobacterium* the principal genera detected in the sequence analysis. One of the most dominant OTU's observed in both control and treatment libraries was assigned to Candidate division WPS-1, an unclassified phylum, indicating that a large portion of the trout microbiome is still yet to be fully characterized.

Principal coordinate analyses, when visualized based on the thetaYC distance matrix comparing similarities in community structure, showed that samples were broadly indistinguishable according to diet, with the treatment and control samples clustering close together (Figure 3). This trend was examined using both the parsimony (Schloss & Handelsman 2006) and unweighted Unifrac (Lozupone & Knight 2005) analyses performed in mothur and confirmed that the microbial community structures were not significantly different between dietary treatments (ParsSig = 0.269, UWSig = 0.49). The community structure between the feed pellets and the intestinal microbiome were however significantly different when analysed statistically (ParsSig = 0.025, WSig = <0.001, UWSig = 0.004). The microbial community structure of the tank biofilm sample was also found to be significantly different from that of the trout intestinal microbiome samples (WSig = <0.001).

Although intestinal community structures were not statistically different between control and treatment fish, metastats (White et al 2009) analyses revealed that a number of OTU's were discriminatory according to dietary treatment and hence were differentially represented according to dietary regime (Table 2). These OTU's were *Leuconostoc* ( $p = 0.009$ ),

*Streptococcus* ( $p = 0.009$ ), *Weissella* ( $p = 0.048$ ), Candidate Division WPS-1 ( $p = 0.006$ ),  
*Lactobacillus* ( $p = 0.010$ ), *Enterobacter* ( $p = 0.034$ ), *Lactococcus* ( $p = 0.046$ ) and *Bacillus* ( $p$   
 $= 0.047$ ). Furthermore, sequences representing each of these OTU's were significantly more  
abundant in the treatment group (Figure 4). Both the LEfSe (Segata et al 2011) and Indicator  
(McCune et al 2002) statistical algorithms also confirmed the same phlotypes as  
discriminatory according to diet, with the exception of *Weissella*, where  $p > 0.05$  for both  
metrics. *Acetanaerobacterium* and *Brevinema* were also selected due to obvious differences in  
overall mean sequence abundances and because of their high prevalence in the sequence  
libraries, but these phlotypes were not found to be discriminatory according to diet (Figure  
S3).



#### 4. Discussion

The findings from this study suggest that 5% dietary microalgal supplementation altered levels of bacterial diversity and individual populations of microbes, but not the overall microbial community structure within the intestine of rainbow trout. No significant differences were recorded in growth and condition between control and treatment fish. These results improve our understanding of the interactions between the rainbow trout GI microbiome and novel dietary ingredients such as microalgae in aquaculture. Dietary supplements will undoubtedly continue to be included in future aquaculture feed formulations as the industry's supply of existing sources of fishmeal and fish oil decline. This research found that all of the individual rainbow trout analysed from both test groups possessed broadly similar intestinal microbial community compositions, after fifteen weeks of feeding. However, there were statistically significant differences in the representation of specific bacterial taxa between the control and treatment groups. Within the treatment group a trend towards an increase in microbial diversity was observed, however this pattern was not observed in all fish within this group and consequently was not statistically significant. Nonetheless, the pattern of increased microbial diversity could be indicative of the microbial community within the intestine of these fish responding to the availability of a different dietary ingredient, and perhaps an additional fermentable substrate in the form of the whole cell microalgal supplement.

It has previously been reported that gut microbial diversity increases from carnivorous to omnivorous to herbivorous fish species, a pattern similar to that observed in mammals (Ley, Hamady, Lozupone, Turnbaugh, Ramey, & Bircher 2008). The reason for this pattern is still poorly understood, but may be correlated with the length of the GI tract in each fish species and hence the overall transit time of food through the gut. In carnivorous fish with short digestive systems, such as rainbow trout, food travels quickly through the gut and hence less time is available for microbial fermentation of dietary ingredients. However, in omnivorous

and herbivorous fish, there is a much slower transit time of food through the convoluted GI tract, enabling a greater level of microbial fermentation to occur and precipitating an increase in microbial diversity. Smriga et al (2010) reported that the intestinal microbiome of the herbivorous whitecheek surgeonfish *Acanthurus nigricans*, whose primary diet consists of algae and detritus, exhibited a far greater level of microbial diversity than that of the strictly carnivorous red snapper *Lutjanus bohar*. Similarly the omnivorous yellowbelly rockcod *Notothenia coriiceps* was shown to possess a greater intestinal microbial diversity than the carnivorous blackfin icefish *Chaenocephalus aceratus* (Ward et al 2009). In this study, it is not unreasonable to posit that the changes in microbial diversity observed in the treatment group were indicative of the microbiome adapting to digesting whole cell microalgae and its constituent polysaccharides. Whilst gut transit time was not measured in this study, similar trends towards an increased microbial diversity in the intestine of trout fed plant-based diets have been recorded (Desai et al 2012; Green et al 2013). A high level of microbial diversity in the intestine has been advocated as being beneficial to host health in that it provides a wider range of potential responses to stressful situations or provides individual resilience to acceptance of different dietary ingredients (Backhed, Ley, Sonnenburg, Peterson & Gordon, 2005). The presence of a more diverse microbiome in the microalgae fed fish could therefore represent a reflection of the need for additional plasticity in the structure of the microbiome in these fish, in order to aid digestion and the breakdown of the microalgal meal included in their diet.

The Tenericutes were the dominant microbial phylum in the vast majority of samples, followed by the Firmicutes and Spirochaetes. Within the Tenericutes, the Mollicutes were the most prominent class, with *Mycoplasma* being the dominant genus. This microbe has previously been recorded in the intestinal tract of both marine and freshwater fish species (Kim, Brunt & Austin 2007; Moran, Turner & Clements 2005; Bano, Derae-Smith, Bennett, Vasquez &

Hollibaugh, 2007; Holben, Williams, Gilbert, Saarinen, Sarkilahti. & Apajalahti 2002; Suhanova, Dzyuba, Triboy, Nikiforova, Denikina & Belkova 2011; Xing, Hou, Yuan, Liu, Qu & Liu 2013, Carda-Diequez, Mira & Fouz 2014). More recent analyses employing high throughput sequencing have reported similar findings to those of the present study, in that the Mycoplasmataceae appear to dominate read libraries from the distal intestinal microbiome of Atlantic salmon (Green et al 2013, Zarkasi, Abell, Taylor, Neuman, Hatje, Tamplin, Katouli & Bowman, 2014) and rainbow trout (Lowrey, Woodhams, Tacchi & Salinas 2015; Ozorio, Kopecka-Pilarczyk, Peixoto, Lochmann, Santos, Santos, Weber, Calheiros, Ferrez-Arruda, Vaz-Pires & Goncalves 2015). *Mycoplasma* do not, however, appear to be significantly affected by diet composition, as they were present in all fish sampled in this trial, irrespective of treatment. Furthermore, large numbers of Tenericutes have been documented in the gut of other aquatic animals such as oysters (King, Judd, Kuske & Smith 2012) and in terrestrial animals such as pigs (Leser, Amenuvor, Jensen, Lindecrona, Boye & Moller 2002).

The genus *Mycoplasma* are Gram positive bacteria that are closely related to the Bacilli/Clostridium branch of the phylum Firmicutes. These fastidious microbes lack cell walls, have a fermentative metabolism, a high G-C content and possess a genome size (~580Kbp) that is amongst the smallest in self-replicating microorganisms. Owing to this extremely small genome, it is unlikely that they perform many complex metabolic functions within the fish intestine, and may primarily be obligate commensals within the gut ecosystem. However *Mycoplasma* have previously been reported to produce lactic acid and acetic acid as their major metabolites (Freundt & Razin 1958). It is thus also possible that the dominance of *Mycoplasma* in the intestine of trout is a result of a long established symbiosis in which this microbe benefits from easy access to a multitude of fermentable substrates (e.g. cytoplasmic secretions) and the fish benefits from the acetic acid and lactic acid metabolites produced as a result. Extreme

genome reduction in bacterial symbionts residing within terrestrial animal hosts is a well described phenomenon, which may also occur in rainbow trout.

Previous studies analysing the effect of dietary alterations on rainbow trout microflora have reported that whilst slight differences are often observed, these are somewhat negligible in terms of their effect on the 'core' microbial community, and the population structure between control and test populations are usually quite similar (Wong et al 2013 ; Zarkasi et al 2014). However, these authors did report subtle effects of the different diets on the relative abundance of select groups of bacterial taxa. Similarly, the principal coordinate analysis data obtained in this study provides evidence of a very minor effect of different diets on the structure of the microbial community within the intestine of rainbow trout, with only a limited number of taxonomic groups being significantly affected by dietary alteration. Furthermore, analysis of the microbial communities of the diets themselves showed that they were very similar in structure, but were significantly different from the fish intestinal samples. It thus appears to be unlikely that the observed differences in microbiota composition between control and treatment fish could be due to the microbiota structure of the dietary pellets. Therefore, it appears that diet composition only had a minor effect on the intestinal microbiome. Others have also reported that switching dietary regimes, including nutritional substitution, can alter microbial diversity, community membership and/or structure to varying degrees (Ringo and Olsen 1999; Ringo, Sperstad, Myklebust, Refstie, & Krogdahl 2006; Askarian, Zhou, Olsen, Sperstad & Ringo 2012; Sullam, Essinger, Lozupone, O'Connor, Rosen, Knight, Kilham & Russell 2012).

Statistical analyses revealed that *Streptococcus*, *Leuconostoc*, *Weissella*, *Lactobacillus*, *Candidate Division WPS-1* and *Lactococcus* were significantly discriminatory between diets in this study. Each of these genera, most of which are members of the lactic acid bacteria (LAB), were significantly elevated in the microalgae fed fish. LAB are frequently recorded in the intestines of fish, including rainbow trout, albeit at low levels of abundance (Merrifield,

Balcazar, Daniels, Zhou, Carnevali, Sun, Hoseinifar & Ringo 2014). More recent research on the effect of diet on the rainbow trout microbiome using deep sequencing platforms have found that this group appears to be amongst the most responsive to dietary alterations. Ingerslev et al (2014a, b) reported that *Streptococcus*, *Leuconostoc*, *Weissella* and *Lactobacillus* were responsive to dietary shifts, and were significantly elevated in the microbiome of trout fed high levels of plant-based ingredients. Similarly, both Desai et al (2012) and Wong et al (2013) reported that levels of *Lactobacillus*, *Streptococcus*, *Weissella*, *Clostridia* and *Staphylococcus* were discriminatory according to plant based and grain based diets respectively. The same microbial groups, with the exception of *Staphylococcus*, were discriminatory by diet in the present study, indicating the possible development of a distinct trend in the literature towards dietary influences on lactic acid bacterial populations in the rainbow trout intestine, in spite of their perceived rarity within this ecosystem.

These bacterial taxa are generally considered to be beneficial organisms associated with a healthy intestinal epithelium, and many of the genera recorded in this research have been tested elsewhere for their potential probiotic capabilities in rainbow trout aquaculture (Joborn, Olssen, Westerdahl, Conway & Kjelleberg 1997; Irianto & Austin 2002; Panigrahi, Kiron, Kobayashi, Puangkaew, Satoh & Sugita 2004; Kim & Austin 2006, 2008; Balcazar, de Blas, Ruiz-Zarzuela, Vendrell, Girones & Muzquiz, 2007; Vendrell, Balcazar, de Blas, Ruiz-Zarzuela, Girones & Muzquiz 2008; Balcazar, Vendrell, de Blas, Ruiz-Zarzuela & Muzquiz 2009; Merrifield et al 2010; Perez-Sanchez, Balcazar, Merrifield, Carnevali, Gioacchini, de Blas & Ruiz-Zarzuela 2011). LAB are hypothesized to improve the health of rainbow trout in aquaculture by enhancing feed conversion efficiency and conferring protection against pathogenic bacteria via mechanisms of competitive exclusion. In addition, the production of organic acids (e.g. acetic acid, lactic acid) and compounds such as bacteriocins and enzymes can further protect the intestinal epithelium and aid in the digestion of resistant dietary

ingredients (Nayak 2010). The LAB have however been observed to represent only a minor constituent of the fish intestinal microbiome and so potential methods of manipulating and enriching these populations are of great interest in improving intestinal health and consequently fish performance in aquaculture.

Overall, the results presented showed that the inclusion of dietary microalgae did not impair rainbow trout growth or negatively impact the distal intestinal microbiome. The dominance of *Mycoplasma* in the microbial libraries of all fish analysed suggests that this phylotype is well adapted to life in the rainbow trout intestine, and hence further research into its potential functional role is undoubtedly required. The altered microbial diversity observed in the microalgae fed fish suggested a flexibility in the intestinal microbiome of these fish which may represent a response to the breakdown and digestion of this novel dietary ingredient. Whilst the ‘global’ microbiome structure was similar in both groups, there were statistically significant differences noted in community membership, with distinct microbial groups observed to be discriminatory according to diet, particularly members of the LAB such as *Weissella*, *Streptococcus*, *Lactococcus*, *Lactobacillus* and *Leuconostoc*. This represents a further indication of a possible, albeit subtle, dietary effect of the microalgae on these populations. The potential manipulation of microbial communities through dietary supplementation may represent a promising method for improving gut health and hence nutrient utilization in farmed rainbow trout. Whilst the data presented is certainly supportive of the inclusion of microalgae in farmed rainbow trout diets, further work is required to clarify the optimal level of inclusion to beneficially manipulate the intestinal microflora of these fish.

## 5. Acknowledgements

This study was fully funded by Stirling University Institute of Aquaculture and Alltech Biotechnology inc. as part of their Margin of Excellence PhD Program. The authors would like to thank Mr. Niall Auchinachie for his technical assistance during the aquarium phase of this research.

## 6. Conflicts of interest

The authors declare no conflicts of interest

## 7. References

- Askarian, F., Zhou, Z., Olsen, R.E., Sperstad, S. and Ringo, E. (2012) Culturable autochthonous gut bacteria in Atlantic salmon (*Salmo salar* L.) fed diets with or without chitin. Characterization by 16S rRNA gene sequencing, ability to produce enzymes and in vitro growth inhibition of four fish pathogens. *Aquaculture*, **326–329**(0), pp. 1-8.
- Austin, B. (2006) The bacterial microflora of fish, revised. *TheScientificWorldJournal*, **6**, pp. 931-945.
- Backhed, F., Ley, R.E., Sonnenburg, J.L., Peterson, D.A. and Gordon, J.I. (2005) Host-bacterial mutualism in the human intestine. *Science (New York, N.Y.)*, **307**(5717), pp. 1915-1920.
- Baeverfjord, G. and Krogdahl, A. (1996) Development and regression of soybean induced enteritis in Atlantic salmon, *Salmo salar* L., distal intestine – A comparison with the intestines of fasted fish. *Journal of Fish Diseases* **19**(5), pp. 375-387
- Bakke-McKellep, A.M., Penn, M.H., Salas, P.M., Refstie, S., Sperstad, S., Landsverk, T., Ringo, E. and Krogdahl, A. (2007) Effects of dietary soyabean meal, inulin and oxytetracycline

538 on intestinal microbiota and epithelial cell stress, apoptosis and proliferation in the teleost  
539 Atlantic salmon (*Salmo salar* L.). *The British Journal of Nutrition*, **97**(4), pp. 699-713.

540 Balcazar, J.L., de Blas, I., Ruiz-Zarzuela, I., Vendrell, D., Girones, O. and Muzquiz, J.L. (2007)  
541 Enhancement of the immune response and protection induced by probiotic lactic acid bacteria  
542 against furunculosis in rainbow trout (*Oncorhynchus mykiss*). *FEMS Immunology and Medical*  
543 *Microbiology* **51**, 185-193

544 Balcazar, J.L., Vendrell, D., de Blas, I., Ruiz-Zarzuela, I. and Muzquiz, J.L. (2009) Effect of  
545 *Lactococcus lactis* CLFP 100 and *Leuconostoc mesenteroides* CLFP 196 on *Aeromonas*  
546 *salmonicida* infection in brown trout (*Salmo trutta*). *Journal of Molecular Microbiology and*  
547 *Biotechnology* **17**, 153-157

548 Bano, N., Derae Smith, A., Bennett, W., Vasquez, L. and Hollibaugh, J.T. (2007) Dominance  
549 of Mycoplasma in the guts of the Long-Jawed Mudsucker, *Gillichthys mirabilis*, from five  
550 California salt marshes. *Environmental Microbiology*, **9**(10), pp. 2636-2641.

551 Bernstein, A.M., Ding, E.L., Willett, W.C. and Rimm, E.B. (2012) A meta-analysis shows that  
552 docosahexaenoic acid from algal oil reduces serum triglycerides and increases HDL-  
553 cholesterol and LDL-cholesterol in persons with coronary heart disease. *Journal of Nutrition*  
554 **142**(1), 99-104

555 Bolnick, D.I., Snowberg, L.K., Hirsch, P.E., Lauber, C.L., Knight, R., Caporaso, J.G. and  
556 Svanback, R. (2014) Individuals' diet diversity influences gut microbial diversity in two  
557 freshwater fish (threespine stickleback and Eurasian perch). *Ecology Letters*, **17**(8), pp. 979-  
558 987.

559 Bradbury, J. (2011) Docosahexaenoic acid (DHA): an ancient nutrient for the modern human  
560 brain. *Nutrients* **3**(5), 529-554



561 Buddington, R.K., Krogh, A. and Bakke-McKellep, A.M. (1997) The intestines of  
 562 carnivorous fish: structure and functions and the relations with diet. *Acta physiologica*  
 563 *Scandinavica. Supplementum*, **638**, pp. 67-80.

564 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,  
 565 S.M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J.A., Smith, G. and Knight, R.  
 566 (2012) Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq  
 567 platforms. *The ISME Journal*, **6**(8), pp. 1621-1624.

568 Carda-Diequez, M., Mira, A. and Fouz, B. (2014) Pyrosequencing survey of intestinal  
 569 microbiota diversity in cultured sea bass (*Dicentrarchus labrax*) fed functional diets. *FEMS*  
 570 *Microbiology Ecology*, **87**(2), pp. 451-459.

571 Cerezuela, R., Fumanal, M., Tapia-Paniagua, S.T., Meseguer, J., Morinigo, M.A. and Esteban,  
 572 M.A. (2012) Histological alterations and microbial ecology of the intestine in gilthead sea  
 573 bream (*Spaurus aurata* L.) fed dietary probiotics and microalgae. *Cell and Tissue Research*  
 574 **350**, pp. 477-489

575 Choat, J.H., and Clements, K.D. (1998) Vertebrate herbivores in marine and terrestrial  
 576 environments: A nutritional ecology perspective. *Annual Review of Ecology and Systematics*  
 577 **29** (159), 375-403

578 Clements, K.D., Pasch, I.B.Y., Moran, D. and Turner, S.J. (2007) Clostridia dominate 16S  
 579 rRNA libraries prepared from the hindgut of temperate marine herbivorous fishes. *Marine*  
 580 *Biology* **150**: pp.1431-1440

581 Clements, K.D., Angert, E.R., Montgomery, W.L. and Choat, J.H. (2014) Intestinal microbiota  
 582 in fishes: what's known and what's not. *Molecular Ecology*, **23**(8), pp. 1891-1898.

583 Desai, A.R., Links, M.G., Collins, S.A., Mansfield, G.S., Drew, M.D., Van Kessel, A.G. and  
 584 Hill, J.E. (2012) Effects of plant-based diets on the distal gut microbiome of rainbow trout  
 585 (*Oncorhynchus mykiss*). *Aquaculture*, **350–353**(0), pp. 134-142.

586 Dimitroglou, A., Merrifield, D.L., Carnevali, O., Picchietti, S., Avella, M., Daniels, C.L.,  
 587 Guroy, D. and Davies, S.J. (2011) Microbial manipulations to improve fish health and  
 588 production: A Mediterranean perspective. *Fish and Shellfish Immunology* **30**, 1-16

589 Dyall, S.C., Michael, G.J. and Michael-Titus, A.T. (2010) Omega-3 fatty acids reverse age-  
 590 related decreases in nuclear receptors and increase neurogenesis in old rats. *Journal of*  
 591 *Neuroscience Research* 88(10), pp.2091-2102

592 Dzyuba, E.V., Suhanova, E.V., Denikina, N.N. and Belkova, N.L. (2014) Comparative analysis  
 593 of gut microbiocenoses of salmonid fish with different feeding strategies. *Doklady biological*  
 594 *sciences: proceedings of the Academy of Sciences of the USSR, Biological sciences sections /*  
 595 *translated from Russian*, **11**, pp. 2429-2433.

596 Edgar, R.C., Haas, B.J., Clemente, J.C., Quince, C. and Knight, R. (2011) UCHIME improves  
 597 sensitivity and speed of chimera detection. *Bioinformatics* **27** (16): 2194-200

598 Freundt, E.A. and Razin, S. (1958) Genus *Mycoplasma*. In: Krieg, N.R., Holt, J.G. (eds)  
 599 *Bergey's Manual of Systematic Bacteriology*, vol.1. Williams and Wilkins, Baltimore, pp 742-  
 600 770

601 Gatesoupe, F.J., Huelvan, C., Le Bayon, N., Sevre, A., Aasen, I.M., Degnes, K.F., Mazurais,  
 602 D., Panserat, S., Zambonino-Infante, J.L. and Kaushik, S.J. (2014) The effects of dietary  
 603 carbohydrate sources and forms on metabolic response and intestinal microbiota in sea bass  
 604 juveniles, *Dicentrarchus labrax*. *Aquaculture* **422-423**: pp.47-53

605 Ghanbari, M., Kneifel, W. and Domig, K. (2015) A new view of the fish gut microbiome:  
606 Advances from next-generation sequencing. *Aquaculture* **448** pp. 464-475

607 Green, T.J., Smullen, R. and Barnes, A.C. (2013) Dietary soybean protein concentrate-induced  
608 intestinal disorder in marine farmed Atlantic salmon, *Salmo salar* is associated with alterations  
609 in gut microbiota. *Veterinary Microbiology*, **166**(1-2), pp. 286-292.

610 Holben, W.E., Williams, P., Gilbert, M., Saarinen, M., Sarkilahti, L.K. and Apajalahti, J.H.,  
611 (2002) Phylogenetic analysis of intestinal microflora indicates a novel *Mycoplasma* phylotype  
612 in farmed and wild salmon. *Microbial Ecology*, **44**(2), pp. 175-185.

613 Ingerslev, H., Von Gersdorff Jorgensen, L., Lenz Strube, M., Larsen, N., Dalsgaard, I., Boye,  
614 M. and Madsen, L. (2014a) The development of the gut microbiota in rainbow trout  
615 (*Oncorhynchus mykiss*) is affected by first feeding and diet type. *Aquaculture*, **424–425**(0), pp.  
616 24-34.

617 Ingerslev, H.C., Strube, M.L., Jorgensen, L., Dalsgaard, I., Boye, M. and Madsen, L., 2014b.  
618 Diet type dictates the gut microbiota and the immune response against *Yersinia ruckeri* in  
619 rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, **40**(2), pp. 624-633.

620 Irianto, A. and Austin, B. (2002) Use of probiotics to control furunculosis in rainbow trout,  
621 *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases* **25**, 333-342

622 Joborn, A., Olssen J.C., Westerdahl, A., Conway, P.L. and Kjelleberg, S. (1997) Colonization  
623 in the fish intestinal tract and production of inhibitory substances in intestinal mucous and  
624 faecal extracts by *Carnobacterium* sp. Strain K1. *Journal of Fish Diseases* **20**, 383-392

625 Kim, D.H. and Austin B. (2006) Innate immune responses in rainbow trout (*Oncorhynchus*  
626 *mykiss*, Walbaum) induced by probiotics. *Fish and Shellfish Immunology* **21**, 513-524

627 Kim, D.H., Brunt, J. and Austin, B. (2007) Microbial diversity of intestinal contents and mucus  
628 in rainbow trout (*Oncorhynchus mykiss*). *Journal of Applied Microbiology*, **102**(6), pp. 1654-  
629 1664.

630 Kim., D.H. and Austin, B., 2008. Characterization of probiotic carnobacteria isolated from  
631 rainbow trout (*Oncorhynchus mykiss*) intestine. *Letters in Applied Microbiology*, **47**(3), pp.  
632 141-147.

633 King, G.M., Judd, C., Kuske, C.R. and Smith, C. (2012) Analysis of stomach and gut  
634 microbiomes of the eastern oyster (*Crassostrea virginica*) from coastal Louisiana, USA. *PLoS*  
635 *One* **7**: e51475

636 Kormas, K.A., Meziti, A., Mente, E. and Frentzos, A. (2014) Dietary differences are reflected  
637 on the gut prokaryotic community structure of wild and commercially reared sea bream (*Sparus*  
638 *aurata*). *MicrobiologyOpen*, **3**(5), pp. 718-728.

639 Leser, T.D., Amenuvor, J.Z., Jensen, T.K., Lindecrona, R.H., Boye, M. and Moller, K. (2002)  
640 Culture-independent analysis of gut bacteria: the pig gastrointestinal tract microbiota revisited.  
641 *Applied and Environmental Microbiology* **68**: pp.673-690

642 Ley, R.E., Hamady, M., Lozupone, C., Turnbaugh, P.J., Ramey, R.R. and Bircher, S. (2008)  
643 Evolution of mammals and their gut microbes. *Science* **320** (5883) pp. 1647-1651

644 Lowrey, L., Woodhams, D.C., Tacchi, L. and Salinas, I. (2015) Topographical mapping of the  
645 rainbow trout microbiome reveals a diverse bacterial community in the skin with antifungal  
646 properties. *Applied and Environmental Microbiology* 018126-15 (need to check)

647 Lozupone, C. and Knight, R. (2005) Unifrac: A new phylogenetic method for comparing  
648 microbial communities. *Applied and Environmental microbiology* **71** (12) 8228-8235

649 Mansfield, G.S., Desai, A.R., Nilson, S.A., Van Kessel, A.G., Drew, M.D. and Hill, J.E. (2010)  
 650 Characterization of rainbow trout (*Oncorhynchus mykiss*) intestinal microbiota and  
 651 inflammatory marker gene expression in a recirculating aquaculture  
 652 system. *Aquaculture*, **307**(1–2), pp. 95-104.

653 McCune, B., Grace, J.B. and Urban, D.L. (2002) Analysis of ecological communities. MjM  
 654 Software, Gleneden Beach, Oregon USA

655 Merrifield, D.L., Dimitroglou, A., Foey, A., Davies, S.J., Baker, R.T.M., Bogwald, J., Castex,  
 656 M. and Ringo, E. (2010) The current status and future focus of probiotic and prebiotic  
 657 applications for salmonids. *Aquaculture*, **302**(1–2), pp. 1-18.

658 Merrifield, D.L., Olsen, R.E., Myklebust, R. and Ringo, E. (2011) Dietary effects of soybean  
 659 products on gut histology and microbiota of fish. In: *Soybean and Nutrition* (ed. H. El-Shemy),  
 660 InTech, pp. 231-251

661 Merrifield, D.L., Balcazar, J.L., Daniels, C., Zhou, Z., Carnevali, O., Sun, Y.Z., Hoseinifar, H.  
 662 and Ringo, E. (2014) Indigenous lactic acid bacteria in fish and crustaceans. In: *Aquaculture*  
 663 *Nutrition: Gut Health, Probiotics and Prebiotics*, First Edition pp. 128-156

664 Miyake, S., Ngugi, D.K. and Stingl, U. (2015) Diet strongly influences the gut microbiota of  
 665 surgeonfishes. *Molecular Ecology*, **24**(3), pp. 656-672.

666 Moran, D., Turner, S.J. and Clements, K.D. (2005). Ontogenetic development of the  
 667 gastrointestinal microbiota in the marine herbivorous fish *Kyphosus sydneyanus*. *Microbial*  
 668 *Ecology*, **49**(4), pp. 590-597.

669 National Research Council (2011) Nutrient requirements of fish and shrimp. National  
 670 Academy Press, Washington D.C.

671 Navarrete, P., Magne, F., Araneda, C., Fuentes, P., Barros, L., Opazo, R., Espejo, R. and  
672 Romero, J. (2012) PCR-TTGE analysis of 16S rRNA from rainbow trout (*Oncorhynchus*  
673 *mykiss*) gut microbiota reveals host-specific communities of active bacteria. *PloS one*, **7**(2), pp.  
674 e31335

675 Nayak, S.K. (2010) Role of gastrointestinal microbiota in fish. *Aquaculture Research*, **41**(11),  
676 pp. 1553-1573.

677 Ozorio, R.O.A., Kopecka-Pilarczyk, J., Peixoto, M.J., Lochmann, R., Santos, R.J., Santos, G.,  
678 Weber, B., Calheiros, J., Ferrez-Arruda, L., Vaz-Pires, P. and Goncalves, J.F.M. (2015) Dietary  
679 probiotic supplementation in juvenile rainbow trout (*Oncorhynchus mykiss*) reared under cage  
680 culture production: effects on growth, fish welfare, flesh quality and intestinal microbiota.  
681 *Aquaculture Research* **doi:** 10.1111/are.12724

682 Panigrahi, A., Kiron, V., Kobayashi, T., Puangkaew, J., Satoh, S. and Sugita, H. (2004)  
683 Immune responses in rainbow trout *Oncorhynchus mykiss* induced by a potential probiotic  
684 bacteria *Lactobacillus rhamnosus* JCM 1136. *Veterinary Immunology and*  
685 *Immunopathology*, **102**(4), pp. 379-388.

686 Panigrahi, A., Kiron, V., Puangkaew, J., Kobayashi, T., Satoh S. and Sugita, H. (2005) The  
687 viability of probiotic bacteria as a factor influencing the immune response in rainbow trout  
688 *Oncorhynchus mykiss*. *Aquaculture* **243**, 241-254

689 Perez-Sanchez, T., Balcazar, J.L., Merrifield, D.L., Carnevali, O., Gioacchini, G., de Blas, I.  
690 and Ruiz-Zarzuela, I. (2011) Expression of immune related genes in rainbow trout  
691 (*Oncorhynchus mykiss*) induced by probiotic bacteria during *Lactococcus garviae* infection.  
692 *Fish and Shellfish Immunology* **31**, 196-201

693 Ringo, E. and Olsen, R.E. (1999) The effect of diet on aerobic bacterial flora associated with  
694 intestine of Arctic charr (*Salvelinus alpinus* L.). *Journal of Applied Microbiology*, **86**(1), pp.  
695 22-28.

696 Ringo, E., Sperstad, S., Myklebust, R., Refstie, S. and Krogdahl, A. (2006) Characterization of  
697 the microbiota associated with the intestine of Atlantic cod (*Gadhus morhua* L.): The effect of  
698 fish meal, standard soybean meal and a bioprocessed soybean meal. *Aquaculture* **261**, 829-841

699 Ringo, E., Olsen, R.E., Gifstad, T.O., Dalmo, R.A., Amlund, H., Hemre, G.I. and Bakke, A.M.  
700 (2010) Prebiotics in aquaculture: a review. *Aquaculture Nutrition* **16**: pp.117-136

701 Ringo, E., Zhou, Z., Olsen, R.E. and Song, S.K. (2012) Use of chitin and krill in aquaculture:  
702 the effect on gut microbiota and the immune system. A review. *Aquaculture Nutrition* **18**:  
703 pp.117-131

704 Schloss, P.D. and Handelsman, J. (2006) Introducing TreeClimber, a test to compare microbial  
705 community structures. *Applied and Environmental Microbiology* **72** (4) 2379-2384

706 Schloss, P.D. (2009) Introducing mothur: Open-source, platform-independent, community-  
707 supported software for describing and comparing microbial communities. *Appl Environ*  
708 *Microbiol.* **75**(23):7537-41

709 Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S. and Huttenhower,  
710 C. (2011) Metagenomic biomarker discovery and explanation. *Genome Biology* **12**: R60

711 Smriga, S., Sandin, S. and Azam, F. (2010) Abundance, diversity and activity of microbial  
712 assemblages associated with coral reef fish guts and feces. *FEMS Microbiology Ecology* **73** (1)  
713 pp. 31-42

714 Spanggaard, B., Huber, I., Nielsen, J., Nielsen, T., Appel, K.F. and Gram, L. (2000) The  
 715 microflora of rainbow trout intestine: a comparison of traditional and molecular identification.  
 716 *Aquaculture*, **182**(1–2), pp. 1-15.

717 Suhanova, E.V., Dzyuba, E.V., Triboy, T.I., Nikiforova, T.I., Denikina, N.N. and Belkova,  
 718 N.L. (2011) Molecular genetic and culture diagnosis of *Mycoplasma* in fish family  
 719 Thymallidae. *Doklady biological sciences: proceedings of the Academy of Sciences of the*  
 720 *USSR, Biological sciences sections / translated from Russian*, **440**, pp. 287-289.

721 Sullam, K.E., Essinger, S.D., Lozupone, C.A., O'Connor, M.P., Rosen, G.L., Knight, R.,  
 722 Kilham, S.S. and Russell, J.A. (2012) Environmental and ecological factors that shape the gut  
 723 bacterial communities of fish: a meta-analysis. *Molecular Ecology* **21**, 3363-3378

724 Tacon, A. G. J. and Metian, M. (2008) Global overview on the use of fish meal and fish oil in  
 725 industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* **285**: pp. 146-  
 726 158

727 Vendrell, D., Balcazar, J.L., de Blas, I., Ruiz-Zarzuela, I., Girones, O. and Muzquiz, J.L. (2008)  
 728 Protection of rainbow trout (*Oncorhynchus mykiss*) from lactococcosis by probiotic bacteria.  
 729 *Comparative Immunology, Microbiology and Infectious Diseases* **31**, 337-345

730 Ward, N., Steven, B., Penn, K., Methe, B. and Detrich, W. (2009) Characterization of the  
 731 intestinal microbiota of two Antarctic notothenioid fish species. *Extremophiles* **13** (4), 679-685

732 White, J.R., Nagarajan, N. and Pop, M. (2009) Statistical methods for detecting differentially  
 733 abundant features in clinical metagenomics samples. *PLoS Computational Biology* **5**,  
 734 e1000352

735 Wong, S., Waldrop, T., Summerfelt, S., Davidson, J., Barrows, F., Kenney, P.B., Welch, T.,  
 736 Wiens, G.D., Snekvik, K., Rawls, J.F. and Good, C. (2013) Aquacultured rainbow trout



(*Oncorhynchus mykiss*) possess a large core intestinal microbiota that is resistant to variation in diet and rearing density. *Applied and Environmental Microbiology*, **79**(16), pp. 4974-4984.

Xiao, Y., Wang, L., Xu, R.J., Chen, Z.Y. (2006) DHA depletion in rat brain is associated with impairment on spatial learning and memory. *Biomedical and Environmental Science* **19**(6), 474-480

Xing, M., Hou, Z., Yuan, J., Liu, Y., M Qu, Y. and Liu, B. (2013) Taxonomic and functional metagenomics profiling of gastrointestinal tract microbiome of the farmed adult turbot (*Scophthalmus maximus*) *FEMS Microbiology Ecology* **86** pp. 432-443

Yue, J.C. and Clayton, M.K. (2005) A similarity measure based on species proportions. *Communications in Statistics Theory and Methods* **34** (11) pp. 2123-2131

Zarkasi, K., Abell, G., Taylor, R., Neuman, C., Hatje, E., Tamplin, M., Katouli, M. and Bowman, J.P. (2014) Pyrosequencing based characterization of gastrointestinal bacteria of Atlantic salmon (*Salmo salar* L.) within a commercial mariculture system. *Journal of Applied Microbiology* **117**(1), pp. 18-2

## Figure Legends

**Figure 1.** Mean relative % sequence abundance of microbial phyla recorded in distal intestine of fish fed a) control and b) treatment diet

**Figure 2.** Relative % sequence abundance of tank biofilm, diet and intestinal microbial classes in rainbow trout fed a) control and b) treatment diet.

**Figure 3.** Principal coordinate analysis (PCoA) depicting differences in microbial community structure between control and treatment fish, tank biofilm and feed pellet samples from both diets, based on ThetaYC distance matrix.

**Figure 4.** Bacterial taxa identified by metastats, LEfSe and Indicator analysis as discriminatory between experimental conditions. The data are plotted as mean percentage relative abundance  $\pm$  standard error of the mean (SEM). \* $P < 0.05$  \*\* $P < 0.01$ .

**Figure S1.** Growth performance data for control and treatment fish populations. Mean final weight and condition factor (K)  $\pm$  SEM at the end of the 15 week trial period are shown (n=12). Condition factor was calculated according to Fulton's method.

**Figure S2.** Rarefaction analysis of a) control and b) treatment group sequence libraries. Samples were rarefied according to the library with the lowest number of reads (n=314961, A F6)

**Figure S3.** Mean relative abundance  $\pm$  SEM of sequences attributed to *Acetanaerobacterium* and *Brevinema* in the intestinal microbiome of both control and microalgae fed rainbow trout. Metastats, LEfSe and Indicator analyses did not identify these differences as statistically different according to diet administered, despite differences noted in relative sequence abundances between diets.

**Figure S4.** Heatmap of abundant bacterial genera recorded in this study. Fish are numbered 1-12, green for treatment group, and red for control group. Within the heatmap, RED colours indicate communities that are more similar between samples, whilst BLACK indicates dissimilarity between samples, based on ThetaYC distance matrix.

**Table 1. Alpha diversity estimates of rainbow trout intestinal microbiomes**


Sample	OTU	Coverage	Simpson		Inverse Simpson		Chao 1	
			$\mu$	$\sigma$	$\mu$	$\sigma$	$\mu$	$\sigma$
C F1	97	0.999917	0.64	0.000054	1.55	0.00013	157.35	2.43
C F2	73	0.999908	0.93	0.000026	1.07	0.00003	140.37	2.17
C F3	150	0.999908	0.35	0.000076	2.87	0.00062	206.02	1.88
C F4	90	0.999905	0.38	0.000125	2.63	0.00085	199.86	4.21
C F5	92	0.999917	0.84	0.000038	1.18	0.00005	178.19	4.03
C F6	58	0.999937	0.90	0.000031	1.10	0.00003	128.80	3.39
C F7	68	0.999914	0.94	0.000025	1.06	0.00003	159.93	4.29
C F8	62	0.999937	0.88	0.000033	1.13	0.00004	117.70	2.17
C F9	94	0.99993	0.89	0.000032	1.12	0.00004	143.69	1.96
C F10	173	0.999838	0.80	0.000017	1.25	0.00002	272.98	1.03
C F11	177	0.999882	0.67	0.000049	1.49	0.00010	237.94	1.89
C F12	62	0.999943	0.93	0.000028	1.07	0.00003	114.63	2.20
<b>MeanC</b>	99.6	0.999911	0.76	0.000045	1.46	0.00016	171.45	2.63
A F1	132	0.999835	0.49	0.000115	2.04	0.00047	284.33	4.73
A F2	87	0.999921	0.89	0.000031	1.12	0.00003	168.74	3.04
A F3	185	0.999851	0.65	0.000052	1.52	0.00012	248.07	1.74
A F4	228	0.999895	0.69	0.000051	1.45	0.00010	318.00	3.90
A F5	87	0.999902	0.85	0.000037	1.17	0.00005	182.45	3.47
A F6	192	0.99981	0.44	0.00004	2.28	0.00001	294.62	2.12
A F7	255	0.999816	0.56	0.000064	1.77	0.00020	332.80	1.93
A F8	77	0.999952	0.88	0.000034	1.13	0.00004	127.85	2.43
A F9	134	0.999886	0.59	0.00006	1.68	0.00017	219.07	2.77
A F10	80	0.999917	0.67	0.000052	1.49	0.00011	141.80	2.69
A F11	86	0.999975	0.38	0.000072	2.61	0.00048	103.96	1.01
A F12	83	0.999927	0.90	0.000033	1.11	0.00004	140.91	2.51
<b>MeanA</b>	135.5	0.999891	0.66	0.000053	1.61	0.00015	213.55	2.69

Normalized mean values ( $\mu$ ) and standard deviations ( $\sigma$ ) for the number of OTU's, Sample coverage, Simpson Index, Inverse Simpson Index and Chao 1 richness. Normalized values were obtained by random resampling via rarefaction analysis according to the smallest sample size ( $n=314961$ , A F6) and standard errors were obtained by bootstrapping. OTU's are clustered according to a 97% sequence similarity cut-off value. C = Control samples, A= Treatment samples (Algae).

Phylotype	p value		
	Metastats	LEfSe	Indicator
<i>Acetanaerobacterium</i>	0.94	-	0.68
<i>Brevinema</i>	0.41	-	0.39
<i>Streptococcus</i>	0.009	0.009	0.042
<i>Leuconostoc</i>	0.009	0.013	0.046
<i>Weissella</i>	0.043	-	0.12
<i>Candidate division WPS-1</i>	0.006	0.005	0.024
<i>Lactobacillus</i>	0.010	0.007	0.034
<i>Lactococcus</i>	0.046	0.026	0.058
<i>Enterobacter</i>	0.034	0.049	0.078

**Table 2.** Phylotypes identified as discriminatory according to diet by three separate statistical algorithms within mothur (Metastats, LEfSe and Indicator). Statistical significance was accepted on two levels;  $p < 0.05$  and  $p < 0.01$ . *Acetanaerobacterium* and *Brevinema* were not discriminatory by diet.

a)



b)





Figure 1

a)



b)






Figure 2.



**Figure 3.**



**Figure 4.**

808 **Table S1. Ingredient composition and nutrient analysis of diets**

	<b>Diet 1 (Control)</b>	<b>Diet 2 (Treatment)</b>
<b>Ingredient</b>	<b>% Inclusion</b>	<b>% Inclusion</b>
Fish meal 68	22	22
Wheat meal	15	14
Wheat gluten	10	10
Soybean meal 47	14	14
Soybean concentrate 65	20	19
Alltech algae meal	0	5
Fish oil	15	12
Monocalcium phosphate	1.3	1.3
Mineral and vitamin premix	1	1
Lysine	0.6	0.6
Methionine	0.5	0.5

809


<b>Dietary component</b>		
Moisture	6.0	6.1
Protein	45.4	45.2
Fat	18.4	18.3
Ash	5.9	6.0
Fibre	1.1	1.1
NFE	20	19.2

810

811


812





**Figure S1**

a)



b)





Figure S2



**Figure S3**

