

- 1 The harlequin ladybird, *Harmonia axyridis*: global perspectives on invasion history and
2 ecology
3
4 Helen E. Roy¹
5 Peter M.J. Brown²
6 Tim Adriaens³
7 Nick Berkvens⁴
8 Isabel Borges⁵
9 Susana Clusella-Trullas⁶
10 Patrick De Clercq⁷
11 Richard F. Comont⁸
12 Rene Eschen⁹
13 Arnaud Estoup¹⁰
14 Edward W. Evans¹¹
15 Benoit Facon¹⁰
16 Mary M. Gardiner¹²
17 Artur Gil⁵
18 Audrey A. Grez¹³
19 Thomas Guillemaud^{14, 15, 16}
20 Danny Haelewaters^{17, 18}
21 Annette Herz¹⁹
22 Alois Honek²⁰
23 Andy G. Howe²¹
24 Cang Hui^{22, 23}
25 William D. Hutchison²⁴
26 Marc Kenis⁹
27 Robert L. Koch²⁴
28 Jan Kulfan²⁵
29 Lori Lawson Handley²⁶
30 Eric Lombaert^{14, 15, 16}
31 Antoon Loomans²⁷
32 John Losey²⁸
33 Alexander O. Lukashuk²⁹
34 Dirk Maes³
35 Alexandra Magro³⁰

36 Katie M. Murray³¹
 37 Gilles San Martin³²
 38 Zdenka Martinkova²⁰
 39 Ingrid A. Minnaar⁶
 40 Oldřich Nedved³³
 41 Marina J. Orlova-Bienkowskaja³⁴
 42 Wolfgang Rabitsch³⁵
 43 Hans Peter Ravn²¹
 44 Gabriele Rondoni³⁶
 45 Steph L. Rorke¹
 46 Sergey K. Ryndevich³⁷
 47 May-Guri Saethre³⁸
 48 John J. Sloggett³⁹
 49 Antonio Onofre Soares⁵
 50 Riaan Stals⁴⁰
 51 Matthew C. Tinsley³¹
 52 Axel Vandereycken⁴¹
 53 Paul van Wielink⁴²
 54 Sandra Viglášová^{1,24}
 55 Peter Zach²⁴
 56 Ilya A. Zakharov⁴³
 57 Tania Zaviezo⁴⁴
 58 Zihua Zhao⁴⁵

59
60 ¹Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB, UK

61 ²Animal & Environment Research Group, Anglia Ruskin University, Cambridge, CB1 1PT, UK

62 ³Research Institute for Nature and Forest (INBO), Kliniekstraat 25, B-1070 Brussels, Belgium

63 ⁴Institute for Agricultural and Fisheries Research, Dpt. Plant Crop Protection Entomology,
64 Burgemeester Van Gansberghelaan 96, B-9820 Merelbeke

65 ⁵cE3c - ABG - Center for Ecology, Evolution and Environmental Changes and Azorean Biodiversity
66 Group, Department of Biology, University of the Azores, 9501-801 Ponta Delgada, Portugal

67 ⁶Centre for Invasion Biology, Department of Botany and Zoology, Stellenbosch University, Private
68 Bag X1, Matieland 7602, South Africa

69 ⁷Department of Crop Protection, Ghent University, Coupure Links 653, B-9000 Ghent, Belgium

70 ⁸Bumblebee Conservation Trust, UK

71 ⁹CABI, Delémont, Switzerland

72 ¹⁰Inra, UMR 1062 Cbgp, Montpellier, France

73 ¹¹Department of Biology, Utah State University, Logan, Utah, 84322-5305, USA

74 ¹²Department of Entomology, The Ohio State University Ohio Agricultural Research and
75 Development Center, 1680 Madison Avenue, Wooster, Ohio USA

76 ¹³Facultad de Ciencias Veterinarias y Pecuarias, Universidad de Chile, Casilla 2-15, Santiago, Chile

77 ¹⁴Inra, UMR 1355 ISA, Sophia-Antipolis, France

78 ¹⁵Université de Nice Sophia Antipolis, UMR ISA, Sophia-Antipolis, France

79 ¹⁶CNRS, UMR 7254 ISA, Sophia-Antipolis, France

80 ¹⁷Department of Organismic and Evolutionary Biology, Harvard University, Cambridge,
81 Massachusetts 02138, USA

82 ¹⁸Farlow Reference Library and Herbarium of Cryptogamic Botany, Harvard University, 22 Divinity
83 Avenue, Cambridge, Massachusetts 02138, USA

84 ¹⁹Institute for Biological Control, JKI, Heinrichstr 243, D-64287 Darmstadt, Germany

85 ²⁰Crop Research Institute, Prague-Ruzyne, Czech Republic

86 ²¹Department of Geosciences and Natural Resource Management, University of Copenhagen,
87 Rolighedsvej 23, Frederiksberg 1958, Denmark

88 ²²Centre for Invasion Biology, Department of Mathematical Sciences, Stellenbosch University,
89 Matieland 7602

90 ²³The African Institute for Mathematical Sciences, Cape Town 7945, South Africa

91 ²⁴Department of Entomology, University of Minnesota, Saint Paul, MN 55108 USA

92 ²⁵Institute of Forest Ecology, Slovak Academy of Sciences, Slovakia

93 ²⁶Evolutionary Biology Group, Department of Biological Sciences, University of Hull, Hull, UK

94 ²⁷Department of Entomology, National Reference Centre, National Plant Protection Organization,
95 P.O. Box 9102, 6700 HC Wageningen, Netherlands

96 ²⁸Department of Entomology, Cornell University, Ithaca, NY 14853 USA

97 ²⁹Berezinskiy Biosphere Reserve, Domzheritsy, Lepel distr., Vitebsk reg., 211188, Belarus

98 ³⁰University of Toulouse - ENFA, UMR CNRS / UPS / ENFA 5174 Évolution et diversité biologique, F-
99 31326 Castanet-Tolosan, France

100 ³¹Biological and Environmental Sciences, Stirling University, Stirling FK9 4LA, UK

101 ³²Walloon Agricultural Research Centre, Life Sciences Department, Plant Protection and
102 Ecotoxicology Unit, Rue de Liroux 2, B-5030 Gembloux, Belgium

103 ³³University of South Bohemia and Institute of Entomology, České Budějovice, Czech Republic

- 104 ³⁴A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, 33 Leninskiy
105 Prospect, Moscow 119071, Russia
- 106 ³⁵Environment Agency Austria, Vienna, Austria
- 107 ³⁶Department of Agricultural, Food and Environmental Sciences, University of Perugia, Borgo XX
108 Giugno 74, 06121 Perugia, Italy
- 109 ³⁷Baranovich State University, Voykova str. 21, Baranovich, 225404, Brest reg., Belarus
- 110 ³⁸Bioforsk - NIBIO-Norwegian Institute of Bioeconomy Research, Biotechnology and Plant Health
111 Division, Department of Entomology and Nematology, Høgskoleveien 7, 1432 Ås, Norway
- 112 ³⁹Maastricht Science Programme, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The
113 Netherlands
- 114 ⁴⁰South African National Collection of Insects, ARC-Plant Protection Research, Private Bag X134,
115 Queenswood, Pretoria 0121, South Africa
- 116 ⁴¹University of Liege, Gembloux Agro-Bio Tech, Functional and Evolutionary Entomology, Passage des
117 Déportés 2, 5030 Gembloux
- 118 ⁴²Tobias Asserlaan 126, 5056 VD Berkel-Enschot, The Netherlands
- 119 ⁴³Vavilov Institute of General Genetics, Moscow, Gubkin Str. 3, 119991, Russia
- 120 ⁴⁴Facultad de Agronomía e Ingeniería Forestal, Pontificia Universidad Católica de Chile, Casilla 306-
121 22, Santiago, Chile
- 122 ⁴⁵Department of Entomology, China Agricultural University, Beijing 100193, China

Abstract

The harlequin ladybird, *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae), is native to Asia but has been intentionally introduced to many countries as a biological control agent of pest insects. In numerous countries, however, it has been introduced unintentionally. The dramatic spread of *H. axyridis* within many countries has been met with considerable trepidation. It is a generalist top predator, able to thrive in many habitats and across wide climatic conditions. It poses a threat to biodiversity, particularly aphidophagous insects, through competition and predation, and in many countries adverse effects have been reported on other species, particularly coccinellids. However, the patterns are not consistent around the world and seem to be affected by many factors including landscape and climate.

Research on *H. axyridis* has provided detailed insights into invasion biology from broad patterns and processes to approaches in surveillance and monitoring. An impressive number of studies on this alien species have provided mechanistic evidence alongside models explaining large-scale patterns and processes. The involvement of citizens in monitoring this species in a number of countries around the world is inspiring and has provided data on scales that would be otherwise unachievable.

Harmonia axyridis has successfully been used as a model invasive alien species and has been the inspiration for global collaborations at various scales. There is considerable scope to expand the research and associated collaborations, particularly to increase the breadth of parallel studies conducted in the native and invaded regions. Indeed a qualitative comparison of biological traits across the native and invaded range suggests that there are differences which ultimately could influence the population dynamics of this invader. Here we provide an overview of the invasion history and ecology of *H. axyridis* globally with consideration of future research perspectives. We reflect broadly on the contributions of such research to our understanding of invasion biology while also informing policy and people.

Keywords: Coccinellidae, biocontrol, species traits, competitive interactions, invasion history

Introduction

The harlequin ladybird, *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae), is native to Asia and is a polymorphic, eurytopic species with a broad dietary range (Roy, Brown 2015a). It was widely introduced as a biological control agent of pest aphids, but has spread to many countries within which it was not intentionally released (Brown et al. 2008a). The global invasion of *H. axyridis* has been rapid (Brown et al. 2011b). Concerns about the adverse impact of *H. axyridis* particularly on biodiversity but also to people have motivated research across the world. There have been a number of reviews considering the biology and ecology of *H. axyridis* (Kenis et al. 2008; Koch 2003; Koch, Galvan 2008; Majerus et al. 2006; Pell et al. 2008; Pervez, Omkar 2006; Roy, Brown 2015a; Sloggett 2012) often from a regional perspective but this species has an almost global distribution. Therefore, it is timely to provide a world overview of research and perspectives. Indeed, the collaborative nature of the research from around the world on this species is inspiring. As concerns increased with respect to the threats posed by *H. axyridis*, two working groups were established through the International Organisation for Biological and Integrated Control (within the Global and Western Palaearctic Regional Sections) in recognition of the need for collaborative research on the benefits and risks of exotic biological control agents. One of the first outputs from these working groups was the publication of a special issue on *H. axyridis* including 19 manuscripts representing authors around the world (Roy, Wajnberg 2008). *Harmonia axyridis* as both a biological control agent and an invasive alien species (IAS) has informed a range of applied ecological themes, from risk assessment to processes of invasion; *H. axyridis* is considered by many as a model species for understanding processes of invasion (Roy, Wajnberg 2008).

Here we first consider *H. axyridis* in the context of biological control. We then provide a brief overview of interactions between *H. axyridis* and other species before exploring its current global distribution. We document, on a regional basis, the occurrence and invasion history of *H. axyridis* providing an insight into research priorities and knowledge gaps identified by scientists from around the world. Finally we qualitatively explore the observed traits of *H. axyridis* from both the native and invaded range reflecting on the potential for future collaborations on a global scale.

Harmonia axyridis and biological control

There has been a long history of using ladybirds as biological control agents against various pest insects around the world, beginning with the successful introduction of the vedalia ladybird, *Rodolia cardinalis* (Mulsant) to citrus groves of California (Caltagirone, Doult 1989). *Harmonia axyridis* has a wide diet breadth (reviewed by Hodek and Evans 2012) including many species of aphids, which are

the main prey (Osawa 2000) but also other insect taxa (Ohgushi, Sawada 1998). It has been used extensively in classical, augmentative and conservation biological strategies around the world (Koch 2003; Lombaert et al. 2008). There has been considerable research focus on the pest control services provided by *H. axyridis* in the native range, particularly in China (Wang et al. 2007a), but also recently in Japan where adults and larvae of flightless *H. axyridis* derived through artificial selection from a Japanese wild population have been used for biological control of aphids mainly in greenhouses (Seko et al. 2014).

Harmonia axyridis was first introduced as a biological control agent in the USA. It is among a number of introduced species of ladybirds that now dominate in many agroecosystems across the USA (Lucas et al. 2007; Obrycki et al. 2009) and *H. axyridis* is considered an important predator of aphid pests in several crops, including pecan (LaRock et al. 2003; Tedders, Schaefer 1994), apple (Brown, Miller 1998), citrus (Michaud 2002) and potatoes (Alyokhin, Sewell 2004). In particular, research has focused on the effect of *H. axyridis* on the suppression of the alien soybean aphid, *Aphis glycines* Matsumura (Hemiptera: [Aphididae](#)) (Ragsdale et al. 2011) in soybean, where *H. axyridis* has recently become one of the most abundant coccinellids (Costamagna, Landis 2007; Gardiner, Landis 2007; Gardiner et al. 2009a; Gardiner et al. 2009b; Hesler 2014; Liere et al. 2014; Varenhorst, O'Neal 2012). This system, including the overwintering host of *A. glycines*, the European buckthorn *Rhamnus cathartica* L. ([Rhamnaceae](#)) and *H. axyridis*, has been cited as an example of an invasional meltdown, whereby multiple IAS interact synergistically (Heimpel et al. 2010). *Harmonia axyridis* is known to readily consume other predators and parasitoids of aphids (Chacón et al. 2008), but there is a lack of evidence that this impacts on pest control in soybean fields (Costamagna et al. 2008).

From the mid 1990s, *H. axyridis* was commercialized by a number of biological control suppliers in Western Europe for augmentative biological control of aphid pests in greenhouse crops and urban ecosystems (Coutanceau 2006b; Poutsma et al. 2008). Most commercial suppliers in Europe stopped selling the beetle in late 2003 to mid 2004, with the first reports of nuisance problems and increasing concerns about adverse environmental effects of its establishment. In France, the original strain of *H. axyridis* used since 1995 for commercial biological control was replaced in 2000 with a flightless strain developed by INRA (Coutanceau 2006b; Tourniaire et al. 2000). The flightless strain was effectively used to control aphids in hops (Weissenberger et al. 1999). However, the species was never a major player on the European biological control market: at the peak of its commercialisation, it took perhaps 5% of the market share of aphidophagous natural enemies (De Clercq, Bale 2011).

The role of *H. axyridis* in suppressing pest insects in Europe has received less attention than in the USA. In the Czech Republic it has been reported that *H. axyridis* controls pear psylla, *Cacopsylla pyri* (L.) (Hemiptera: Psyllidae), in commercial orchards more effectively than other ladybird species (Nedvěd 2014). However, much of the research across Europe has considered the occurrence of *H. axyridis* in crop systems without detailed consideration of effects on aphid populations. In Belgium, the population dynamics of *H. axyridis* in crop systems (wheat, corn, broad bean and potato crops) was studied through field surveys. It has been shown that *H. axyridis* is recorded in such crops 7-8 days after the dominant native ladybirds (Jansen, Hautier 2008; Vandereycken et al. 2013). A one-year study involving field observations in wheat and bean crops in southern England reported an absence of *H. axyridis* in wheat (aphid abundance was reported as low), but presence of *H. axyridis* co-occurring with other ladybirds in bean crops (Wells 2011). *Harmonia axyridis* was the most common aphid enemy species in bean crops and the presence of this species was correlated with high aphid abundance (Wells 2011), perhaps unsurprisingly since the prey biomass required by each single larva to reach the adult stage is very high (Soares et al. 2001, 2003, 2004).

Even though *H. axyridis* can be considered an effective biological control agent at least from the US studies, the Insurance Hypothesis predicts that control will in the long term be better achieved with a diverse array of natural enemies (Loreau et al. 2003). Since *H. axyridis* tends to depress the diversity of coccinellid assemblages (see below), consistently effective biological control may be threatened by the invasion of *H. axyridis*. Further research is needed to unravel the role of *H. axyridis* as a biological control agent of insect pests. Comparison of aphid populations before and after the arrival of *H. axyridis*, or among places with higher and lower abundances of *H. axyridis*, may provide useful information in this respect. Importantly, landscape composition has been found to influence the effectiveness of biological control by *H. axyridis* and other predators. Indeed, soybean fields embedded within diverse landscapes receive a greater pest control service from aphid predators than fields within simplified agricultural landscapes (Gardiner et al. 2009c) compared to landscapes dominated by forests and grasslands where soybean field size was reduced (Woltz, Landis 2014). Similar studies have been carried out in Chile, in regions where *H. axyridis* dominated the aphidophagous communities and again it has been demonstrated that biological control was related to landscape composition with benefits seen through positive associations with the abundance of woodland and urban habitats, but not with fruit crops, in the landscape (Grez et al. 2014a).

Harmonia axyridis and declines of native ladybirds

The wide diet breadth and recognition that *H. axyridis* is a top predator (Pell et al. 2008) has driven predictions that *H. axyridis* has the potential to adversely affect aphidophagous guilds. A number of large-scale analyses have indicated that declines of native ladybirds correlate with the establishment of *H. axyridis*. Indeed declines of native ladybirds have been reported across the USA (Alyokhin, Sewell 2004; Bahlai et al. 2014; Colunga-Garcia, Gage 1998; Evans 2004; Harmon et al. 2007; Hesler, Kieckhefer 2008; Losey et al. 2014; Majerus et al. 2006; Wheeler Jr, Hoebeke 1995) and Europe (Roy et al. 2012b). Following the establishment of *H. axyridis* in Michigan a decrease in populations of three species of ladybird has been reported: *Brachiacantha ursina* (F.), *Cycloneda munda* (Say) and *Chilocorus stigma* (Say) (Colunga-Garcia, Gage 1998), followed more recently by a decline in *Coleomegilla maculata* (DeGeer) (Coccinellidae) (Bahlai et al. 2015). However, this last study also noted that declines of several species reported previously from the same site (Colunga-Garcia, Gage 1998) appear to have stabilized or reversed, having become statistically undetectable (Bahlai et al. 2015). Also in the USA, Michaud (2002) reported *H. axyridis* to be displacing *Cycloneda sanguinea* (L.) in Florida citrus orchards. In the United Kingdom (UK) there is a strong correlation between the declines of seven out of eight native species of ladybird assessed and co-occurrence with *H. axyridis* (Roy et al. 2012b). In Chilean alfalfa fields the abundance of native ladybird species declined after *H. axyridis* was first observed in this crop in 2008. Furthermore, total ladybird species and diversity in alfalfa have also declined during this period (Grez & Zaviezo unpublished data). It is thought that the spread of *H. axyridis* has caused the decline of native species, *Adalia bipunctata* (L.) and *Propylea quatuordecimpunctata* (L.) in the Ukraine (Verizhnikova 2011) and *C. septempunctata* and *A. bipunctata* in Moldova (Iazlovețchii, Sumencova 2013), but this requires confirmation through further research.

In contrast, some long-term studies (Honek et al. submitted) have highlighted that some ladybirds native to Central Europe had already been declining before the arrival of *H. axyridis*. Indeed, the diversity of native ladybird communities were similar before (Honek et al. 2014) and after (Honěk, Martinkova et al. submitted) the arrival of *H. axyridis* in the Czech Republic. Furthermore, in Switzerland, on-going long-term population studies have highlighted that, so far, only *A. bipunctata* has significantly declined since the arrival of *H. axyridis* (Kenis and Eschen, unpublished data) although risk assessments (Kenis et al. 2010) predicted that three other ladybirds sharing the same ecological niches with *H. axyridis* (*A. decempunctata*, *Oenopia conglobata* (L.) and *C. decemguttata*) were at risk.

In Belgium, high niche overlap between *H. axyridis* and generalist native species, particularly *A. bipunctata* and *P. quatuordecimpunctata*, was observed, suggesting a high potential for impact of *H. axyridis* on those species. Large-scale mapping data showed substantial range contraction of *A. bipunctata*, *Adalia decempunctata* (L.), *Calvia quatuordecimguttata* (L.), *Exochomus quadripustulatus* (L.) and *P. quatuordecimpunctata* after the arrival of *H. axyridis*. As a consequence of the invasion, systematic surveying was set up in Brussels using standardized beating of trees in parks, avenues and roadsides. With the exception of *C. quatuordecimguttata*, these abundance data reflected the reported large scale trends (Roy et al. 2012b). *Adalia bipunctata* exhibited a 57% decline in its extent of occurrence in the last decade and according to a conservative application of the IUCN guidelines has now become a good example of species that was formerly widespread but now meets the criteria of a red list species (Adriaens et al. 2015). However, as for Czech Republic, *A. bipunctata* and *P. quatuordecimpunctata* were already in decline prior to the invasion.

The clear differences in the response of ladybird assemblages to the arrival of *H. axyridis* between countries highlight the need for comparative studies. In a recent study it was concluded that differences in species trends between central (Czech Republic) and western Europe (UK) could be attributed to suboptimal environmental conditions in the UK, which is the edge of the biogeographic range for many ladybirds, exacerbating the negative effects of *H. axyridis* (Brown, Roy *In press*). Such interactions between drivers of change are undoubtedly important in population dynamics, and the negative effects of *H. axyridis* are likely to be the result of a complex range of interactions and processes including resource competition and intra-guild predation (IGP) (Majerus et al. 2006).

Long-term and large-scale data on the distribution and abundance of ladybirds are critical for the detection of population changes. To support the collection of long-term survey data, citizen science programmes such as the UK Ladybird survey, Lost Ladybug Project and Buckeye Lady Beetle Blitz in the USA, Chinita arlequin in Chile and the harlequin ladybird survey in Norway have been established to track changes in ladybird populations (Gardiner et al. 2012; Losey et al. 2007; Roy, Brown 2015a; Saethre et al. 2010).

Harmonia axyridis and direct competitive interactions

The majority of research examining interactions between *H. axyridis* and other aphidophagous species has focused on direct interference competition on other ladybirds, proposing that decline is

due to strong asymmetric IGP of eggs and larvae in favour of *H. axyridis* (Pell et al. 2008). This hypothesis is supported by many laboratory and field cage studies around the world (Cottrell 2004; Gardiner, Landis 2007; Hoogendoorn, Heimpel 2002; Katsanis et al. 2013; Roy et al. 2008b; Snyder et al. 2004; Soares, Serpa 2007; Ware, Majerus 2008; Ware et al. 2009; Yasuda et al. 2004). Under open field conditions in Ohio, USA, eggs of the exotic *H. axyridis* were subject to lower predation relative to the eggs of native species (Smith, Gardiner 2013). However, out of 342 attacks, video surveillance illustrated that only two were attributable to alien ladybirds (*H. axyridis* feeding on conspecific eggs). Instead, a diverse guild of predators (dominated by Opiliones, Tettigoniidae and the native *C. maculata*) were responsible for the majority of egg attacks (Smith, Gardiner 2013). Importantly, this study accounted only for egg predation but alien and native ladybird species may compete directly through consumption of other life stages. Other tools, such as alkaloid (defensive chemicals within coccinellids) sequestration analyses, frass analysis, and PCR-based gut content analyses, have been developed to quantify the actual extent of interference competition occurring among native and alien species (Brown et al. 2015; Davidson, Evans 2010; Gagnon et al. 2011; Hautier et al. 2008; Hautier et al. 2011; Sloggett et al. 2009; Thomas et al. 2013). For example, Hautier et al. (2011) found that 20.5% of 590 *H. axyridis* larvae in Belgium tested positive for native ladybird alkaloids. Thomas et al. (2013) detected the DNA of native ladybirds within *H. axyridis* in the UK, with 3.7-22.7% of 156 *H. axyridis* found to have consumed native species over three years. Gagnon et al. (2011) used molecular gut content analysis to illustrate that IGP among larval ladybirds in soybean fields can be very high; 52.9% of sampled ladybirds contained the DNA of one or more other ladybird species. PCR primers have been developed and used to track aphid and ladybird predation by *H. axyridis* fourth-instar larvae collected in lime trees in Italy, and it was found that 7% of sampled individuals contained the DNA of one of two native ladybird species (Rondoni et al. 2014).

There has been less research on the interactions between *H. axyridis* and species within the aphidophagous guild beyond ladybirds (Pell et al. 2008). However, laboratory and field studies have also indicated direct interactions with non-coccinellid aphidophagous predators. *Harmonia axyridis* is an intra-guild predator of *Episyrphus balteatus* (Zeller) (Diptera: Syrphidae) with the strength of this asymmetric interaction increasing with developmental stage of *H. axyridis* and decreasing in the presence of extraguild prey (Ingels, De Clercq 2011). The aphid-specific pathogenic fungus *Pandora neoaphidis* (Remaudiere and Hennebert) Humber (Zygomycota: Entomophthorales: Entomophthoraceae) is consumed by *H. axyridis* (Roy et al. 2008a) whereas most other aphid predators avoid consumption of infected aphids (Roy et al. 1998). Interestingly cage experiments in Italy have been used to study predation on immature stages of *H. axyridis*, and it has been observed

that ants exhibit high levels of predation (Burgio et al. 2008a). Further research has demonstrated the influence of many factors on IGP, including intrinsic (such as feeding history of the species) and extrinsic (such as habitat complexity), and highlighted the importance of addressing such factors when considering the ecological relevance and extent of IGP (Ingels et al. 2015).

Harmonia axyridis and indirect competitive interactions

Harmonia axyridis may also adversely affect aphidophagous species through exploitative competition for shared resources (Evans et al. 2011). Studies to date have failed to show that *H. axyridis* negatively affects native North American ladybirds through exploitative competition among larvae (Hoogendoorn, Heimpel 2004; Yasuda et al. 2004), however, field and microcosm studies reveal reduced fitness in the predatory bug *Anthocoris nemoralis* (F.) (Hemiptera: Anthocoridae) due to competition with *H. axyridis* (Howe et al. 2015). Evans (2004) proposed that following the introduction of alien competitors (specifically *C. septempunctata*), native species abandoned croplands due to increased competition for shared prey. Adults of native species were thought to be displaced to ancestral (refuge) habitats not heavily exploited by alien competitors (Evans 2004; Snyder 2009). Evans (2004) tested this hypothesis by increasing aphid abundance in a Utah alfalfa fields and found that native ladybirds were drawn back into the field, apparently from refuges maintained in the surrounding landscape. In Japan *Harmonia axyridis* coexists with other predators within natural habitats, without adverse negative impacts on co-occurring species (Osawa 2011).

Alien ladybirds and landscape-scale processes

Landscape change can strongly influence species declines within a region (Lindborg, Eriksson 2004). Factors such as habitat degradation, reduction of habitat patch size, and increased isolation of habitats (loss of connectivity) can also lead to declines in the abundance and diversity of coccinellid species within a landscape (Gardiner et al. 2009b; Grez et al. 2013; Grez et al. 2014b). Ladybirds may forage across several habitats within a given landscape including forests, grasslands, and croplands. Landscape changes that alter the distribution of these habitats may affect ladybirds by influencing prey populations, overwintering habitats, or by facilitating the invasion of alien competitors. Observations from several countries suggest that a high proportion of native ladybirds occur in less disturbed habitats, while a high proportion of alien ladybirds (including *H. axyridis*) occurs in human-modified habitats (Grez et al. 2013; Panigaj et al. 2014).

Landscape composition and heterogeneity can also affect competitive interactions among coccinellids (Gardiner et al. 2009a; Gardiner et al. 2009b). In the USA, alien species were more abundant within soybean fields embedded within forested landscapes, and native species more common in agricultural landscapes with significant forage and grassland habitat. A large-scale study assessing the influence of landscape factors on the spread of *H. axyridis* across the UK indicated that coniferous woodland may negatively affect the spread of this species (Purse et al. 2014). Further research is needed to examine how the composition of the surrounding landscape influences the stability of native populations in the invaded regions.

Harmonia axyridis as a household and agricultural pest

Many ladybirds in temperate regions migrate for overwintering to elevated and conspicuous elements in landscape, or even to hill tops (Hodek et al. 2012). There they seek crevices in rocks, in bark of prominent trees and other shelters including south-facing anthropogenic structures such as buildings and monuments (Koch, Galvan 2008; Roy, Brown 2015a; Wang et al. 2011). *Harmonia axyridis* forms large aggregations during the winter months across the native and invaded range in natural areas but also within homes and other structures (Nalepa 2007; Roy, Brown 2015a). A recent study from Poland highlighted the use of wind turbines as overwintering sites for *H. axyridis* (Dudek et al. 2015). It shows hypsotactic behaviour (moving towards prominent objects on the horizon) and a clear preference for contrasting visual elements; vertically positioned stripes being more attractive than horizontal ones (Nalepa et al. 2005; Obata 1986). The relative attractiveness of different surface colours has been studied; white is the most attractive colour followed by yellow and black then green then red and finally natural wood (Wang et al. 2011). Volatile aggregation pheromones are not involved in this orientation (Nalepa et al. 2000) but it is apparent that contact chemoreception is important in the establishment of large aggregations (Durieux et al. 2012).

Harmonia axyridis accounts for 97% of observations from houses contributed by citizen scientists to the US Lost Ladybug Project (Ramsey, Losey 2012). Infestations in homes can cause staining damage to carpet and furnishings and cause allergic reactions (Goetz 2008; Koch, Galvan 2008), but most often are considered a nuisance. Additionally, although this species does not have a greater propensity to bite humans than other ladybirds, its propensity to aggregate in dwellings in high numbers has resulted in a significant increase in the number of ladybird bites reported (Ramsey, Losey 2012). After infestation of the intensive care unit in a large hospital in Austria, counteractive measures including relocation of patients and temporarily closure of the station generated

considerable financial costs. Management tactics have been developed to mitigate such nuisance problems (Kemp, Cottrell 2015).

Harmonia axyridis has been reported to feed on many fruit crops in many parts of the world including grapes, stone fruit, apples, pumpkins and berry crops (Koch et al. 2004a; Koch et al. 2006b). Depending on fruit type, this feeding damage includes primary injury as well as secondary feeding on wounded tissue. In China *H. axyridis* has been documented foraging on pollen, nectar and young plant tissues, occasionally causing serious damage to fruits (Guo, Wan 2001; Yang et al. 2006). In contrast in Japan *H. axyridis* does not consume orchard fruits and overwintering *H. axyridis* are not regarded as a household pest. However, economically, *H. axyridis* presents the largest threat as a contamination pest in wine grape production (Koch, Galvan 2008). Beetles can be found within vineyards largely between the onset of ripening and harvest (Galvan et al. 2008). Adults aggregate and feed on injured fruit clusters (Koch, Galvan 2008). When fruit are harvested and crushed, *H. axyridis* release methoxypyrazines (MPs) which create an unpleasant odour and taste in the wine produced (Botezatu et al. 2013; Galvan et al. 2008). In the Midwestern US, the dramatic increase in *H. axyridis* in wine grapes generally follows population declines in nearby soybean and maize fields (Bahlai, Sears 2009; Galvan et al. 2006; Galvan et al. 2008); however, more research is necessary to document this dispersal hypothesis.

A push-pull strategy including artificial injury of selected fruit clusters on the vineyard margin and spraying the adjacent rows with bisulphite has been proposed as a management option (Glemser et al. 2012; Nedvěd 2014). In addition, progress has been made in developing both natural and synthetic corks that can significantly reduce MP concentrations in wine (Pickering et al. 2010). However, more research is needed to evaluate this approach for additional varietal sources and cost effectiveness. The establishment of *H. axyridis* in regions where the production of wine is economically important, including Crimea, the Caucasus, Canada, Europe, South Africa and South America, continues to be a concern, and close monitoring of the populations in these countries is critical.

Global distribution and regional research priorities

Harmonia axyridis has become one of the world's most widely distributed ladybirds (Figure 1) and is now found on all continents except Antarctica and notably is absent from a few large countries such as Australia. Outside of its native range (Asia), it has become very widely established in North

America, South America and Europe (Figure 2), and in limited parts of Africa. There is a small distance between the western edge of the native range and the eastern edge of the invaded range (see *Russia and adjacent countries* below), and it seems inevitable that these populations will meet in the near future.

In this section we document the distribution of *H. axyridis* around the world and its invasion history (summarized in Table 1), including the perspectives on regions where it has failed to invade. Additionally, current and future research priorities in different countries are presented.

Asia

Harmonia axyridis is native to East Asia (Mongolia, parts of China and Russia, northern Vietnam, Japan and Korea) (Figure 1).

China

Harmonia axyridis has a wide geographical range in China, especially in the north (Wang et al. 2007a). Current research on *H. axyridis* in China is mainly focused on agricultural applications, including the development of artificial diet for *H. axyridis* to maximise production of this species for field release in augmentative biological control (Wang et al. 2007b). *Harmonia axyridis* has also been used extensively for testing pesticides and specifically examining the sensitivity of natural enemies to pesticides (Tang et al. 2014). There is also ongoing research on the genetics of *H. axyridis* colour polymorphism, specifically how colour variation interacts with environmental factors.

Japan

In Japan, *H. axyridis* occurs on the islands of Hokkaido, Honshu, Shikoku, Kyushu, Tsushima, Iki and Gotoh, while its sibling species *H. yedoensis* occurs on Honshu, Shikoku, Kyushu, Yakushima and the Ryukyus (Kurosawa et al. 1985). Future research in Japan will focus on intensive field studies to establish the mechanisms which enable coexistence of *H. axyridis* and other ladybird species in its native range.

Asian part of Russia, Kazakhstan and Kyrgyzstan

The northern part of the native range of *H. axyridis* occupies the south of Siberia and the Far East. The type locality of the species is in Siberia (Pallas 1771). The geographical distribution and colour variability of this species was first studied in the 1920s (Dobzhansky 1924). A recent map of the native range has been compiled (Orlova-Bienkowskaja et al. 2015). *Harmonia axyridis* populations

west of Baikal Lake (in West Siberia and the western part of East Siberia) differ both genetically and morphologically from those living east of Baikal (Lombaert et al. 2014; Lombaert et al. 2011; Vorontsov, Blekhman 2001).

The native range of *H. axyridis* includes in the Altai Mountains, situated in north-eastern Kazakhstan and West Siberia. In some recent studies, the south-east of Kazakhstan was regarded as part of the native range (Loiseau et al. 2009; Lombaert et al. 2014; Lombaert et al. 2011). However, this is incorrect because *H. axyridis* was not recorded in the south-east of Kazakhstan in the nineteenth century or in the first half of the twentieth century. In 1968–1970 attempts to introduce *H. axyridis* from the Far East to the south-east of Kazakhstan for biological control of aphids (Savojskaja 1971) failed. Now *H. axyridis* is a common species in the south-east of Kazakhstan and was recently detected in the neighbouring region of Kyrgyzstan. Both morphological and genetic studies strongly indicate that the *H. axyridis* occurring in this region are not descendants of the released beetles, since current populations are similar to West-Siberian populations and differ from Far East populations. It is hypothesised that *H. axyridis* appeared in south-eastern Kazakhstan and Kyrgyzstan after the construction of the Turkestan-Siberian Railway, and that the beetles spread along this railway (Orlova-Bienkowskaja et al. 2015).

European part of Russia, Moldova, Ukraine and Belarus

Releases of *H. axyridis* for biological control of aphids in the Soviet Union began about 80 years ago. Attempts to introduce *H. axyridis* were made in Transcaucasia in the 1930s, in the Chernovtsy region (Ukraine) in 1964, in the Kiev region (Ukraine) in the 1960s, in Crimea in 1969, in the Tashkent region (Uzbekistan) in 1969, in Minsk (Belarus) in 1968–1970, in south-eastern Kazakhstan in 1968–1970 and in Adjara and Mtsheta region (Georgia) in 1982–1988. It is thought that none of the intentionally released populations established (Izhevsky 1990; Verizhnikova, Shylova 2013a). Established populations are thought to have originated from secondary spread eastward of *H. axyridis* from western Europe to European Russia and neighbouring countries. Established populations were detected in the Kaliningrad region (Russian enclave in the Baltic region) (Zakharov et al. 2011). Recently, *H. axyridis* has been observed spreading rapidly in the Caucasus and the south of European Russia (Orlova-Bienkowskaja 2014). Established populations have been found in Adygea, Krasnodar region, Abkhazia, the Stavropol region, Georgia (Belyakova, Reznik 2013), Crimea (Rybalchenko, personal communication), Daghestan (Ilyina, personal communication) and Rostov region (Arzanov, personal communication). Individual specimens of *H. axyridis* have also been found in the central

belt of European Russia (Belgorod and Lipetsk regions) (Orlova-Bienkowskaja 2013; Ukrainsky, Orlova-Bienkowskaja 2014), but it is unknown if the species is established there.

In 2003, several specimens of *H. axyridis* f. *spectabilis* were found in the wild in Kiev (Ukraine) and since 2007 a stable population of *H. axyridis* has existed there (Verizhnikova, Shylova 2013b). In 2009 *H. axyridis* was detected in several locations in western Ukraine, and there has been rapid expansion since, with the species now abundant all over the country (Nekrasova, Tytar 2014). The spread of *H. axyridis* has been so rapid that it has been observed to occur across a whole country before being detected: *H. axyridis* was not detected in Moldova until 2011, when it had already become common throughout the country (Timuş, Stahi 2013).

Harmonia axyridis has been recently detected in Belarus, in the Brest region. The first specimen was found in 2011, but 14 individuals were subsequently found in 2014 (Lukashuk and Ryndevich, unpublished).

Current research in Russia and the Ukraine relates to the development and effects of photoperiod on development and maturation (Belyakova, Reznik 2013; Reznik et al. 2015), genetics (Zakharov et al. 2011), morphological variability (Blekhman 2008), population dynamics (Nekrasova, Tytar 2014) and current expansion of the range in Russia and adjacent countries (Ukrainsky, Orlova-Bienkowskaja 2014). There is also considerable interest in elucidating the routes of invasion by genetic methods as has been done for many other regions (Lombaert et al. 2014). It is possible that the populations in the Caucasus and in the south of European Russia include hybrids between invasive European individuals and those introduced from the Far East for use in biological control. As the secondary range is expanding eastwards, it is predicted that it will soon reach the western border of the native range. Thus there is an opportunity to observe the consequences of interactions between native and invasive populations in West Siberia.

North America

Hamonia axyridis was repeatedly introduced throughout the twentieth century to the USA with the aim of establishing sustaining populations (Gordon 1985; Harmon et al. 2007; Koch et al. 2006b). Intentional releases of *H. axyridis* include a number of states but notably California in 1916, and multiple eastern states from 1978 to 1992 (Chapin, Brou 1991; Gordon 1985; McClure 1987; Tedders, Schaefer 1994). *Harmonia axyridis* was first detected beyond intentional release sites in the United States in 1988 in south-eastern Louisiana and eastern Mississippi (Chapin, Brou 1991). A

second, independent set of releases led to establishment in the 1980s in the Pacific Northwest (LaMana, Miller 1996; Lombaert et al. 2014; Lombaert et al. 2010). It is unknown whether all established populations resulted from these releases or from additional, accidental introductions. Nonetheless by the mid-1990s, *H. axyridis* had been found across the country with detection reports from 45 of the 48 contiguous states by 2007 (Dreistadt et al. 1995; Hesler et al. 2001; Krafur et al. 1997; LaMana, Miller 1996; Mizell III 2007) and most recently from Montana in 2009 (Foley et al. 2009) and Arizona in 2008 (Fothergill et al. 2010). *Harmonia axyridis* was first detected in Canada in 1994 (Coderre et al. 1995). Currently this species is found throughout most of North America north of Mexico with Labrador, Saskatchewan, Alaska and Wyoming the only areas where it has not yet been reported (Foley et al. 2009; Fothergill et al. 2010; Hicks et al. 2010; Koch et al. 2006b). *Harmonia axyridis* is also established and widespread in Mexico (Brown et al. 2011b).

Outside of the native range, North America has the longest history of experience with *H. axyridis* and may serve as case study of potential impacts, positive or negative, that this species may cause in other invaded areas (Koch et al. 2006a; Koch 2003; Koch, Galvan 2008). Some recent lines of research in North America are reviewed elsewhere in this paper. Briefly, research continues on attaining benefits from *H. axyridis* as a biological control agent of pests, and on understanding and minimizing adverse impacts of the invasion (pest of fruit and wine production, nuisance household invader, and impacts on native fauna).

South America

Harmonia axyridis is reported in most countries in South America (Amat-García et al. 2011; González, Kondo 2012; Koch et al. 2011; Kondo, González 2013; Nedvěd, Krejčík 2010; Saini 2004; Solano, Arcaya 2014). Based on factors such as climate and habitat, it has been predicted that *H. axyridis* would become established across broad areas of South America (Koch et al. 2006b). The invasion by *H. axyridis* has been followed intensively in Chile.

Chile

In 1998, a flightless strain of *H. axyridis* was introduced from France to central Chile for biological control in greenhouses, but those populations did not establish (Grez et al. 2010) (Table 1). In 2003, large numbers of flying *H. axyridis* associated with aphids in poplar trees were reported near the release sites. In 2008, surveys of ladybird populations in alfalfa revealed only one individual *H. axyridis* (from a sample of approximately 90,000 ladybirds), but rapid population increase was observed over the next two years in alfalfa and other habitat or crop types (Grez et al. 2010).

Harmonia axyridis is currently one of the two most abundant species in alfalfa fields in central Chile, representing 50 - 90% of ladybirds (Grez et al. 2014a; Grez et al. 2014b). It is expanding into other habitats, including native vegetation, such as sclerophyllous matorral, although at very low densities (Grez et al. 2013). *Harmonia axyridis* is spreading toward southern Chile, colonizing regions with colder climates. In contrast, few records are reported towards northern Chile, where the Atacama Desert is probably acting as a barrier for its invasion. From its original distribution in 2010, covering a range of 250 km (north to south) near Santiago, it has expanded its distribution to ~2600 km (north to south), and from sea level to >3000 m a.s.l., with a north to south spread of approximately 160 km year⁻¹. Genetic analyses suggest that current Chilean populations come from the East North America strain (Lombaert et al. 2014), representing an accidental introduction. Only f. *succinea* has been found in Chile.

Current research in Chile relates to the impacts of this species on the diversity of ladybird assemblages and abundance, including native species, in different habitats, as well as its potential damage to vineyards. There are also extensive field and laboratory experiments in progress to assess IGP and competition as mechanisms for understanding the dominance of *H. axyridis* over other ladybirds. Studies of physiological performance and life history traits under different temperature conditions in the laboratory have also been carried out (Barahona et al. submitted). The invasion process is being systematically tracked through surveys coordinated by Grez and Zaviezo, including data from the Surveillance Department of the National Agrarian and Livestock Service (SAG), and from media and social networks specifically developed for this purpose (web page: www.chinita-arlequin.com; Facebook: <https://www.facebook.com/chinita.arlequin>; twitter: <https://twitter.com/chinitaarlequin>). The high interest and participation of citizens through these media suggest that the presence of *H. axyridis* is of growing concern to Chileans.

Spatial distribution models are also being developed to forecast the future distribution of *H. axyridis* in the country. Chile is a long (4329 km) and narrow (180 km on average) country, isolated by the Atacama Desert to the north, the Pacific Ocean to the west, and the Andes mountain range to the east. The Chilean environment has strong latitudinal gradients in abiotic conditions, from the most arid desert in the world in the North, followed by a semi-arid region, Mediterranean type ecosystems in central Chile, the temperate rainforests toward the South, and the southernmost sub-Antarctic ecosystems (Luebert, Plischoff 2006). This gradient offers a unique opportunity to study how climate modulates the distribution and abundance of this invasive alien species. Also, along this

gradient, a highly diverse native fauna of coccinellids (~115 species) (González 2008) offers a unique opportunity to evaluate the possible changes of native communities after *H. axyridis* arrival.

The main gaps in knowledge in Chile relate to natural enemies and possible control or containing methods. The only observed natural enemy thus far is the parasitoid *D. coccinellae*. Also, there are many unanswered questions in relation to the ecology and biology of *H. axyridis* during summer time, when this species disappears from crops, apparently estivating or migrating to other, as yet undetermined, locations. Its role as a biological control agent of aphids and other pests is still unknown.

Europe

A network of scientists was established through a working group (Benefits and Risks of Exotic Biological Control Agents) of the Western-Palearctic Regional Section of the International Organisation of Biological Control in response to the rapid spread of *H. axyridis* throughout Europe. Consequently there has been ongoing collaborations to map the distribution throughout Europe (Brown et al. 2008a) and so here we present information reflecting this activity on a country by country basis. There is considerable scope to enlarge the network and research collaborations and we look forward to doing so in the coming years.

Austria

Harmonia axyridis was first recorded in eastern Austria in 2006 (Rabitsch, Schuh 2006) and soon reported from other provinces, where it may have been present but undetected for some time. It was never imported or sold as a biological control agent in the country and the arrival via natural spread from north-western Europe is most likely. After almost a decade, however, the species has been found in all federal provinces and there is no doubt that it has become the most abundant ladybird species in Austria.

Unfortunately, no systematic surveillance or monitoring data are available to trace the expansion within Austria or to document any impact on native biodiversity. Personal observations from entomologists, however, indicate that the species has strongly increased in abundance and range over the last decade. The establishment and use of citizen science initiatives in gathering information are currently under development.

Belgium

In Belgium, *H. axyridis* was first applied as a biological control agent in 1997. The species was first observed in the wild in 2001 (Adriaens et al. 2003) and this led to the end of its commercial use in Belgium. In less than five years the species invaded the entire country, its distribution covering all Belgian ecoregions (Adriaens et al. 2008). The area of occupancy showed an average rate of increase of 189% (5000 km²) per year from 10.000 km² in 2002 to 31.000 km² in 2006. *Harmonia axyridis* has become the most abundant species in (semi-)natural systems and also dominates the aphidophagous guilds in certain agro-ecosystems (Vandereycken et al. 2013).

At the time of arrival and spread of *H. axyridis* in Belgium a country-wide validated citizen-science survey (Gardiner et al. 2012) focussing on the ecology and distribution of 40 native ladybird species was running. This allowed almost real-time monitoring of this insect invasion as well as detailed analysis of niche overlap with native species (Adriaens et al. 2008). Future research on the decline of native coccinellids should also consider factors such as land use change, climate change, habitat quality and effects of pesticides as potential contributors to this phenomenon.

Czech Republic

Although *H. axyridis* was released in the Czech Republic for protection of hops in the early 2000s, it did not establish. The first occurrence of the invasive population dates back to 2006. In 2007, it occurred near the western borders of the country and in a few cities, but in 2008 it was found in many cities and towns including the eastern-most (Špyňar 2008). In 2009, the species could be found in most anthropogenic-altered landscapes, but not in remote areas such as mountains over 1100 m, continuous forests and closed military zones. Mass infestation of houses during autumn migrations raised public awareness and a few cases of biting and allergic reactions were reported. Unintended transfers of single specimens over hundreds of kilometres were recorded (Nedvěd 2014).

Denmark

Harmonia axyridis was first recorded in Copenhagen 2006, and presumed to have arrived from Germany (Brown et al. 2008a; Pedersen et al. 2008). In the following years the beetle spread within the Greater Copenhagen area, becoming established across the island of Zealand. By 2007/8, its spread continued westwards to the islands of Lolland, Langeland, Ærø, and in the east to Bornholm (perhaps arriving from Poland or Germany) where populations are now established (Howe 2015; Howe et al. 2015; Steenberg, Harding 2009b; Steenberg et al. 2009). By 2008, *H. axyridis* had reached Jutland in the far west of Denmark (Steenberg et al. 2009). The most northern record to date (latitude 57° N) was from produce in a supermarket, highlighting the potential for spread

through anthropogenic pathways. Data from Zealand indicates stable populations with *H. axyridis* dominating ladybird assemblages where established. Greatest densities occur within urban areas of central Copenhagen (Ravn and Howe, unpublished data).

It is interesting to note that the distribution of *H. axyridis* in Denmark is limited in the north of the country. Indeed, following ten years' of establishment within Denmark, aside from the aforementioned supermarket record there are no records of *H. axyridis* north of 56 ° N (Ravn and Howe, unpublished data), which corresponds with a lack of records from Scotland (UK). Whether this represents a true limit to the northern distribution attributable to present climatic conditions requires further research, particularly in relation to future predictions of spread based on expected climate warming.

France

In France, *H. axyridis* was first introduced for biological control in 1982 but very few records of feral populations exist until 2004 (Coutanceau 2006a), when the species started to spread across the country from the north, close to its probable introduction point in Belgium (Adriaens et al. 2003). Maps of first observations of *H. axyridis* at particular locations in France suggest a heterogeneous process of diffusion, with some regions rapidly colonized whereas in others there is a lag in invasion, or *H. axyridis* remains absent (see maps in http://vinc.ternois.pagesperso-orange.fr/cote_nature/Harmonia_axyridis). First observation records have been used in sophisticated modelling approaches to make inferences about the relative impact of various environmental and anthropogenic factors on that spread (Veran et al. 2015) and confirmed that the rate of colonization of *H. axyridis* in France was heterogeneous in time and space. Anthropogenic factors explained more variation of the diffusion process than environmental ones. The relative surface of urbanized area was the major anthropogenic factor increasing the probability of colonization. More specifically, low urban densities, corresponding to rural areas, represented unfavourable habitats. Finally, average summer temperature was the main environmental factor affecting colonization, with negative impact for both high and low values.

IAS expanding their range provide unique opportunities to explore the effect of spatial spread on life-history traits, making it possible to test for a spatial arrangement of dispersal abilities along the expanding range (Phillips et al. 2007; Purse et al. 2014). Moreover, the question of the evolution of dispersal capacity in invasive populations is highly relevant, because it may accelerate spread, with serious applied and theoretical consequences (Phillips et al. 2010; Travis, Dytham 2002). Using

controlled experiments in laboratory conditions, clear evidence was found of a strong, rapid increase in flight speed with range expansion of *H. axyridis* from the core of the invaded area in Western Europe (Belgium) to the front of this invaded area in South and Western parts of France (Lombaert et al. 2014). This shift towards a higher flying speed at the invasion front was remarkably rapid, as it was demonstrated after only eight years of expansion, corresponding to about 16 generations (Koch 2003).

Using population genetic approaches, it has been shown that the *H. axyridis* sampled across France (see below for exceptions) belonged to a single genetic unit that has invaded the Western part of Europe (Lombaert et al. 2010; Lombaert et al. 2011). This invasive population bears traces of genetic admixture between an eastern North American wild source, which was found to have served as a “bridgehead source” for many worldwide invasive outbreaks, and a biological control strain used in Europe (Lombaert et al. 2010). Experimental studies in the laboratory have investigated the phenotypic impacts of such genetic admixture (Facon et al. 2011; Tayeh et al. 2013; Tayeh et al. 2012; Tayeh et al. 2015; Turgeon et al. 2011). Biological control individuals were found to display classic *r*-selected traits with a shorter lifespan and an earlier egg production when compared to native and US invasive individuals. European invasive individuals have shown phenotypic traces of the genetic admixture between US invasive and biological control individuals. For most traits (such as age at the start of reproduction, total adult and reproductive lifespan), European invasive individuals displayed intermediate values between both parents. Thus genetic admixture has had a long-lasting effect in the wild by shaping the life-history strategy of the European invasive individuals of *H. axyridis*. In France, and more generally Western Europe, the exact role of admixture with the European biological control strain in the process of invasion remains poorly known. The single eastern North American origin of one South African and two South American outbreaks indicates that the genetic admixture observed in Western Europe is not required for an eastern North American propagule to establish and start an invasive population in diverse ecological contexts (Lombaert et al. 2014). It seems therefore unlikely that admixture in Western Europe has radically changed the outcome from failed to successful invasion. Such admixture has probably simply modulated the rate (or impact) of an invasion process that would have been successful anyway.

In addition to the single vast admixed population which invaded Western Europe, a genetically distinct population established in South East of France has been identified (Lombaert et al. 2014). This population was first observed in 2005, and it appeared to have originated exclusively from the European biological control strain introduced into Europe from 1982. This is surprising, at first sight,

because European biological control individuals have long been thought to be unable to survive in the wild (Ferran et al. 1997). In support of this, it is apparent that this distinct population does not seem to have expanded spatially, unlike most of the other *H. axyridis* outbreaks known all over the world. This locally established population might attest, however, to the ability of the European biological control strain to found small overwintering populations in the wild, in areas with clement winters.

Germany

In 2000 the first German record of *H. axyridis* in the wild was reported from Frankfurt City in 2000. This was followed two years later by reports of *H. axyridis*, in higher abundance than the first records, from Frankfurt region and also Hamburg (Klausnitzer 2002). It has spread rapidly into other regions across Germany and is now considered established throughout Germany, although there has been no nation-wide monitoring to confirm this assumption. In South-West Germany near Frankfurt peak population densities were observed in early autumn until 2009 and also in 2012 but there seems to have been declines in numbers since then. However, this information is not derived from systematic monitoring.

Several German populations from 2008 and 2009 were examined for the presence of antagonistic microorganisms (bacteria, fungi, microsporidia) and invertebrate parasites (Herz, Kleespiel 2012).

Italy

In Italy, *H. axyridis* was released in protected crops as a biological control agent from 1995 to 1999. However, the use of *H. axyridis* for biological control was suspended because concerns over adverse impacts became apparent. The first occurrence of establishment was in 2006 in Piedmont followed by subsequent observations in Emilia-Romagna in 2008 and rapid establishment in 16 out of 20 Italian regions with particularly widespread occurrences across northern and central Italy (Burgio et al. 2008b; Cornacchia, Nardi 2012). The origin of the populations is unknown; individuals could be either offspring of those released *in situ* or immigrants from other parts of Italy, France or Switzerland (Burgio et al. 2008b).

Research studies in Italy have focused on *H. axyridis* biology (Bazzocchi et al. 2004) and specifically life table parameters. Research on the occurrence of parasitoids attacking *H. axyridis* is ongoing (Francati 2015a, b; Rondoni et al. 2013a; Rondoni et al. 2013b). Other studies are focusing on the development of a liver-based artificial diet, which could assist in maintenance of cultures of adult *H.*

axyridis (Sighinolfi et al. 2008), and on the susceptibility of *H. axyridis* larvae to lambda-cyhalothrin insecticide (Benelli et al. 2015).

Norway

Harmonia axyridis was assessed as a potential biological control agent for use in Norwegian greenhouses in 2001 (Statens landbrukstilsyn 2001). It was concluded that *H. axyridis* might become established outdoors and thereby pose a risk to the environment. The assessment was therefore negative with respect to import and commercial use of *H. axyridis* in Norway.

The first record of *H. axyridis* in Norway was in Oslo in 2006, the adult female f. *succinea* arrived as a stowaway on horticultural plants, *Thuja* sp. (Cupressaceae) imported from the Netherlands to Norway (Staverløkk 2006). In late 2007 and throughout 2008 several adults were found indoors and outdoors at a number of locations in the urban and suburban areas of Oslo (Sæthre et al. 2010).

Observations in areas some distance from Oslo, such as Tvedestrand in Aust-Agder County (2008), Våle in Vestfold County (2008) and Trondheim (2009) (the latter about 600 km north of Oslo) revealed further spread or separate introduction to new areas (Sæthre et al. 2010). Repeated introductions (probably on imported plants) are likely to be the most important factor for introduction of the species to new areas in Norway. Natural geographic barriers and long distances within Norway limit the species possibilities for rapid natural dispersal. However, anthropogenic spread facilitates dispersal and of particular note was the occurrence of between 2000 and 3000 adult *H. axyridis* in a cargo of timber imported to Åndalsnes (Møre og Romsdal County) in March 2008 from Pennsylvania, USA. Some specimens were also recorded at Snåsa (Nord-Trøndelag County) on the imported timber which was transported from Åndalsnes. According to the importing timber company, actions had been taken to eradicate the beetles.

In late 2008 a website was launched to engage the public in submitting observations on-line and this has made a major contribution to documenting the distribution of *H. axyridis* in Norway (Sæthre et al. 2010). Records are available at <http://www.artsportalen.artsdatabanken.no/#/Harmonia+axyridis/7468>. In autumn 2015 high numbers of *H. axyridis* were reported from overwintering aggregations across the city of Oslo. However, so far, the distribution of *H. axyridis* in Norway appears to be limited to urban and suburban regions, and it has to date not been recorded or reported in commercial crops or in natural

habitats. Further studies on the biology, ecology, cold tolerance and winter survival will contribute to better predictions of the dispersal and establishment potential of *H. axyridis* in Scandinavia.

Slovakia

Harmonia axyridis was first recorded in Slovakia in 2008 (Majzlan 2008). It arrived as an unintentional introduction by secondary spread, following the spread across Austria (Rabitsch, Schuh 2006), Poland (Przewozny et al. 2007) and the Czech Republic (Špyňar 2008). There was less than one year between the first record of establishment and widespread occurrence of *H. axyridis*. By the end of 2009, it was recorded across Slovakia, and by the end of 2012 it occurred in numerous habitats, particularly gardens, orchards and urban areas. The records from 2008–2012 document the invasion clearly (Panigaj et al. 2014). The distribution and time sequence of the records support the maximum rate of the spread of *H. axyridis* to be approximately 200 km year⁻¹, the spread being accelerated by human movement (Panigaj et al. 2014). The local topography played a crucial role in the spread: 47% of the records of the coccinellid were from lowlands (94–200 m), 36% from low hilly areas (200–400 m), 11% from moderate altitudes (400–600 m) and only 6% from higher areas (600–1250 m a.s.l.) (Panigaj et al. 2014). Despite great efforts in 2013 and 2014, only a single specimen of *H. axyridis* was recorded from altitudes above 1000 m.

Wildlife records in Slovakia are mostly shared through popular naturalist's web pages (e.g. www.nahuby.sk, www.fotonet.sk) which also provide information about *H. axyridis*. The Facebook page Lienky Slovenska (Ladybirds of Slovakia) was launched in 2015 to encourage the public to take part in ladybird surveys focussing on *H. axyridis*. The international cooperation with scientists and collaboration with volunteers within the home country is critical for the progress in the further research of *H. axyridis*.

Switzerland

The first *H. axyridis* adult was found in 2004 in Basel (Klausnitzer 2002), but it was only in 2006 that establishment was confirmed in several locations (Eschen et al. 2007). From then, the ladybird rapidly invaded all areas of low and middle altitudes within two years. A long-term inventory was initiated in the northwest of Switzerland in 2006 to record the impact of *H. axyridis* on native ladybirds (Eschen et al. 2007). Ladybird populations were monitored using standardised sampling methods at 45 sites: 15 broadleaved hedgerows, 15 meadows and 15 conifer sites several times a year. The monitoring was interrupted only in 2014 but has since restarted. Other surveys were made in other habitats on an irregular basis, in particular on urban trees. Since 2008, *H. axyridis* has

become by far the most abundant ladybird on broad-leaved shrubs and trees, accounting for 60 to 80% of all ladybirds collected throughout the year. In contrast, in meadows and on conifers, *H. axyridis* still remains rather uncommon, except on some specific plants such as nettle (Kenis and Eschen, unpublished data).

Further studies in Switzerland have also focused on the impact of *H. axyridis* on native ladybird populations. The occurrence of intraguild predation in the field using polymerase chain reaction (PCR) to identify target prey DNA within a predator's gut has been investigated (Aebi et al. 2011). IGP between *H. axyridis* and eleven native non-target European ladybirds in laboratory experiments has been studied (Katsanis 2011).

The Netherlands

Harmonia axyridis was first released as a biological control agent in greenhouses in 1995, on outdoor crops, arboriculture and in urban areas since 1996, largely for the control of aphid pests (Cuppen et al. 2004). Releases were stopped by the end of 2003. *Harmonia axyridis* has established at some sites very rapidly with the first report of *H. axyridis* in the wild in 2002 in Groesbeek, Gelderland (as a pupa on a leaf of *Hedera helix* L. (Araliaceae); Cuppen et al. 2004) followed by a specimen in Rotterdam, South Holland in 2003, and around the same time another specimen was collected in Reimerswaal, Zeeland. In July 2003 the first adult *H. axyridis* was collected on a lighted white sheet at night at a nature reserve (De Kaaistoep, North Brabant) (Cuppen et al. 2004; van Wielink, Spijkers 2013). In this locality, up to 2014, 6516 specimens have been collected using light, 71.9% of which were males (yearly range between 65 and 80%) (van Wielink, unpublished data).

From 2004 onwards, numbers of *H. axyridis* started to rise dramatically and monitoring efforts were increased (via <http://www.knnv.nl>, <http://www.stippen.nl> and <http://www.waarneming.nl>). While up until 2004 (mainly) the south part of the country was invaded, by 2007 more than 2000 records had been received covering the entire country, including the Wadden Islands (Brown et al. 2008a). *Harmonia axyridis* is now widely distributed and the dominant ladybird species, predominantly on trees and shrubs but also on herbs in urban and anthropogenic habitats, it is less abundant in (semi-) natural areas such as heathland and grasslands. In agricultural areas *H. axyridis* is considered the dominant ladybird in corn, but not in cereals. During the early years of invasion, until 2010, local abundance was particularly high during early summer and autumn but this has not been so apparent in recent years.

Research in the Netherlands has focused on natural enemies (Haelewaters et al. 2012; Raak-van den Berg et al. 2014; Sloggett 2010) and the high overwintering survival of *H. axyridis* (70.8-88.2%) (Raak-van den Berg et al. 2012). The latter study found that overwintering survival was higher (1) at sheltered places compared to exposed sites and (2) when ladybirds were overwintering at south-western sides of buildings. As a comparison, winter survival of *A. bipunctata* in the Netherlands is 17-78% (Brakefield 1985).

United Kingdom

The UK has a long history of involvement of volunteers in gathering information on wildlife through biological recording (Pocock et al. 2015; Roy et al. 2015a). The Coccinellidae Recording Scheme (hosted by the Biological Records Centre which is part of the Centre for Ecology & Hydrology) was established in 1971 (Roy et al. 2011a). The legacy of ladybird recording in the UK provided a unique dataset through which to explore the impacts of *H. axyridis* on other ladybirds. *Harmonia axyridis* arrived in Britain through dispersal and introduction events from regions (mainly Europe) in which it was deliberately released as a biological control agent (Brown et al. 2008b). *Harmonia axyridis* was first recorded in the UK in 2003 (Roy et al. 2012d) and was established by 2005. An on-line survey (www.ladybird-survey.org) was launched to monitor the spread of *H. axyridis* while promoting the continued recording of other ladybirds. Tens of thousands of people have provided records of *H. axyridis* and other species of ladybirds (Roy et al. 2015b), providing an invaluable large-scale and long-term dataset which has been used to explore the invasion process and concomitantly trends in the distribution of other ladybirds (Comont et al. 2014a; Comont et al. 2012; Purse et al. 2014; Roy et al. 2012c). For example, declines in the distribution of seven (of eight assessed) native species of ladybird have been demonstrated, and correlated with the arrival of *H. axyridis*, using the records collated through the UK Ladybird Survey (Roy et al. 2012c).

The rapid spread of *H. axyridis* - more than 100 km per year across the UK (Brown et al. 2008b) - has been attributed to its high natural dispersal capability through both flight (Jeffries et al. 2013; Lombaert et al. 2014; Maes et al. 2014) and anthropogenic transport (Brown et al. 2011b). A number of factors are considered to have contributed to the successful establishment and dominance of this polymorphic species within aphidophagous guilds across the UK, including high reproductive capacity, intra-guild predation, eurytopic nature and high resistance to natural enemies within the invaded range (Roy, Brown 2015b).

Considerable attention has been given to experimental research (Comont et al. 2014b) and systematic field surveys (Brown et al. 2011a) to further understanding of the interactions between *H. axyridis* and other species. Future work will reflect the opportunities presented by *H. axyridis* to explore the complex and dynamic role of natural enemy interactions in the invasion process through community (network) approaches (Roy, Lawson Handley 2012). The role of citizen scientists in gathering information on species interactions is considered an important component of this research.

Africa

In North Africa, *H. axyridis* was introduced for biological control use in Tunisia around 1990 and in Egypt before 2000 (El-Arnaouty et al. 2000). It is thought to be established in limited areas of the latter, but not the former (Brown et al. 2011b).

According to unpublished records and misplaced voucher specimens discovered only in mid-2015, *H. axyridis* was intentionally released in South Africa around 1980 (Stals unpublished). This contradicts the previously published view that the species had never been intentionally introduced to this country (Stals and Prinsloo 2007, Stals 2010). The beetles were sourced from the USA, apparently originating from Japan. The release was made in an attempt to control the black pine aphid, *Cinara cronartii* Tissot & Pepper, in the Sabie area, in the present-day Mpumalanga Province. In the newly found records, a later, but undated, entry notes that the species had not established. No other information is presently available. It seems probable that this release failed, since no specimens of *H. axyridis* collected in southern Africa before 2001 are present in any public insect collection in South Africa.

Unaware of the above, (Stals, Prinsloo 2007) announced that *H. axyridis* was first recorded in South Africa in 2004, in the Western Cape Province. However, museum records later revealed that adults and immature stages had been found in the Cape Town area as early as 2001 (Stals 2010). This remains the earliest known date and location of establishment in southern Africa. Sabie, the 1980s release point, is c. 1,800 km away from Cape Town. *Harmonia axyridis* specimens from the Sabie area were only recorded in the austral summer of 2008 (Stals unpublished), when the contemporary spread of the invader into Mpumalanga was already well underway (Stals 2010).

The introduction pathway of the contemporary invaders in South Africa is unknown, but all analysed South African populations originated from eastern North America (Lombaert et al. 2014; Lombaert

et al. 2010). Introduction was likely unintentional, but it is unknown whether there was more than one introduction to South Africa, disregarding the almost certainly failed introduction of the 1980s.

The only coordinated data-gathering initiatives for insects in South Africa are for Lepidoptera and Neuroptera. Nonetheless, recording of the range expansion through southern Africa mainly depends upon volunteer contributions. Until 2010, contributions were encouraged through sustained calls in popular media (Stals 2008; Stals 2010). Since 2011, the citizen science web application iSpot (Silvertown et al. 2015) has been exploited as a recording platform for southern Africa and became the source of many high-quality observations of *H. axyridis*. An expert Coleoptera taxonomist assesses all contributions, querying contributors where necessary.

Harmonia axyridis rapidly spread widely through much of South Africa (Stals 2010; unpublished). It seems established largely in cooler and more mesic parts of the country, viz. the south-western, southern and interior-eastern Cape regions, and the more northern eastern and east-central areas, with few records from the semi-arid western and west-central reaches or the hot northern regions. No records have as yet come from the subtropical eastern coastal belt. The invader is well established and commonly encountered in the Fynbos and Grassland Biomes; established and not infrequently encountered in the Savanna Biome; but infrequently reported from other biomes. In South Africa, the majority of records come from urban and rural gardens or dwellings, and from agricultural land. Other southern African countries with records of *H. axyridis* are Lesotho (first record June 2008, Stals 2010) and Swaziland (first record November 2013, Stals unpublished), and the ladybird has likely established in both these countries.

In the rest of sub-Saharan Africa, *H. axyridis* has only been recorded in Kenya (Nedvěd et al. 2011) and Tanzania (Nedvěd, Háva 2016). In Kenya, a population was discovered in December 2010 at a coastal holiday centre and may represent an established population (Nedvěd et al. 2011). In Tanzania, only two *H. axyridis* individuals were found at a beach resort in Zanzibar in April 2014 and may represent a transient introduction with no establishment. These occurrences suggest that the invasion of tropical Africa is possible. The origin of the beetles in East Africa has not yet been investigated and all examined specimens were of the f. *succinea*.

Research on *H. axyridis* in South Africa is scarce but it is recognised that there is a need to assess its biology and ecology in order to evaluate its impact on native communities and agroecosystems. Researchers from the Centre for Invasion Biology, Stellenbosch University, are focusing on the

thermal biology and life history of *H. axyridis* in South Africa and compare this species' traits to those of native aphidophagous ladybirds. This information can highlight characteristics that promote the invasiveness of *H. axyridis* in southern Africa and provide data for modelling its potential spread within and beyond borders (Shinner 2014). The study of its behavioural responses and adaptation to climate variation will remain a focus in the years to come as well as modes and mechanisms of introduction and range expansion investigated using molecular techniques.

However, many research aspects that are key for evaluating the establishment, spread and impact of *H. axyridis* are lacking. No systematic field surveys are taking place and therefore the invader's abundance across habitat types is also unknown. More importantly, data on the native ladybird communities (species richness and abundance) occurring in areas with and without *H. axyridis* are not being collected and the impacts on these communities are thus unknown. Of overriding importance may be the complete lack of baseline data prior to this invasion; it is possible that no readily comparable and uninvaded habitats may remain for urgent collection of baseline information. In addition, the taxonomy and phylogeny of the native Coccinellini need to be examined, in particular those African species currently placed in the genus *Harmonia*.

Under South African national legislation, *H. axyridis* is a Category 1b Listed Invasive Species (Department of Environmental Affairs South Africa 2014a), which in terms of the National Environmental Management: Biodiversity Act, 2004, legally is a "species which must be controlled" (Department of Environmental Affairs South Africa 2014b). How such control is to be achieved is unclear at present.

Failure of Harmonia axyridis to establish in some regions

Harmonia axyridis is much more commonly and widely distributed in the northern than in the southern hemisphere, perhaps unsurprising given its Asian origin. While *H. axyridis* distribution extends to boreal regions in its native range, its invasive distribution in the north and south of Europe, and also in northern Canada and Alaska, is more limited and suggests that climatic factors may be important in limiting the spread of this species. Globally, there are very few records from tropical regions (23.4°N–23.4°S); where it has been reported from the tropics it is not widely spread (e.g., Colombia, Venezuela, Kenya, Tanzania). In both South America and Africa, limits to its distribution apparently include warm tropical but also arid environments.

In Europe there are some habitats and regions that appear to be resistant to the establishment of *H. axyridis*. Some examples are the limited evidence of establishment in Greece (Kontodimas et al. 2008), Turkey (Bukejs, Telnov 2015), Spain, including the Canary Islands (Goldarazena, Calvo 2007; Jacas et al. 2006; Pons et al. 2015), Bosnia and Herzegovina (Kulijer 2010), Portugal, including the Madeira and Azores archipelagos (Garcia 1986; Soares et al. 2008), Northern Ireland (Murchie et al. 2008), the Republic of Ireland (<http://www.invasivespeciesireland.com>), and Réunion (Quilici, pers. com). It is as important to consider the regions in which *H. axyridis* has failed to establish as those in which it has succeeded. Only recently has consideration been given to understanding invasion failures within the context of invasion processes (Zenni, Nuñez 2013). The factors limiting invasion by *H. axyridis* in some geographic areas are worthy of exploration and could potentially provide insights into whether the southern and northern European ecosystems are more resilient to invasion than other parts of Europe and beyond. Investigation of equivalent patterns on the southern African subcontinent would also be informative.

Records of successful breeding by *H. axyridis* are very limited in Scotland (Roy, Brown 2015b) and although climatic conditions are not thought to have been a barrier to the colonization and spread of *H. axyridis* in southern Britain, it is possible that climate has limited its abundance not only in northern England and Scotland (Brown et al. 2008b; Roy, Brown 2015b), but also Denmark (Steenberg, Harding 2009b) and further north throughout Scandinavia. The combination of lower temperatures and higher precipitation in Scotland compared to England could be limiting the distribution of *H. axyridis* within Scotland. The Orkney and Shetland Islands are considered climatically unsuitable for *H. axyridis* (Poutsma et al. 2008), and in support of this there have only been occasional records of adults, arriving on produce imported from the mainland, from these northern islands (Ribbands et al. 2009). There are no records of immature stages of *H. axyridis*, or other ladybird species, on these islands.

Although thousands of individuals of both f. *succinea* and f. *conspicua* have been released in the Azores (Garcia 1986; Schanderl et al. 1992), *H. axyridis* has not become established there (Evans et al. 2011; Soares et al. 2008). It has been hypothesised that prey features related to local plant habitats and landscape structure, together with the ladybird characteristics in terms of body size, might explain why the invader is absent (Hemptinne et al. 2012). The coastal terrestrial habitats that form 9% of the landscape are the richest in terms of food resources for aphidophagous ladybirds. It is predicted that other large and medium-sized species of ladybirds (*Coccinella undecimpunctata* L.), which have requirement for high prey consumption, will also decline because of shortage of prey

(Borges et al. 2006; Cabral et al. 2006; Sebastião et al. 2015; Soares et al. 2001) for example *C. septempunctata* is considered extinct from the Azores. The Azorean communities are nowadays dominated by minute Scymnini species and larger species are no longer recorded. Despite the competitive advantage of *H. axyridis* against the native *C. undecimpunctata* (Felix, Soares 2004; Nória et al. 2008), it is apparent that Azorean habitat characteristics and the high feeding rate of *H. axyridis* are hampering its invasion.

Comparison of traits

There have been a number of studies examining the influence of life-history traits on invasion and *H. axyridis* is no exception (Comont et al. 2014a; Comont et al. 2012). The traits databases compiled for these studies provides a rare opportunity to explore variation in life-history traits between localities (native and invaded) around the world. Extending and combining traits databases to a global scale will provide intriguing insights. The invasion process and ecological attributes (Table 1) and the compilation of life history traits of *H. axyridis* (Table 2) represents the start of this process and highlights gaps in understanding. However, qualitative comparisons of the traits of *H. axyridis* across the invaded range and with the native range reveals patterns that are worthy of further investigation.

Habitat

The habitat of *H. axyridis* is wide and although poorly documented in many regions the exploitative and opportunistic nature of this species is evident (Table 1). During early stages of invasion it has been noted that *H. axyridis* is more prevalent within urban and agricultural landscapes than in semi-natural landscapes (Brown et al. 2008b; Grez et al. 2014a; Grez et al. 2014b). *Harmonia axyridis* is common within gardens and parks throughout the year and its presence in agricultural fields, orchards or vineyards has been documented globally and local damage due to quality loss of fruits is known (Koch et al. 2004a). Across Europe and Japan *H. axyridis* is commonly associated with trees and shrubs. For example in urban areas in Europe it is the most abundant ladybird on lime (*Tilia* spp.) and maple trees (*Acer* spp.) but is also frequently found on Scots pine, *Pinus sylvestris* L. (Pinaceae). The habitats of *H. axyridis* in Japan are disturbed areas such as agricultural fields, orchards, parks, residential yards and gardens (Osawa 2011) and *H. axyridis* is generally uncommon in natural forests (Osawa 2011).

In winter *H. axyridis* has a propensity to aggregate in buildings. Across Europe adults usually start aggregating in October and leave overwintering sites in April. The aggregating behaviour has been

shown to depend on two blends of long chain hydrocarbon molecules, one leading conspecifics towards aggregation sites and the other ensuring cohesion of the aggregations (Durieux et al. 2012). These findings, with the identification of a volatile sex pheromone in female *H. axyridis* (Verheggen et al. 2007), might offer some potential in the development of specific control methods for *H. axyridis*.

It is critical that we have a better understanding of the habitat preference and suitability of *H. axyridis*; this is especially relevant when facing current global environmental change, especially urbanization, agricultural intensification and climate change which will undoubtedly affect the spread and distribution of *H. axyridis*. Habitat suitability could be estimated using Species Distribution Models based on fine-scale records on the presence/absence or densities of *H. axyridis* populations. Such results would have wide relevance to invasion biology.

Diet breadth

The wide diet breadth of *H. axyridis* is evident from Table 1 and, as already stated, includes pest and non-pest insects but also fruits. Perhaps of most interest, and relevance to understanding the threat posed by this species to biodiversity, is the range of non-pest insects consumed and associated population-level effects. Much research has focussed on interactions between *H. axyridis* and other coccinellids (Pell et al. 2008) but there is a need for further work on other taxonomic groups for example expanding the research on butterflies such as the studies on the monarch butterfly *Danaus plexippus* (L.) (Koch et al. 2006a; Koch et al. 2003). Furthermore it is critical that future research considers the population-level effects of *H. axyridis* on different species and implications for ecosystem function. Current studies have not clearly documented the diet preference, adaptability and feeding efficiency of *H. axyridis*, which are essential for assessing its potential impact in recipient ecosystems.

Thermal tolerance

Only a few studies, representing populations of *H. axyridis* from a few regions, have considered thermal tolerance (Table 2) (Koch et al. 2004b). Of particular importance would be to measure lower and upper temperature tolerance of several life stages using ecologically relevant conditions for the population examined. Indeed to build Species Distribution Models, thermal tolerance needs to be carefully measured in the laboratory so that the reaction norm of physiological/behavioural performance as a function of ambient temperature can be established. However, *H. axyridis* usually overwinters in shelters and aggregations and cold tolerance measurements should be interpreted in

regards to ecologically relevant conditions. Indeed *H. axyridis* can be found at a range of altitudes: in Chile individuals have been found at >3000 m a.s.l. while in central Europe it is regularly present from the eastern lowlands (114 m a.s.l.) to montane/subalpine forests (approx. 1600 m a.s.l., unpubl. data) in the Alps, with one isolated record of a probable wind-drifted individual at 2280 m a.s.l. in Carinthia.

Although laboratory studies show that invasive populations of *H. axyridis* do not survive at 34°C (Benelli et al. 2015), the species has been documented in Kenya (Nedvěd et al. 2011) and Tanzania (Nedvěd, Háva 2015) but there is no evidence that this species is abundant in the tropics. In addition, the wide spread and occurrence of *H. axyridis* in mesic cool climates or pockets of urban gardens within drier environments in its novel range suggests that climatic factors other than temperature may also play a role in shaping its distribution.

Desiccation resistance (Nedved, Kalushkov 2011) and the potential cross-effects of temperature and humidity on temperature limits may provide useful information for modelling its future distribution in face of climate change (Hoffmann et al. 2013). Perhaps more importantly, the study of the plasticity and evolutionary adaptation of tolerance limits in *H. axyridis* using both native and invasive populations would provide insights into its adaptive capacity to future climatic challenges. The effect of different climate regimes on the body weight and fat body content of *H. axyridis* and *C. septempunctata* has been the focus of laboratory studies in Germany (Krengel et al. 2012). This study concluded that that *C. septempunctata* has life history adaptations that would confer an advantage over *H. axyridis* at elevated temperatures.

Reproductive potential

A number of traits (voltinism, fecundity and egg hatching success) reveal the high reproductive potential of *H. axyridis* (Table 2). In the native range, *H. axyridis* is generally considered to have two generations per year, although it can have three and occasionally up to eight generations in some years and localities (Osawa 2011). In Kyoto, in the centre of Japan, the overwintering adults mate and lay eggs in spring (Osawa 2000). The adults of the first generation emerge in mid May to June (Osawa 2000). In mid summer, the beetles aestivate in small groups in leaf-shelters on trees and the behaviour is regarded to be an adaptation to high temperature (Toda, Sakuratani 2006). In autumn, *H. axyridis* adults fly towards overwintering sites, white or pale objects on hilltops or valleys, where they aggregate from early November (Obata 1986).

Within much of the invaded range, *H. axyridis* only achieves two generations per year, whereas in the native range up to eight generations have been observed. Increases in global temperatures could facilitate an increase in voltinism, and corresponding increase in abundance across the invaded range in the future. On the contrary, increase in temperatures may limit its reproduction due to the high egg mortality observed at temperatures above 30°C (Table 2).

We suggest that the three traits related to population demography and thus invasiveness, namely voltinism, fecundity and hatching success, need to be recorded in a standardised unit so that population dynamics can be estimated from matrix population models. Furthermore, the trade-offs between, and rates of adaptation in, life-history traits are of intrinsic theoretical value (e.g. the trade-off between these three demographic traits among populations in different environments) and colour polymorphism of *H. axyridis* (Table 2) is also an interesting model system in population genetics. Overall, we suggest that the strength of biotic interactions centralised around *H. axyridis* (i.e. diet preference and foraging efficiency) and the three demographic traits are key variables for monitoring/assessing its performance and impact in invaded ecosystems.

The reproduction success of individual females has been measured in variable ways in *H. axyridis* according to regions (Table 2). Usually reproduction is measured as lifetime fecundity (total number of eggs laid by female), but this requires very long breeding and observation, since many females of *H. axyridis* live and reproduce for over four months (Awad et al. 2013). Alternatively, fecundity during the first month of reproduction (beginning by the day of the first egg laying, not including pre-oviposition period) is used. Both measures require feeding the females *ad libitum* with a suitable prey. Widely used types of prey that may become a standard are the pea aphid *Acyrtosiphon pisum* (Harris) (Hemiptera: Aphididae) and frozen eggs of the flour moth *E. kuehniella* (Berkvens et al. 2010a; Kögel et al. 2012). Other measures of reproductive performance could include daily fecundity (after a standard time period for example two weeks after the start of egg laying) and cluster size (again after a standard time period and perhaps an average over two weeks).

Fertility is defined as the product of fecundity and hatching rate. Hatching rate (percentage of eggs that hatch to the first larval instar) should be calculated over the first month of reproduction, since it declines strongly later (Awad et al. 2013). Hatching rate in *H. axyridis* in different countries varied from 15% in Belgium (Berkvens et al. 2008) to 100% in Italy (unpublished data), although it has not been measured in most countries (Table 2). Hatching success may depend on many factors for

example food source, colour morph, endosymbionts and multiple fertilisation. Thus permanent presence of one or more males with each female is recommended.

Colour forms

More than 200 colour patterns of *H. axyridis* elytra have been observed in the native range (China), but most of them can be categorized into four major colour forms (Du, Chen 2010; Tang et al. 2012). Controlled crossing experiments have shown that the majority of color patterns are controlled by 15 alleles at one locus with multiple alleles (Komai 1956). However, all but four of these alleles are rare in natural populations, with a combined frequency of less than one per cent. The four major alleles are *succinea*, *axyridis*, *conspicua*, *spectabilis*, all but the first of these being melanic forms (Michie et al. 2010): 1) yellow to red background colour of elytra with the number of black dots ranging from 0 to 19 (f. *succinea*); 2) black background colour of elytra with many orange-red dots (of varying size and position) (f. *axyridis*); 3) black background of elytra with one large orange-red dot in the top-centre of each elytron (f. *conspicua*); 4) black background colour of elytra with two orange-red dots on each elytron and the top one larger than the bottom one (f. *spectabilis*). The impressive diversity of colour patterns documented in China has been attributed to the diverse environmental conditions in the region as well as to complex interaction between rare allelic forms (Tan 1946). A hierarchical dominance with respect to phenotype expression was demonstrated between the four major colour alleles: *conspicua* > *spectabilis* > *axyridis* > *succinea* (Michie et al. 2010; Tan, Li 1934; Tan 1946). More specifically, the expression of the colour pattern in the heterozygous individuals (i.e. individuals bearing two different allelic forms) conforms to the rule of mosaic dominance heredity, which states that any portion of the elytra which has black pigment in the homozygote for a given allele will have black pigment also in the heterozygotes in which that allele is present (Tan 1946). Interestingly, Michie, Mallard et al. (2010) reported that the non-melanic morph *succinea* dramatically increases its degree of melanization at cold temperatures, and that there is genetic variation in reaction norms, with different families responding to temperature in different ways. The exact genomic architecture and gene content of the multiallelic locus controlling color morph variation and phenotypic plasticity of the non-melanic morph remains unknown.

In some of the earliest research in population genetics, the frequencies of these alleles were found to vary hugely across the native range of *H. axyridis* (Dobzhansky 1933). The geographical variation appeared to be linked to climate, with the non-melanic form *succinea* being found most often in hot, arid regions and melanic forms being more frequent in cooler, more humid ones (Dobzhansky 1933). This might be considered as a consequence of climatic adaptation. However, a different pattern was

found in Japan, where *succinea* decreases in frequency from northeast to south-west without any significant correlation to temperature or other climatic factors (Komai 1956). Moreover, in north China *f. succinea* dominates, whereas in south China the melanic colour forms dominate (Du, Chen 2010; Tang et al. 2012). In addition to this geographic variation, small to large seasonal changes in allele frequencies have been reported in some native populations (Jiang et al. 2007; Osawa, Nishida 1992; Tang et al. 2012)). For example, in Beijing over half the population is melanic in the spring, but this drops to less than one-fifth by the autumn. Therefore, the melanic individuals presumably have a large fitness advantage in the winter and a disadvantage in the summer. This has been attributed to the effects of thermal melanism, with melanism being costly in summer and beneficial in winter, possibly mediated by mate choice (Wang et al. 2009).

The range of colour forms is limited within the invaded range. Indeed in most invaded regions only the non-melanic morph *f. succinea* is present, except in Europe where the melanic morphs *f. conspicua* and *spectabilis* are also present (and rarely *f. axyridis*), probably due to the genetic admixture with European biological control strains which occurred in this region (Lombaert et al. 2014; Lombaert et al. 2010). So far, it is not possible to say whether the predominance of *f. succinea* observed in the invaded range is due to historical and/or demographical contingencies (i.e. simple random effects) or to any selective process that would have favoured the establishment and the spatial spreading of *f. succinea*, for example the phenotypic plasticity of this morph (Michie et al. 2010). Geographical and seasonal variation of melanic and non-melanic morphs remains rather sparsely studied in Europe (Purse et al. 2014). In Belgium, the initial point of introduction of the species, the most commonly encountered morph is *f. succinea*, and the proportion of melanics (*f. spectabilis*, *f. conspicua*) is about 25% (Adriaens et al. 2008). In this particular region of Europe, strong cold hardiness and differences in several fitness parameters between melanics and non-melanics from field populations have been demonstrated (Berkvens et al. 2008; Berkvens et al. 2010a).

Natural enemies

Many regional reports have been published about one or several natural enemies of *H. axyridis* both within the native and invaded range but much could be revealed from a systematic approach comparing the native and invaded range (Table 1). There are a number of theories relating to IAS and their interactions with natural enemies but perhaps the most widely known is the Enemy Release Hypothesis (ERH) (Jeffries, Lawton 1984; Roy et al. 2011b; Roy, Lawson Handley 2012). The ERH predicts that an alien species introduced to a new region will increase in distribution and

abundance because of reduced impacts from natural enemies. Adopting a 'biogeographical' (compare richness and impacts of enemies in native and introduced populations of an alien host) or 'community' (compare native and alien species occurring within the same community) approach for *H. axyridis* would be informative (Colautti et al. 2004). With both 'biogeographical' and 'community' approaches it is essential to consider how the loss of enemy diversity translates into population regulation. A small number of enemies may have large effects and so functional diversity of enemies may be a better predictor of impacts upon hosts than overall diversity. There is considerable scope for global collaboration in exploring *H. axyridis* within the context of ERH.

Generalist predators do attack ladybirds, for example sparrows have been confirmed as predators of *H. axyridis* in Slovakia (Veselý et al. in press), but parasites are the dominant natural enemies of ladybirds. Ladybirds are attacked by over 100 species of hymenopteran and dipteran parasitoids and several fungi (Ceryngier et al. 2012; Herz, Kleespiel 2012; Riddick et al. 2009). Of these, the majority of research globally has focused on the parasitoid *Dinocampus coccinellae* (Schrank) (Hymenoptera: Braconidae), which attacks both native and alien ladybirds (Ceryngier et al. 2012). *Dinocampus coccinellae* is a parasitoid of ladybirds that has a global distribution (Ceryngier et al. 2012) but currently the influence of alien ladybirds on the abundance and distribution of this parasitoid is unknown. Ladybirds vary in their quality as a host (Comont et al. 2014b; Koyama, Majerus 2008), but *D. coccinellae* does not discriminate between suitable and unsuitable ladybirds and will attack individuals differentially based on colour and movement (Cartwright et al. 1982; Obrycki et al. 1998; Richardson, Deloach 1972). Thus, the presence of unsuitable alien ladybird hosts such as *H. axyridis* may function as an ecological trap for the parasitoid, with positive effects on native ladybirds, or alternatively increase the abundance of the parasitoid, with adverse consequences for native ladybirds. It is apparent that *H. axyridis* is less susceptible to *D. coccinellae* than are other ladybirds (Berkvens et al. 2010b; Comont et al. 2014b) but seems to be the most abundant and frequent *H. axyridis* parasitoid in many countries (Table 1) (Francati 2015a, b). Ongoing work is exploring the differences in immune response between *H. axyridis* and other ladybirds (Murray et al. 2015).

A semi-field study from the UK confirmed low rates of parasitism of *H. axyridis* by parasitoids, particularly in comparison to the native *C. septempunctata* (Comont et al. 2014b). Pupae of *H. axyridis* were parasitized, primarily by *P. fasciata* and *Phalacrotophora berolinensis* Schmitz (Diptera: Phoridae), at an exceptionally low level (1.73%) and adults were not found to be parasitized at all in this study; parasitism of the co-occurring *C. septempunctata* was high (20.91% pupae, 5.67% adults). Surveys in the Netherlands from 2003 revealed a number of natural enemies (*H. virescens*,

Parasitylenchus bifurcatus Poinar & Steenberg (Nematoda: Allantonematidae), *Coccipolipus hippodamiae* and *Dinocampus coccinellae*) attacking *H. axyridis* but only from 2008 onwards (Raak-van den Berg, van Wielink et al. 2014) (Table 1). The suggestion is made that after a time lag of six years these natural enemies are starting to use *H. axyridis* as a novel host, following host shift from native ladybirds and adaptation to *H. axyridis*.

Recent research has demonstrated that *H. axyridis* individuals contain high numbers of obligate parasitic microsporidia (Vilcinskas et al. 2013). It appears that these microsporidia do not adversely affect *H. axyridis* but cause high mortality when artificially injected into *C. septempunctata* (Vilcinskas et al. 2013). Artificial injection, as used within this study to transmit the microsporidia from *H. axyridis* to *C. septempunctata*, is far removed from the natural mechanisms involved in microsporidia transmission (Solter et al. 2013), so further studies are required to explore the ecological relevance. Indeed further research is needed to understand how the presence of all established alien ladybirds influence parasitism and pathogen infection of declining native ladybirds.

The fungal pathogen *B. bassiana* commonly infects native species of ladybird (such as *C. septempunctata*), but again, *H. axyridis* seems highly resistant (Roy et al. 2008c). However, in Denmark several entomopathogenic fungi were isolated from overwintering and early season *H. axyridis* (larvae, pupae, adults) including *B. bassiana*, *Isaria farinosa* (Holmsk.) Fr. (Ascomycota: Cordycipitaceae), *Lecanicillium lecanii* (Zimm.) Zare & W. Gams (Ascomycota: Cordycipitaceae), and *L. muscarium* (Petch) Zare & W. Gams (Steenberg, Harding 2009a) (Table 1). Similarly, mid-summer surveys in Denmark of all *H. axyridis* life stages revealed infections from the same fungal assemblage with *I. farinosa* most prevalent, followed by *Lecanicillium* spp. (Howe, Ravn, Jensen, Meyling, unpublished data).

Additional work has assessed the sexually-transmitted ectoparasitic mite *Coccipolipus hippodamiae* (McDaniel & Morrill) (Acari: Podapolipidae) as a biological control candidate against *H. axyridis* (Roy et al. 2011c). This mite, which occurs naturally in Europe and North America (Table 1), causes sterility in female *H. axyridis*, but some native ladybird species are also susceptible to the mite, and thus it does not represent a suitable control strategy for *H. axyridis* (Rhule et al. 2010).

The obligate ectoparasitic fungus *H. virescens* has been capturing the imagination of a number of scientists working on *H. axyridis*. This species has historically received very little attention and so is of particular note. While most Laboulbeniales exhibit a high degree of host specificity, *H. virescens*

has been reported from as many as 30 ladybird species in 17 genera (Bernardi et al. 2014; Ceryngier et al. 2012; Ceryngier, Twardowska 2013; Haelewaters et al. 2012; Haelewaters et al. 2015b). Transmission of *H. virescens* occurs mainly during sexual contact, as exemplified by the non-random distribution of thalli on the body of males and females (Riddick et al. 2009; Welch et al. 2001); infection can be considered as a sexually transmitted disease (Welch et al. 2001). In *H. axyridis*, however, *H. virescens* is also socially transmitted; in overwintering aggregations, transmission of *H. virescens* through direct physical contact is the most important mechanism of spread (Nalepa, Weir 2007; Riddick 2006). Infection is caused by grooming, resulting in high thallus densities on older hosts (Haelewaters et al. 2012). *Hesperomyces virescens* was reported for the first time on *H. axyridis* in Ohio in summer 2002 (Garcés, Williams 2004). *Harmonia axyridis* is multivoltine, promiscuous, and overwinters in aggregations, all of which contribute to the rapid spread of *H. virescens* and higher infection prevalence on this host, compared to other ladybird hosts (De Kesel 2011). Interestingly, the parasite prevalence of *H. virescens* on *H. axyridis* varies between locations and between years (Haelewaters et al. 2012; Raak-van den Berg et al. 2014). In Belgium, for example, an increase from 0.5% to 96.5% of parasite prevalence was noted after only four years (De Kesel 2011). Currently, *H. virescens* infection of *H. axyridis* is widespread in Western Europe, the eastern United States and to a lesser extent in South Africa (D. Haelewaters unpublished). *Hesperomyces virescens* has also been reported on *H. axyridis* in its native range with one record from China, (Haelewaters et al. 2014). Ongoing work is assessing the influence of (dual) fungal infections on *H. axyridis* and *Olla v-nigrum* (Mulsant), a North American native ladybird species (Haelewaters et al. 2015a).

There is clearly much to uncover about the interactions between *H. axyridis* and natural enemies; global collaborations will provide unique opportunities for exploring these on biogeographic scales.

Conclusions and future directions

Harmonia axyridis has inspired global collaborations and has also been the impetus for understanding biological invasions within and between countries. Many countries have documented the distribution and noted the rapid spread of *H. axyridis* following establishment. Further research is required to improve our understanding of the factors involved in determining the global patterns of invasion by *H. axyridis* which will have wide relevance for invasion biology. Many countries have engaged members of the public in monitoring the distribution of *H. axyridis* and the lessons learnt from such initiatives have been shared and proved informative for developing approaches to citizen science (Gardiner et al. 2012; Pocock et al. 2015; Roy et al. 2012a) and inspiring new projects both

within and between countries. Indeed the role of volunteers in monitoring IAS is recognised (Roy et al. 2015b) and there is considerable scope to share resources and technology (August et al. 2015) to increase involvement in monitoring ladybirds and other IAS around the world.

The interactions between *H. axyridis* and other species have fascinated ecologists both in the native and invaded ranges of this species. There has been considerable focus on the potential impacts of *H. axyridis* on biodiversity, particularly intra-guild interactions and specifically IGP. Competitive interactions have received less attention but are worthy of future research emphasis. There is still much to unravel about the interactions between *H. axyridis* and its natural enemies. Comparisons between the native and invaded range will be particularly fruitful. Molecular studies have provided insights into genetic aspects of invasion and there is certainly more that can be revealed from detailed studies at a global scale. It is exciting to consider the potential of ongoing research to sequence the genome of *H. axyridis* and consider the opportunities that this research might present for future studies (Chown et al. 2015) (<http://www.agence-nationale-recherche.fr/?Project=ANR-13-EBID-0001>).

Many of the studies examining the interactions between *H. axyridis* and other species have involved mesocosm experiments under laboratory conditions. There is a need to increase the scale of such studies to consider the negative and positive ecosystem-level impacts of *H. axyridis*. *Harmonia axyridis* contributes pest control services in a number of crop systems but there is a lack of evidence in relation to ecosystem function and resilience of invaded systems. New molecular methods provide opportunities for detailed studies on the interactions between *H. axyridis* and the diverse range of species with which it interacts. Ecological network analysis represents an appealing and exciting way to explore these complex communities (Roy, Lawson Handley 2012). The coupling of citizen science approaches with global collaborations between researchers will provide the scale of information required to address some of the complex ecological questions that remain unanswered.

Acknowledgements

The paper had its origin at a workshop on “Drivers, impacts, mechanisms and adaptation in insect invasions” hosted by the DST-NRF Centre of Excellence for Invasion Biology in Stellenbosch, South Africa, in November 2014. We thank all our collaborators, and particularly the volunteer community, who have contributed to research around the world on *H. axyridis*. The number of references included reflects the range of inspiring studies on *H. axyridis* from so many people—we look forward to new and continued collaborations in the future. We are grateful to the editors of this special issue

for inviting this review and providing an opportunity to explore ideas through the “Invasive Insects Workshop funding (NRF South Africa, CIB)”. The UK Ladybird Survey and associated co-authors are supported by the Biological Records Centre (part of the Centre for Ecology & Hydrology), which receives support from both the Natural Environment Research Council and the Joint Nature Conservation Committee. The IOBC WPRS and Global Working Groups “Benefits and Risks of Exotic Biological Control Agents” and the COST Action TD1209 “Alien Challenge” have facilitated discussions and collaborations on *H. axyridis*. This study was supported by the French Agropolis Fondation (Labex Agro-Montpellier, BIOFIS Project Number 1001-001) and by a grant from the ERA-Net BiodivERsA, with the national funders ANR (France), DFG (Germany) and BELSPO (Belgium), as part of the 2012–2013 BiodivERsA call for research proposals. Support has been also received from FONDECYT 1140662 (Chile). The study of M.J. Orlova-Bienkowskaja and I. A. Zakharov was supported by Russian Science Foundation, project № 16-16- 00079. Gabriele Rondoni acknowledges financial support from Fondazione Cassa di Risparmio di Perugia. Riaan Stals acknowledges funding from the Department of Science and Technology, South Africa. The research of Peter Zach and colleagues was funded by the project VEGA 2/0035/13 and VEGA 2/0052/15. A. Honěk and Z. Martinkova were supported by grants GACR 14-26561S and COST CZ LD14084. Research in Switzerland is funded by the Swiss Federal Office for the Environment.

References

- Adriaens T, Branquart E, Maes D (2003) The multicoloured Asian ladybird *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae), a threat for native aphid predators in Belgium? Belgian Journal of Zoology 133:195-196
- Adriaens T, Gomez GMY, Maes D (2008) Invasion history, habitat preferences and phenology of the invasive ladybird *Harmonia axyridis* in Belgium. Biocontrol 53:69-88
- Adriaens T, San Martin y Gomez G, Bogaert J, et al. (2015) Testing the applicability of regional IUCN Red List criteria on ladybirds (Coleoptera, Coccinellidae) in Flanders (north Belgium): opportunities for conservation. Insect Conservation and Diversity 8:404–417
- Aebi A, Brown PMJ, De Clercq P, et al. (2011) Detecting arthropod intraguild predation in the field. BioControl 56:429-440
- Alyokhin A, Sewell G (2004) Changes in a lady beetle community following the establishment of three alien species. Biological Invasions 6:463-471
- Amat-García G, Amat-García E, Ariza-Marín E (2011) Insectos invasores en los tiempos de cambio climático. Innovación y Ciencia 18:45-53
- August T, Harvey M, Lightfoot P, et al. (2015) Emerging technologies for biological recording. Biological Journal of the Linnean Society 115:731-749
- Awad M, Kalushkov P, Nedvědová T, et al. (2013) Fecundity and fertility of ladybird beetle *Harmonia axyridis* after prolonged cold storage. BioControl 58:657-666
- Bahlai C, Sears M (2009) Population dynamics of *Harmonia axyridis* and *Aphis glycines* in Niagara Peninsula soybean fields and vineyards. J Entomol Soc Ont 140:27-39
- Bahlai CA, Colunga-Garcia M, Gage SH, et al. (2014) The role of exotic ladybeetles in the decline of native ladybeetle populations: evidence from long-term monitoring. Biological Invasions:1-20
- Bahlai CA, vander Werf W, O'Neal M, et al. (2015) Shifts in dynamic regime of an invasive lady beetle are linked to the invasion and insecticidal management of its prey. Ecological Applications
- Bazzocchi GG, Lanzoni A, Accinelli G, et al. (2004) Overwintering, phenology and fecundity of *Harmonia axyridis* in comparison with native coccinellid species in Italy. BioControl 49:245-260
- Belyakova NA, Reznik SY (2013) First record of the harlequin ladybird, *Harmonia axyridis* (Coleoptera: Coccinellidae) in the Caucasus. European Journal of Entomology 110:699-702
- Benelli M, Simon R, Francati S, et al. (2015) Effect of two temperatures on biological traits and susceptibility to a pyrethroid insecticide in an exotic and native coccinellid species. Bulletin of Insectology 68:23-29
- Berkvens N, Bonte J, Berkvens D, et al. (2008) Influence of diet and photoperiod on development and reproduction of European populations of *Harmonia axyridis* (Pallas) (Coleoptera : Coccinellidae). BioControl 53:211-221
- Berkvens N, Landuyt C, Deforce K, et al. (2010a) Alternative foods for the multicoloured Asian lady beetle *Harmonia axyridis* (Coleoptera: Coccinellidae). Eur. J. Entomol 107:189-195
- Berkvens N, Moens J, Berkvens D, et al. (2010b) *Dinocampus coccinellae* as a parasitoid of the invasive ladybird *Harmonia axyridis* in Europe. Biological Control 53:92-99
- Bernardi M, Barragán A, Rossi W (2014) New records of Laboulbeniales (Fungi: Ascomycota) from Ecuador and other countries. Webbia 69:281-289
- Blekhman A (2008) Population variation of elytral ridge occurrence in ladybirds *Harmonia axyridis* Pallas. Russ. J. Genet. 44:1351-1354
- Borges I, Soares A, Hemptinne JL (2006) Abundance and spatial distribution of aphids and scales select for different life histories in their ladybird beetle predators. Journal of Applied Entomology 130:461-464
- Botezatu AI, Kotseridis Y, Inglis D, et al. (2013) Occurrence and contribution of alkyl methoxypyrazines in wine tainted by *Harmonia axyridis* and *Coccinella septempunctata*. Journal of the Science of Food and Agriculture 93:803-810

1473 Brakefield PM (1985) Differential winter mortality and seasonal selection in the polymorphic
 1474 ladybird *Adalia bipunctata* (L) in the Netherlands. Biological Journal of the Linnean Society 24:189-
 1475 206

1476 Brown MW, Miller SS (1998) Coccinellidae (Coleoptera) in apple orchards of eastern West Virginia
 1477 and the impact of invasion by *Harmonia axyridis*. Entomological News 109:143-151

1478 Brown PMJ, Adriaens T, Bathon H, et al. (2008a) *Harmonia axyridis* in Europe: spread and
 1479 distribution of a non-native coccinellid. BioControl 53:5-21

1480 Brown PMJ, Frost R, Doberski J, et al. (2011a) Decline in native ladybirds in response to the arrival of
 1481 *Harmonia axyridis*: early evidence from England. Ecological Entomology 36:231-240

1482 Brown PMJ, Ingels B, Wheatley A, et al. (2015) Intraguild predation by *Harmonia axyridis*
 1483 (Coleoptera: Coccinellidae) on native insects in Europe: molecular detection from field samples.
 1484 Entomological Science 18:130-133

1485 Brown PMJ, Roy HE (In press) Reflections on the long-term assessment of ladybird (Coleoptera:
 1486 Coccinellidae) populations in the Czech Republic and the United Kingdom. Soc. Zool. Bohem. 79

1487 Brown PMJ, Roy HE, Rothery P, et al. (2008b) *Harmonia axyridis* in Great Britain: analysis of the
 1488 spread and distribution of a non-native coccinellid. BioControl 53:55-67

1489 Brown PMJ, Thomas CE, Lombaert E, et al. (2011b) The global spread of *Harmonia axyridis*
 1490 (Coleoptera: Coccinellidae): distribution, dispersal and routes of invasion. BioControl 56:623-641

1491 Bukejs A, Telnov D (2015) The first record of the invasive lady beetle *Harmonia axyridis* (Pallas,
 1492 1773)(Coleoptera: Coccinellidae) in Turkey. Zoology and Ecology 25:59-62

1493 Burgio G, Lanzoni A, Accinelli G, et al. (2008a) Estimation of mortality by entomophages on exotic
 1494 *Harmonia axyridis* versus native *Adalia bipunctata* in semi-field conditions in northern Italy.
 1495 BioControl 53:277-287

1496 Burgio G, Santi F, Lanzoni A, et al. (2008b) *Harmonia axyridis* recordings in northern Italy. Bulletin of
 1497 Insectology 61:361-364

1498 Cabral S, Soares AO, Moura R, et al. (2006) Suitability of *Aphis fabae*, *Myzus persicae* (Homoptera :
 1499 Aphididae) and *Aleyrodes proletella* (Homoptera : Aleyrodidae) as prey for *Coccinella*
 1500 *undecimpunctata* (Coleoptera : Coccinellidae). Biological Control 39:434-440

1501 Caltagirone L, Doult R (1989) The history of the vedalia beetle importation to California and its
 1502 impact on the development of biological control. Annual review of entomology 34:1-16

1503 Cartwright B, Eikenbary R, Angalet G (1982) Parasitism by *Perilitus coccinellae* [Hym.: Braconidae] of
 1504 indigenous coccinellid hosts and the introduced *Coccinella septempunctata* [Col.: Coccinellidae], with
 1505 notes on winter mortality. Entomophaga 27:237-243

1506 Ceryngier P, Roy HE, Poland RL (2012) Natural enemies of ladybird beetles. In: Hodek I, van Emden
 1507 HF and Honek A (eds) Ecology and behaviour of the ladybird beetles (Coccinellidae). John Wiley &
 1508 Sons,

1509 Ceryngier P, Twardowska K (2013) *Harmonia axyridis* (Coleoptera: Coccinellidae) as a host of the
 1510 parasitic fungus *Hesperomyces virescens* (Ascomycota: Laboulbeniales, Laboulbeniaceae): A case
 1511 report and short review. Eur. J. Entomol 110:549-557

1512 Chacón J, Landis D, Heimpel G (2008) Potential for biotic interference of a classical biological control
 1513 agent of the soybean aphid. Biological control 46:216-225

1514 Chapin JB, Brou VA (1991) *Harmonia axyridis* (Pallas), the 3rd species of the genus to be found in the
 1515 United States (Coleoptera, Coccinellidae). Proceedings of the Entomological Society of Washington
 1516 93:630-635

1517 Chown SL, Hodgins KA, Griffin PC, et al. (2015) Biological invasions, climate change and genomics.
 1518 Evolutionary Applications 8:23-46

1519 Coderre D, Lucas É, Gagné I (1995) The occurrence of *Harmonia axyridis* (Pallas)(Coleoptera:
 1520 Coccinellidae) in Canada. The Canadian Entomologist 127:609-611

1521 Colautti RI, Ricciardi A, Grigorovich IA, et al. (2004) Is invasion success explained by the enemy
 1522 release hypothesis? Ecology Letters 7:721-733

1523 Colunga-Garcia M, Gage SH (1998) Arrival, establishment, and habitat use of the multicolored Asian
1524 lady beetle (Coleoptera: Coccinellidae) in a Michigan landscape. *Environmental Entomology* 27:1574-
1525 1580

1526 Comont R, Roy H, Harrington R, et al. (2014a) Ecological correlates of local extinction and
1527 colonisation in the British ladybird beetles (Coleoptera: Coccinellidae). *Biological Invasions* 16:1805-
1528 1817

1529 Comont RF, Purse BV, Phillips W, et al. (2014b) Escape from parasitism by the invasive alien ladybird,
1530 *Harmonia axyridis*. *Insect Conservation and Diversity* 7:334-342

1531 Comont RF, Roy HE, Lewis OT, et al. (2012) Using biological traits to explain ladybird distribution
1532 patterns. *Journal of Biogeography* 39:1772-1781

1533 Cornacchia P, Nardi G (2012) Nuovi dati su *Harmonia axyridis* in Italia (Coleoptera, Coccinellidae).
1534 *Bollettino dell'Associazione Romana di Entomologia* 67:51-68

1535 Costamagna AC, Landis DA (2007) Quantifying predation on soybean aphid through direct field
1536 observations. *Biological Control* 42:16-24

1537 Costamagna AC, Landis DA, Brewer MJ (2008) The role of natural enemy guilds in *Aphis glycines*
1538 suppression. *Biological Control* 45:368-379

1539 Cottrell TE (2004) Suitability of exotic and native lady beetle eggs (Coleoptera : Coccinellidae) for
1540 development of lady beetle larvae. *Biological Control* 31:362-371

1541 Coutanceau J-P (2006a) *Harmonia axyridis* (Pallas, 1773): une coccinelle asiatique introduite,
1542 acclimatée et en extension en France. *Bulletin de la Société entomologique de France* 111:395-401

1543 Coutanceau JP (2006b) *Harmonia axyridis* (Pallas, 1773): une coccinelle asiatique introduite,
1544 acclimatée et en extension en France. *Bulletin de la Societe Entomologique de France* 111:395-401

1545 Cuppen J, Heijerman T, van Wielink P, et al. (2004) Het lieveheersbeestje *Harmonia axyridis* in
1546 Nederland: Een aanwinst voor onze fauna of een ongewenste indringer (Coleoptera: Coccinellidae)?
1547 *Nederlandse Faunistische Mededelingen* 20:1-12

1548 Davidson LN, Evans EW (2010) Frass analysis of diets of aphidophagous lady beetles (Coleoptera:
1549 Coccinellidae) in Utah alfalfa fields. *Environmental entomology* 39:576-582

1550 De Clercq P, Bale JS (2011) Risks of invertebrate biological control agents—*Harmonia axyridis* as a
1551 case study. *Regulation of Biological Control Agents*. Springer, pp. 243-255

1552 De Kesel A (2011) *Hesperomyces* (Laboulbeniales) and coccinellid hosts. *Sterbeeckia* 30:32-37

1553 Department of Environmental Affairs South Africa (2014a) National Environmental Management:
1554 Biodiversity Act 2004 (Act No. 10 of 2004). Alien and Invasive Species Lists, 2014. Government
1555 Gazette 37886:3–74.

1556 Department of Environmental Affairs South Africa (2014b) National Environmental Management:
1557 Biodiversity Act 2004 (Act No. 10 of 2004). Alien and Invasive Species Regulations, 2014.
1558 Government Gazette 37885:3–27.

1559 Dobzhansky T (1924) Die geographische und individuelle Variabilität von *Harmonia axyridis* Pall. in
1560 ihren Wechselbeziehungen. *Biologisches Zentralblatt* 44:401-421

1561 Dobzhansky T (1933) Geographical Variation in Ladybeetles. *The American Naturalist* 67:97-126

1562 Dreistadt S, Hagen K, Bezark L (1995) *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae), first
1563 western United States record for this Asiatic lady beetle. *Pan-Pacific Entomologist* 71:135-136

1564 Du W, Chen I (2010) Investigations of the Color Variations of *Harmonia axyridis* (Pallas) in Changchun
1565 [J]. *Chinese Agricultural Science Bulletin* 4:057

1566 Dudek K, Dudek M, Tryjanowski P (2015) Wind turbines as overwintering sites attractive to an
1567 invasive lady beetle, *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae) *The Coleopterists Bulletin*
1568 69:1-5

1569 Durieux D, Fischer C, Brostaux Y, et al. (2012) Role of long-chain hydrocarbons in the aggregation
1570 behaviour of *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). *Journal of Insect Physiology*
1571 58:801-807

1572 El-Arnaouty S, Beyssat-Arnaouty V, Ferran A, et al. (2000) Introduction and release of the coccinellid
 1573 *Harmonia axyridis* Pallas for controlling *Aphis craccivora* Koch on faba beans in Egypt. Egyptian
 1574 Journal of biological pest control 10:129-136
 1575 Eschen R, Babendreier D, Nauer S, et al. (2007) Surveys for ladybirds (Coleoptera : Coccinellidae) in
 1576 Switzerland and confirmation of the presence of the invasive alien ladybird species, *Harmonia*
 1577 *axyridis* (Pallas). Mitteilungen der Schweizerischen Entomologischen Gesellschaft 80:7-14
 1578 Evans EW (2004) Habitat displacement of North American ladybirds by an introduced species.
 1579 Ecology 85:637-647
 1580 Evans EW, Soares AO, Yasuda H (2011) Invasions by ladybugs, ladybirds, and other predatory
 1581 beetles. BioControl 56:597-611
 1582 Facon B, Hufbauer RA, Tayeh A, et al. (2011) Inbreeding depression is purged in the invasive insect
 1583 *Harmonia axyridis*. Current Biology 21:424-427
 1584 Felix S, Soares AO (2004) Intraguild predation between the aphidophagous ladybird beetles
 1585 *Harmonia axyridis* and *Coccinella undecimpunctata* (Coleoptera : Coccinellidae): the role of body
 1586 weight. European Journal of Entomology 101:237-242
 1587 Ferran A, Giuge L, Brun J, et al. (1997) Coccinelle *Harmonia axyridis* pallas : mise au point sur son
 1588 introduction et son utilisation en lutte biologique. Adalia 36:21-24
 1589 Foley IA, Ivie MA, Denke PM (2009) The first state record for the multicolored Asian lady beetle,
 1590 *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae), from Montana. The Coleopterists Bulletin
 1591 63:351-352
 1592 Fothergill K, Moore W, Losey J, et al. (2010) First Arizona records of the multicolored Asian lady
 1593 beetle, *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae). The Coleopterists Bulletin 64:51-52
 1594 Francati S (2015a) Native preimaginal parasitism of *Harmonia axyridis*: New record of association
 1595 with *Phalacotrophora fasciata* in Italy. Bull. Insectol. 68:3-6
 1596 Francati S (2015b) Native preimaginal parasitism of *Harmonia axyridis*: New record of association
 1597 with *Phalacotrophora fasciata* in Italy. Bull. Insectol. 68:3-6
 1598 Gagnon AÈ, Doyon J, Heimpel G, et al. (2011) Prey DNA detection success following digestion by
 1599 intraguild predators: influence of prey and predator species. Mol. Ecol. Resour. 11:1022-1032
 1600 Galvan TL, Burkness EC, Hutchison WD (2006) Influence of berry injury on infestations of the
 1601 multicolored Asian lady beetle in wine grapes. Plant Health Prog. doi 10:1094
 1602 Galvan TL, Koch RL, Hutchison WD (2008) Impact of fruit feeding on overwintering survival of the
 1603 multicolored Asian lady beetle, and the ability of this insect and paper wasps to injure wine grape
 1604 berries. Entomologia experimentalis et applicata 128:429-436
 1605 Garcés S, Williams R (2004) First record of *Hesperomyces virescens* Thaxter (Laboulbeniales:
 1606 Ascomycetes) on *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae). J. Kans. Entomol. Soc.
 1607 77:156-158
 1608 Garcia V (1986) Approaches to integrated control of some citrus pests in the Azores and Algarve
 1609 (Portugal). Integrated pest control in citrus-groves: proceedings of the experts' meeting, Acireale
 1610 (Italy), 26-29 Mar 1985. AA Balkema,
 1611 Gardiner M, Landis D (2007) Impact of intraguild predation by adult *Harmonia axyridis* (Coleoptera:
 1612 Coccinellidae) on *Aphis glycines* (Hemiptera: Aphididae) biological control in cage studies. Biological
 1613 control 40:386-395
 1614 Gardiner M, Landis D, Gratton C, et al. (2009a) Landscape diversity enhances biological control of an
 1615 introduced crop pest in the north-central USA. Ecological Applications 19:143-154
 1616 Gardiner M, Landis D, Gratton C, et al. (2009b) Landscape composition influences patterns of native
 1617 and exotic lady beetle abundance. Diversity and Distributions 15:554-564
 1618 Gardiner MM, Allee LL, Brown PM, et al. (2012) Lessons from lady beetles: accuracy of monitoring
 1619 data from US and UK citizen-science programs. Frontiers in Ecology and the Environment 10:471-476
 1620 Gardiner MM, Landis DA, Gratton C, et al. (2009c) Landscape composition influences patterns of
 1621 native and exotic lady beetle abundance. Diversity and Distributions 15:554-564

1622 Glemser EJ, Dowling L, Inglis D, et al. (2012) A novel method for controlling multicolored Asian lady
 1623 beetle (Coleoptera: Coccinellidae) in vineyards. *Environmental entomology* 41:1169-1176
 1624 Goetz DW (2008) *Harmonia axyridis* ladybug invasion and allergy. *Allergy and Asthma Proceedings*
 1625 29:123-129
 1626 Goldarazena A, Calvo D (2007) First record of *Harmonia axyridis* (Coleoptera: Coccinellidae) from the
 1627 Iberian Peninsula. *Boletín Sociedad Entomol Aragonesa* 41:437-439
 1628 González G (2008) Lista y distribución geográfica de especies de Coccinellidae (Insecta: Coleoptera)
 1629 presentes en Chile. *Boletín del Museo Nacional de Historia Natural (Chile)* 57:77-107
 1630 González GF, Kondo T (2012) Primer registro de la especie invasora *Harmonia axyridis*
 1631 (Pallas)(Coleoptera: Coccinellidae) en Ecuador. *Boletín de la Sociedad Entomológica Aragonesa*
 1632 51:310
 1633 Gordon RD (1985) The Coleoptera (Coccinellidae) of America north of Mexico. *Journal of the New*
 1634 *York Entomological Society* 93:1-912
 1635 Grez A, Zaviezo T, González G, et al. (2010) *Harmonia axyridis* en Chile: una nueva amenaza. *Ciencia*
 1636 *e investigación agraria* 37:145-149
 1637 Grez AA, Rand TA, Zaviezo T, et al. (2013) Land use intensification differentially benefits alien over
 1638 native predators in agricultural landscape mosaics. *Diversity and Distributions* 19:749-759
 1639 Grez AA, Zaviezo T, Gardiner MM (2014a) Local predator composition and landscape affects
 1640 biological control of aphids in alfalfa fields. *Biological Control* 76:1-9
 1641 Grez AA, Zaviezo T, Hernández J, et al. (2014b) The heterogeneity and composition of agricultural
 1642 landscapes influence native and exotic coccinellids in alfalfa fields. *Agricultural and Forest*
 1643 *Entomology* 16:382-390
 1644 Guo JY, Wan FH (2001) Effect of three diets on development and fecundity of the ladybeetles
 1645 *Harmonia axyridis* and *Propylaea japonica* *Chinese Journal of Biological Control* 17:116-120
 1646 Haelewaters D, Comont RF, Zhao SY, et al. (2014) *Hesperomyces virescens* (Fungi, Ascomycota,
 1647 Laboulbeniales) attacking *Harmonia axyridis* (Coleoptera, Coccinellidae) in its native range. *Chinese*
 1648 *Science Bulletin* 59:528-532
 1649 Haelewaters D, Shapiro-Ilan DI, Cottrell TE (2015a) Will dual fungal infections increase *Harmonia*
 1650 *axyridis* mortality in natural populations? 3rd Meeting of IOBC-WPRS study group "Benefits and risks
 1651 of exotic biological control agents" 13 to 15 May 2015, Bornholm, Denmark.
 1652 Haelewaters D, van Wielink P, van Zuijlen JW, et al. (2012) New records of Laboulbeniales (Fungi,
 1653 Ascomycota) for the Netherlands. *Entomol. Ber* 72:26-33
 1654 Haelewaters D, Zhao SY, Royer IR, et al. (2015b) Laboulbeniales (Ascomycota) of the Boston Harbor
 1655 Islands I: species parasitizing Coccinellidae and Staphylinidae. *Northeastern Naturalist* 22:459-477
 1656 Harmon JP, Stephens E, Losey J (2007) The decline of native coccinellids (Coleoptera : Coccinellidae)
 1657 in the United States and Canada. *J Insect Conserv* 11:85-94
 1658 Hautier L, Gregoire JC, de Schauwers J, et al. (2008) Intraguild predation by *Harmonia axyridis* on
 1659 coccinellids revealed by exogenous alkaloid sequestration. *Chemoecology* 18:191-196
 1660 Hautier L, San Martin G, Callier P, et al. (2011) Alkaloids provide evidence of intraguild predation on
 1661 native coccinellids by *Harmonia axyridis* in the field. *Biological Invasions* 13:1805-1814
 1662 Heimpel GE, Frelich LE, Landis DA, et al. (2010) European buckthorn and Asian soybean aphid as
 1663 components of an extensive invasional meltdown in North America. *Biological Invasions* 12:2913-
 1664 2931
 1665 Hemptinne J-L, Magro A, Evans E, et al. (2012) Body size and the rate of spread of invasive ladybird
 1666 beetles in North America. *Biological Invasions* 14:595-605
 1667 Herz A, Kleespiel RG (2012) Occurrence of natural enemies in different populations of the invasive
 1668 ladybird *Harmonia axyridis* (Pallas, 1771) (Coleoptera, Coccinellidae) in Germany. *Mitt Dtsch Ges Allg*
 1669 *Angew Entomol* 18:201-206
 1670 Hesler L, Kieckhefer R, Beck D (2001) First record of *Harmonia axyridis* (Coleoptera: Coccinellidae) in
 1671 South Dakota and notes on its activity there and in Minnesota. *Entomological News* 112:264-270

1672 Hesler LS (2014) Inventory and assessment of foliar natural enemies of the soybean aphid
 1673 (Hemiptera: Aphididae) in South Dakota. *Environmental entomology* 43:577-588
 1674 Hesler LS, Kieckhefer RW (2008) Status of exotic and previously common native coccinellids
 1675 (Coleoptera) in South Dakota landscapes. *J. Kans. Entomol. Soc.* 81:29-49
 1676 Hicks B, Majka CG, Moores SP (2010) *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae) found in
 1677 Newfoundland. *The Coleopterists Bulletin* 64:50-50
 1678 Hodek I, van Emden HF, Honek A (2012) Ecology and behaviour of the ladybird beetles
 1679 (Coccinellidae). Wiley-Blackwell, Chichester, West Sussex
 1680 Hoffmann AA, Chown SL, Clusella-Trullas S (2013) Upper thermal limits in terrestrial ectotherms:
 1681 how constrained are they? *Functional Ecology* 27:934-949
 1682 Honek A, Martinkova Z, Dixon AFG, et al. (submitted) Long-term changes in communities of native
 1683 coccinellids: population fluctuations and the effect of competition from an invasive species. *Insect*
 1684 *Conservation and Diversity*
 1685 Honek A, Martinkova Z, Kindlmann P, et al. (2014) Long-term trends in the composition of
 1686 aphidophagous coccinellid communities in Central Europe. *Insect conservation and diversity* 7:55-63
 1687 Hoogendoorn A, Heimpel G (2002) Indirect interactions between an introduced and a native ladybird
 1688 beetle species mediated by a shared parasitoid. *Biological Control* 25:224-230
 1689 Hoogendoorn M, Heimpel GE (2004) Competitive interactions between an exotic and a native
 1690 ladybeetle: a field cage study. *Entomologia Experimentalis et Applicata* 111:19-28
 1691 Howe AG (2015) Harlekinmariehønen-en bornholmsk status. *Natur på Bornholm* 13
 1692 Howe AG, Ransijn J, Ravn HP (2015) A sublethal effect on native *Anthocoris nemoralis* through
 1693 competitive interactions with invasive *Harmonia axyridis*. *Ecological Entomology*
 1694 Iazlovețchii I, Sumencova V (2013) New invasive species in the Republic of Moldova: multicolored
 1695 Asian ladybird *Harmonia axyridis* Pallas (Coleoptera: Coccinellidae). VIII international conference of
 1696 zoologists. Actual problems of protection and sustainable use of the animal world diversity (Chisinau,
 1697 10–12.10. 2013). Book of abstract. Academy of Sciences of Moldova, Chisinau. pp. 136-137
 1698 Ingels B, De Clercq P (2011) Effect of size, extraguild prey and habitat complexity on intraguild
 1699 interactions: a case study with the invasive ladybird *Harmonia axyridis* and the hoverfly *Episyrphus*
 1700 *balteatus*. *BioControl* 56:871-882
 1701 Ingels B, Van Hassel P, Van Leeuwen T, et al. (2015) Feeding History Affects Intraguild Interactions
 1702 between *Harmonia axyridis* (Coleoptera: Coccinellidae) and *Episyrphus balteatus* (Diptera:
 1703 Syrphidae). *PloS one* 10:e0128518
 1704 Izhevsky S (1990) Introduction and applications of entomophags. Moscow, VO "Agropromizdat
 1705 Jacas J-A, Urbaneja A, Viñuela E (2006) History and future of introduction of exotic arthropod
 1706 biological control agents in Spain: a dilemma? *BioControl* 51:1-30
 1707 Jansen JP, Hautier L (2008) Ladybird population dynamics in potato: comparison of native species
 1708 with an invasive species, *Harmonia axyridis*. *BioControl* 53:223-233
 1709 Jeffries DL, Chapman J, Roy HE, et al. (2013) Characteristics and Drivers of High-Altitude Ladybird
 1710 Flight: Insights from Vertical-Looking Entomological Radar. *PLoS ONE* 8(12): e82278
 1711 Jeffries MJ, Lawton JH (1984) Enemy free space and the structure of ecological communities.
 1712 *Biological Journal of the Linnean Society* 23:269-286
 1713 Jiang W-H, Pan X-H, Liu J-X, et al. (2007) Studies on the type of spot in the wings of *Harmonia*
 1714 *axyridis* (Pallas) in Baoding [J]. *Hebei Journal of Forestry and Orchard Research* 2:022
 1715 Katsanis A (2011) Multitrophic Interactions Between the Invasive Coccinellid, *Harmonia axyridis*, and
 1716 Non-target Insects. Bern University,
 1717 Katsanis A, Babendreier D, Nentwig W, et al. (2013) Intraguild predation between the invasive
 1718 ladybird *Harmonia axyridis* and non-target European coccinellid species. *BioControl*:1-11
 1719 Kemp E, Cottrell T (2015) Effect of lures and colors on capture of lady beetles (coleoptera:
 1720 coccinellidae) in tedders pyramidal traps. *Environmental entomology:nvv108*
 1721 Kenis M, Adriaens T, Brown P, et al. (2010) Impact of *Harmonia axyridis* on European ladybirds:
 1722 which species are most at risk. *IOBC/WPRS Bull* 58:57-59

1723 Kenis M, Roy HE, Zindel R, et al. (2008) Current and potential management strategies against
 1724 *Harmonia axyridis*. *BioControl* 53:235-252
 1725 Klausnitzer B (2002) *Harmonia axyridis* (Pallas, 1773) in Germany (Col., Coccinellidae).
 1726 *Entomologische Nachrichten und Berichte* 46:177-183
 1727 Koch R, Burkness E, Burkness SJW, et al. (2004a) Phytophagous preferences of the multicolored
 1728 Asian lady beetle (Coleoptera: Coccinellidae) for autumn-ripening fruit. *Journal of Economic*
 1729 *Entomology* 97:539-544
 1730 Koch R, Venette R, Hutchison W (2006a) Predicted impact of an exotic generalist predator on
 1731 monarch butterfly (Lepidoptera: Nymphalidae) populations: a quantitative risk assessment.
 1732 *Biological Invasions* 8:1179-1193
 1733 Koch RL (2003) The multicolored Asian lady beetle, *Harmonia axyridis*: a review of its biology, uses in
 1734 biological control, and non-target impacts. *J Insect Sci.* 3
 1735 Koch RL, Carrillo MA, Venette RC, et al. (2004b) Cold hardiness of the Multicolored Asian lady beetle
 1736 (Coleoptera : Coccinellidae). *Environmental Entomology* 33:815-822
 1737 Koch RL, Fernandes MG, Dutra CC (2011) First confirmed record of *Harmonia axyridis* (Pallas,
 1738 1773)(Coleoptera: Coccinellidae) in the state of Mato Grosso do Sul, Brazil. *Check List* 7:476-477
 1739 Koch RL, Galvan TL (2008) Bad side of a good beetle: the North American experience with *Harmonia*
 1740 *axyridis*. *BioControl* 53:23-35
 1741 Koch RL, Hutchison WD, Venette RC, et al. (2003) Susceptibility of immature monarch butterfly,
 1742 *Danaus plexippus* (Lepidoptera : Nymphalidae : Danainae), to predation by *Harmonia axyridis*
 1743 (Coleoptera : Coccinellidae). *Biological Control* 28:265-270
 1744 Koch RL, Venette RC, Hutchison WD (2006b) Invasions by *Harmonia axyridis* (Pallas)(Coleoptera:
 1745 Coccinellidae) in the western hemisphere: implications for South America. *Neotropical Entomology*
 1746 35:421-434
 1747 Kögel S, Eben A, Hoffmann C, et al. (2012) Influence of diet on fecundity, immune defense and
 1748 content of 2-isopropyl-3-methoxypyrazine in *Harmonia axyridis* Pallas. *Journal of chemical ecology*
 1749 38:854-864
 1750 Komai T (1956) Genetics of ladybirds. *Advanced Genetics* 8:155-188
 1751 Kondo T, González GF (2013) The multicolored Asian lady beetle, *Harmonia axyridis* (Pallas,
 1752 1773)(Coleoptera: Coccinellidae), a not so new invasive insect in Colombia and South America.
 1753 *Insecta Mundi*:1-7
 1754 Kontodimas DC, Stathas GJ, Martinou AF (2008) The aphidophagous predator *Harmonia axyridis*
 1755 (Coleoptera : Coccinellidae) in Greece, 1994-1999. *European Journal of Entomology* 105:541-544
 1756 Koyama S, Majerus MEN (2008) Interactions between the parasitoid wasp *Dinocampus coccinellae*
 1757 and two species of coccinellid from Japan and Britain. *BioControl* 53:253-264
 1758 Krawfur E, Kring T, Miller J, et al. (1997) Gene Flow in the Exotic Colonizing Ladybeetle *Harmonia*
 1759 *axyridis* in North America. *Biological Control* 8:207-214
 1760 Krengel S, Stangl GI, Brandsch C, et al. (2012) A Comparative Study on Effects of Normal Versus
 1761 Elevated Temperatures During Preimaginal and Young Adult Period on Body Weight and Fat Body
 1762 Content of Mature *Coccinella septempunctata* and *Harmonia axyridis* (Coleoptera: Coccinellidae).
 1763 *Environmental entomology* 41:676-687
 1764 Kulijer D (2010) First record of invasive species *Harmonia axyridis* (Pallas, 1773)(Coleoptera:
 1765 Coccinellidae) in Bosnia and Herzegovina. *Acta entomologica serbica* 15:141-143
 1766 Kurosawa Y, S. H, Sasaji H (1985) The Coleoptera of Japan in color Vol. III. Hoikusha Publishing Co.,
 1767 Osaka (In Japanese).
 1768 LaMana ML, Miller JC (1996) Field Observations on *Harmonia axyridis* Pallas (Coleoptera:
 1769 Coccinellidae) in Oregon. *Biological Control* 6:232-237
 1770 LaRock DR, Mirdad Z, Ellington JJ, et al. (2003) Control of green peach aphids *Myzus persicae*
 1771 (Homoptera : Aphididae) with lady beetles *Harmonia axyridis* on Chile *Capsicum annum* (Coleoptera
 1772 : Coccinellidae) in the greenhouse. *Southw. Entomol.* 28:249-253

1773 Liere H, Kim TN, Werling BP, et al. (2014) Trophic cascades in agricultural landscapes: indirect effects
 1774 of landscape composition on crop yield. *Ecological Applications*
 1775 Lindborg R, Eriksson O (2004) Historical landscape connectivity affects present plant species
 1776 diversity. *Ecology* 85:1840-1845
 1777 Loiseau A, Malausa T, Lombaert E, et al. (2009) Isolation and characterization of microsatellites in
 1778 the harlequin ladybird, *Harmonia axyridis* (Coleoptera, Coccinellidae), and cross-species
 1779 amplification within the family Coccinellidae. *Mol. Ecol. Resour.* 9:934-937
 1780 Lombaert E, Estoup A, Facon B, et al. (2014) Rapid increase in dispersal during range expansion in
 1781 the invasive ladybird *Harmonia axyridis*. *Journal of evolutionary biology* 27:508-517
 1782 Lombaert E, Guillemaud T, Cornuet J-M, et al. (2010) Bridgehead Effect in the Worldwide Invasion of
 1783 the Biocontrol Harlequin Ladybird. *Plos One* 5:e9743
 1784 Lombaert E, Guillemaud T, Thomas CE, et al. (2011) Inferring the origin of populations introduced
 1785 from a genetically structured native range by approximate Bayesian computation: case study of the
 1786 invasive ladybird *Harmonia axyridis*. *Molecular Ecology* 20:4654-4670
 1787 Lombaert E, Malausa T, Devred R, et al. (2008) Phenotypic variation in invasive and biocontrol
 1788 populations of the harlequin ladybird, *Harmonia axyridis*. *BioControl* 53:89-102
 1789 Loreau M, Mouquet N, Gonzalez A (2003) Biodiversity as spatial insurance in heterogeneous
 1790 landscapes. *Proceedings of the National Academy of Sciences* 100:12765-12770
 1791 Losey JE, Allee LL, Stephens E, et al. (2014) Lady beetles in New York: Insidious invasions, erstwhile
 1792 extirpations, and recent rediscoveries. *Northeastern Naturalist* 21:271-284
 1793 Losey JE, Perlman JE, Hoebeke ER (2007) Citizen scientist rediscovers rare nine-spotted lady beetle,
 1794 *Coccinella novemnotata*, in eastern North America. *J Insect Conserv* 11:415-417
 1795 Lucas E, Vincent C, Labrie G, et al. (2007) The multicolored Asian ladybeetle *Harmonia axyridis*
 1796 (Coleoptera : Coccinellidae) in Quebec agroecosystems ten years after its arrival. *European Journal*
 1797 *of Entomology* 104:737-743
 1798 Luebert F, Plischoff P (2006) Sinopsis bioclimática y vegetal de Chile. Editorial Universitaria
 1799 Maes S, Massart X, Grégoire J-C, et al. (2014) Dispersal potential of native and exotic predatory
 1800 ladybirds as measured by a computer-monitored flight mill. *BioControl* 59:415-425
 1801 Majerus M, Strawson V, Roy H (2006) The potential impacts of the arrival of the harlequin ladybird,
 1802 *Harmonia axyridis* (Pallas) (Coleoptera : Coccinellidae), in Britain. *Ecological Entomology* 31:207-215
 1803 Majzlan O (2008) Faunistical contributions from Slovakia, Coleoptera, 4. *Naturae Tutela* 12 12:207–
 1804 210
 1805 McClure MS (1987) Potential of the Asian predator, *Harmonia axyridis* Pallas (Coleoptera:
 1806 Coccinellidae), to control *Matsucoccus resinosae* bean and godwin (Homoptera: Margarodidae) in
 1807 the United States. *Environmental Entomology* 16:224-230
 1808 Michaud JP (2002) Invasion of the Florida citrus ecosystem by *Harmonia axyridis* (Coleoptera :
 1809 Coccinellidae) and asymmetric competition with a native species, *Cycloneda sanguinea*.
 1810 *Environmental Entomology* 31:827-835
 1811 Michie LJ, Mallard F, Majerus MEN, et al. (2010) Melanic through nature or nurture: genetic
 1812 polymorphism and phenotypic plasticity in *Harmonia axyridis*. *Journal of Evolutionary Biology*
 1813 23:1699-1707
 1814 Mizell III RF (2007) Impact of *Harmonia axyridis* (Coleoptera: Coccinellidae) on native arthropod
 1815 predators in pecan and crape myrtle. *Florida entomologist* 90:524-536
 1816 Murchie AK, Moore JP, Moore GA, et al. (2008) The harlequin ladybird (*Harmonia axyridis* (Pallas))
 1817 (Coleoptera: Coccinellidae), found in Ireland. *Ir Nat J* 29:25–26
 1818 Murray K, Roy HE, Tinsley M (2015) Assessing the role of host immunity in harlequin ladybird enemy
 1819 release. In: Brown PMJ, Roy HE, Howe A and De Clercq P (eds) 3rd Meeting of IOBC-WPRS study
 1820 group “Benefits and risks of exotic biological control agents” 13 to 15 May 2015. Bornholm,
 1821 Denmark,
 1822 Nalepa CA (2007) *Harmonia axyridis* (Coleoptera : Coccinellidae) in buildings: Relationship between
 1823 body height and crevice size allowing entry. *Journal of Economic Entomology* 100:1633-1636

1824 Nalepa CA, Kennedy GG, Brownie C (2005) Role of visual contrast in the alighting behavior of
1825 *Harmonia axyridis* (Coleoptera : Coccinellidae) at overwintering sites. Environmental Entomology
1826 34:425-431

1827 Nalepa CA, Kidd KA, Hopkins DI (2000) The Multicolored Asian lady beetle (Coleoptera :
1828 Coccinellidae): Orientation to aggregation sites. J. Entomol. Sci. 35:150-157

1829 Nalepa CA, Weir A (2007) Infection of *Harmonia axyridis* (Coleoptera : Coccinellidae) by
1830 *Hesperomyces virescens* (Ascomycetes : Laboulbeniales): Role of mating status and aggregation
1831 behavior. J. Invertebr. Pathol. 94:196-203

1832 Nedvěd O (2014) Slunéčko východní (*Harmonia axyridis*) – pomocník v biologické ochraně nebo
1833 ohrožení biodiverzity? Jihočeská univerzita, České Budějovice, 65 pp.

1834 Nedvěd O, Háva J (2015) New record of the invasive lady beetle *Harmonia axyridis* in Afrotropical
1835 Region: Tanzania, Zanzibar African Entomology 24

1836 Nedvěd O, Háva J (2016) New record of the invasive lady beetle *Harmonia axyridis* in Afrotropical
1837 Region: Tanzania, Zanzibar. African Entomology 24:1-4

1838 Nedvěd O, Háva J, Kulíková D (2011) Record of the invasive alien ladybird *Harmonia axyridis*
1839 (Coleoptera, Coccinellidae) from Kenya. ZooKeys:77-81

1840 Nedved O, Kalushkov P (2011) Effect of air humidity on sex ratio and development of ladybird
1841 *Harmonia axyridis* (Coleoptera: Coccinellidae). Psyche: A Journal of Entomology 2012:173482

1842 Nedvěd O, Krejčík S (2010) Record of the ladybird *Harmonia axyridis* (Coleoptera: Coccinellidae) from
1843 Uruguay. Klapalekiana 46:203-204

1844 Nekrasova OD, Tytar VM (2014) Longstanding and seasonal population dynamics of the invasive
1845 species *Harmonia axyridis* (Coleoptera, Coccinellidae) in Ukraine The Journal of V.N.Karazin Kharkiv
1846 National University. Series: biology 20:159-162

1847 Nóia M, Borges I, Soares AO (2008) Intraguild predation between the aphidophagous ladybird
1848 beetles *Harmonia axyridis* and *Coccinella undecimpunctata* (Coleoptera: Coccinellidae): the role of
1849 intra and extraguild prey densities. Biological Control 46:140-146

1850 Obata S (1986) Determination of hibernation site in the ladybird beetle, *Harmonia axyridis* Pallas
1851 (Coleoptera: Coccinellidae). . Kontyû 54:218-233

1852 Obrycki JJ, Giles KL, Ormord AM (1998) Experimental assessment of interactions between larval
1853 *Coleomegilla maculata* and *Coccinella septempunctata* (Coleoptera: Coccinellidae) in field cages.
1854 Environmental Entomology 27:1280-1288

1855 Obrycki JJ, Harwood JD, Kring TJ, et al. (2009) Aphidophagy by Coccinellidae: application of biological
1856 control in agroecosystems. Biological Control 51:244-254

1857 Ohgushi T, Sawada H (1998) What changed the demography of an introduced population of an
1858 herbivorous lady beetle? Journal of animal ecology 67:679-688

1859 Orlova-Bienkowskaja MJ (2013) Dangerous invasive harlequin ladybird *Harmonia axyridis* (Pallas,
1860 1773)(Coleoptera, Coccinellidae) in the European Russia. Russian journal of biological invasions
1861 4:190-193

1862 Orlova-Bienkowskaja MJ (2014) The outbreak of harlequin ladybird *Harmonia axyridis* (Pallas,
1863 1773)(Coleoptera, Coccinellidae) in the Caucasus and possible sources of invasion. Russian Journal of
1864 Biological Invasions 5:275-281

1865 Orlova-Bienkowskaja MJ, Ukrainsky AS, Brown PM (2015) *Harmonia axyridis* (Coleoptera:
1866 Coccinellidae) in Asia: a re-examination of the native range and invasion to southeastern Kazakhstan
1867 and Kyrgyzstan. Biological Invasions:1-8

1868 Osawa N (2000) Population field studies on the aphidophagous ladybird beetle *Harmonia axyridis*
1869 (Coleoptera: Coccinellidae): resource tracking and population characteristics. Population Ecology
1870 42:115-127

1871 Osawa N (2011) Ecology of *Harmonia axyridis* in natural habitats within its native range. BioControl
1872 56:613-621

1873 Osawa N, Nishida T (1992) Seasonal variation in elytral color polymorphism in *Harmonia axyridis* (the
1874 ladybird beetle) - the role of nonrandom mating. Heredity 69:297-307

1875 Pallas PS (1771) Reise durch verschiedene Provinzen des russischen Reichs. gedruckt bey der
 1876 Kayserlichen Academie der Wissenschaften

1877 Panigaj L, Zach P, Honěk A, et al. (2014) The invasion history, distribution and colour pattern forms of
 1878 the harlequin ladybird beetle *Harmonia axyridis* (Pall.)(Coleoptera, Coccinellidae) in Slovakia, Central
 1879 Europe. ZooKeys:89

1880 Pedersen J, Runge JB, Jonsén BP (2008) Fund af biller i Danmark, 2006 og 2007 (Coleoptera).
 1881 Entomologiske Meddelelser 76:105-144

1882 Pell JK, Baverstock J, Roy HE, et al. (2008) Intraguild predation involving *Harmonia axyridis*: a review
 1883 of current knowledge and future perspectives. BioControl 53:147-168

1884 Pervez A, Omark (2006) Ecology and biological control application of multicoloured Asian ladybird,
 1885 *Harmonia axyridis*: A review. Biocontrol Science and Technology 16:111-128

1886 Phillips BL, Brown GP, Greenlees M, et al. (2007) Rapid expansion of the cane toad (*Bufo marinus*)
 1887 invasion front in tropical Australia. Austral Ecology 32:169-176

1888 Phillips BL, Brown GP, Shine R (2010) Life-history evolution in range-shifting populations. Ecology
 1889 91:1617-1627

1890 Pickering GJ, Blake AJ, Soleas GJ, et al. (2010) Remediation of wine with elevated concentrations of
 1891 3-alkyl-2-methoxypyrazines using cork and synthetic closures. Journal of Food, Agriculture &
 1892 Environment 8:97-101

1893 Pocock MJO, Roy HE, Preston CD, et al. (2015) The Biological Records Centre: a pioneer of citizen
 1894 science. Biological Journal of the Linnean Society 115:475-493

1895 Pons X, Roca M, Lumbierres B, et al. (2015) Characterization of a newly established aggregation of
 1896 the invasive ladybeetle *Harmonia axyridis* and current status of the invader in Spain. Spanish Journal
 1897 of Agricultural Research 13:1006

1898 Poutsma J, Loomans AJM, Aukema B, et al. (2008) Predicting the potential geographical distribution
 1899 of the harlequin ladybird, *Harmonia axyridis*, using the CLIMEX model. BioControl 53:103-125

1900 Przewozny M, Barlozek T, Bunalski M (2007) *Harmonia axyridis* (Pallas, 1773) (Coleoptera :
 1901 Coccinellidae) new species of ladybird beetle for Polish fauna. Polish Journal of Entomology 76:177-
 1902 182

1903 Purse BV, Comont R, Butler A, et al. (2014) Landscape and climate determine patterns of spread for
 1904 all colour morphs of the alien ladybird *Harmonia axyridis*. Journal of Biogeography 42:575-588

1905 Raak-van den Berg C, Hemerik L, de Jong PW, et al. (2012) Mode of overwintering of invasive
 1906 *Harmonia axyridis* in the Netherlands. BioControl 57:71-84

1907 Raak-van den Berg C, van Wielink PS, de Jong PW, et al. (2014) Invasive alien species under attack:
 1908 natural enemies of *Harmonia axyridis* in the Netherlands. BioControl 59:229-240

1909 Rabitsch W, Schuh R (2006) First record of the multicoloured Asian ladybird *Harmonia axyridis*.
 1910 Pallas:161-164

1911 Ragsdale DW, Landis DA, Brodeur J, et al. (2011) Ecology and management of the soybean aphid in
 1912 North America. Annual review of entomology 56:375-399

1913 Ramsey S, Losey JE (2012) Why is *Harmonia axyridis* the culprit in coccinellid biting incidents? An
 1914 analysis of means, motive, and opportunity. American Entomologist 58:166-170

1915 Reznik SY, Dolgovskaya MY, Ovchinnikov A, et al. (2015) Weak photoperiodic response facilitates the
 1916 biological invasion of the harlequin ladybird *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae).
 1917 Journal of Applied Entomology 139:241-249

1918 Rhule EL, Majerus ME, Jiggins FM, et al. (2010) Potential role of the sexually transmitted mite
 1919 *Coccipolipus hippodamiae* in controlling populations of the invasive ladybird *Harmonia axyridis*.
 1920 Biological Control 53:243-247

1921 Ribbands B, Brown PMJ, Roy HE, et al. (2009) The most northerly record of the Harlequin ladybird
 1922 (Col., Coccinellidae) in the British Isles. Entomologist's Monthly Magazine 145:43-44

1923 Richardson JV, Deloach C, J. (1972) Some aspects of host selection by *Perilitus coccinellae*
 1924 Hymenoptera: Braconidae. Annals of the Entomological Society of America 65:834

1925 Riddick EW (2006) Influence of host gender on infection rate, density and distribution of the parasitic
 1926 fungus, *Hesperomyces virescens*, on the multicolored Asian lady beetle, *Harmonia axyridis*. J Insect
 1927 Sci. 6:15

1928 Riddick EW, Cottrell TE, Kidd KA (2009) Natural enemies of the Coccinellidae: Parasites, pathogens,
 1929 and parasitoids. Biological Control 51:306-312

1930 Rondoni G, Athey KJ, Harwood JD, et al. (2014) Development and application of molecular gut-
 1931 content analysis to detect aphid and coccinellid predation by *Harmonia axyridis* (Coleoptera:
 1932 Coccinellidae) in Italy. Insect science

1933 Rondoni G, Comont RF, Ricci C, et al. (2013a) Parasitisation of coccinellid species in Oxfordshire and
 1934 London and first record of an *Harmonia axyridis* – phorid association from Italy. International
 1935 Symposium "Ecology of Aphidophaga 12". Belgrade, Serbia, pp. 121

1936 Rondoni G, Comont RF, Ricci C, et al. (2013b) Parasitisation of coccinellid species in Oxfordshire and
 1937 London and first record of an *Harmonia axyridis* – phorid association from Italy. International
 1938 Symposium "Ecology of Aphidophaga 12", Belgrade, Serbia, p. 121.

1939 Roy H, Baverstock J, Ware R, et al. (2008a) Intraguild predation of the aphid pathogenic fungus
 1940 *Pandora neoaphidis* by the invasive coccinellid *Harmonia axyridis*. Ecological Entomology 33:175-182

1941 Roy H, Pell J, Clark S, et al. (1998) Implications of predator foraging on aphid pathogen dynamics. J.
 1942 Invertebr. Pathol. 71:236-247

1943 Roy H, Pocock M, Preston C, et al. (2012a) Understanding citizen science and environmental
 1944 monitoring: final report on behalf of UK Environmental Observation Framework.

1945 Roy HE, Adriaens T, Isaac NJB, et al. (2012b) Invasive alien predator causes rapid declines of native
 1946 European ladybirds. Diversity and Distributions 18:717-725

1947 Roy HE, Adriaens T, Isaac NJB, et al. (2012c) Invasive alien predator causes rapid declines of native
 1948 European ladybirds. Diversity and Distributions 18:717-725

1949 Roy HE, Bacon J, Beckmann B, et al. (2012d) Non-Native Species in Great Britain: establishment,
 1950 detection and reporting to inform effective decision making. DEFRA, London,

1951 Roy HE, Baverstock J, Ware RL, et al. (2008b) Intraguild predation of the aphid pathogenic fungus
 1952 *Pandora neoaphidis* by the invasive coccinellid *Harmonia axyridis*. Ecological Entomology 33:175-182

1953 Roy HE, Brown PMJ (2015a) Ten years of invasion: *Harmonia axyridis* (Pallas) (Coleoptera:
 1954 Coccinellidae) in Britain. Ecological Entomology:n/a-n/a

1955 Roy HE, Brown PMJ (2015b) Ten years of invasion: *Harmonia axyridis* (Pallas) (Coleoptera:
 1956 Coccinellidae) in Britain. Ecological Entomology

1957 Roy HE, Brown PMJ, Frost R, et al. (2011a) Atlas of the ladybirds (Coccinellidae) of Britain and
 1958 Ireland. Biological Records Centre, Wallingford, UK.

1959 Roy HE, Brown PMJ, Rothery P, et al. (2008c) Interactions between the fungal pathogen *Beauveria*
 1960 *bassiana* and three species of coccinellid: *Harmonia axyridis*, *Coccinella septempunctata* and *Adalia*
 1961 *bipunctata*. BioControl 53:265-276

1962 Roy HE, Handley L-J, Schoenrogge K, et al. (2011b) Can the enemy release hypothesis explain the
 1963 success of invasive alien predators and parasitoids? BioControl 56:451-468

1964 Roy HE, Lawson Handley L-J (2012) Networking: a community approach to invaders and their
 1965 parasites. Functional Ecology 26:1238–1248

1966 Roy HE, Preston CD, Roy DB (2015a) Foreword: Fifty years of the Biological Records Centre. Biological
 1967 Journal of the Linnean Society 115:469-474

1968 Roy HE, Rhule E, Harding S, et al. (2011c) Living with the enemy: parasites and pathogens of the
 1969 ladybird *Harmonia axyridis*. BioControl 56:663-679

1970 Roy HE, Rorke SL, Beckmann B, et al. (2015b) The contribution of volunteer recorders to our
 1971 understanding of biological invasions. Biological Journal of the Linnean Society 115:678-689

1972 Roy HE, Wajnberg E (2008) From biological control to invasion: the ladybird *Harmonia axyridis* as a
 1973 model species. BioControl 53:1-4

1974 Sæthre M-G, Staverløkk A, Hofsvang T (2010) The history of *Harmonia axyridis* (Pallas 1773) in
 1975 Norway. . IOBC/wprs Bulletin 58:97-104

1976 Saethre MG, Staverløkk A, Hofsvang T (2010) The history of *Harmonia axyridis* (Pallas 1773) in
1977 Norway. IOBC/wprs Bulletin (Working Group “Benefits and Risks of exotic biological Control Agents”)
1978 58:97-104

1979 Saini ED (2004) Presencia de *Harmonia axyridis* (Pallas)(Coleoptera: Coccinellidae) en la provincia de
1980 Buenos Aires. Aspectos biológicos y morfológicos. Instituto Nacional de Tecnología Agropecuaria
1981 Savojkaja GI (1971) On the acclimatization of *Leis axyridis* Pallas (Coleoptera, Coccinellidae) in Ile-
1982 Alatau. Trudy XIII Mezhdunarodnogo entomologicheskogo kongressa 2:181-182

1983 Schanderl H, Ferran A, Coderre D, et al. (1992) Capacidade de dispersão de *Harmonia axyridis*
1984 PALLAS (Col., Coccinellidae) após largada inundativa para controle de afídeos do milho,
1985 *Rhopalosiphum padi* L. e *Sitobion avenae* (F.)(Hom., Aphididae).

1986 Sebastião D, Borges I, Soares AO (2015) Effect of temperature and prey in the biology of *Scymnus*
1987 *subvillosus*. BioControl 60:241-249

1988 Seko T, Sumi A, Nakano A, et al. (2014) Suppression of aphids by augmentative release of larvae of
1989 flightless *Harmonia axyridis*. Journal of Applied Entomology 138:326-337

1990 Shinner R (2014) Thermal biology of the invasive ladybird *Harmonia axyridis*: physiological plasticity,
1991 thermoregulatory behaviour and fitness. MSc thesis, Stellenbosch University, South Africa.

1992 Sighinolfi L, Febvay G, Dindo ML, et al. (2008) Biological and biochemical characteristics for quality
1993 control of *Harmonia axyridis* (Pallas)(Coleoptera, Coccinellidae) reared on a liver-based diet. Archives
1994 of insect biochemistry and physiology 68:26-39

1995 Silvertown J, Harvey M, Greenwood R, et al. (2015) Crowdsourcing the identification of organisms: A
1996 case-study of iSpot. ZooKeys:125

1997 Sloggett JJ (2010) Predation of ladybird beetles by the orb-web spider *Araneus diadematus*.
1998 BioControl 55:631-638

1999

2000 Sloggett JJ (2012) *Harmonia axyridis* invasions: Deducing evolutionary causes and consequences.
2001 Entomol. Sci. 15:261-273

2002 Sloggett JJ, Obrycki JJ, Haynes KF (2009) Identification and quantification of predation: novel use of
2003 gas chromatography–mass spectrometric analysis of prey alkaloid markers. Functional ecology
2004 23:416-426

2005 Smith CA, Gardiner MM (2013) Oviposition habitat influences egg predation of native and exotic
2006 coccinellids by generalist predators. Biological Control 67:235-245

2007 Snyder WE (2009) Coccinellids in diverse communities: Which niche fits? Biological Control 51:323-
2008 335

2009 Snyder WE, Clevenger GM, Eigenbrode SD (2004) Intraguild predation and successful invasion by
2010 introduced ladybird beetles. Oecologia 140:559-565

2011 Soares AO, Borges I, Borges PAV, et al. (2008) *Harmonia axyridis*: what will stop the invader?
2012 BioControl 53:127-145

2013 Soares AO, Coderre D, Schanderl H (2001) Fitness of two phenotypes of *Harmonia axyridis*
2014 (Coleoptera : Coccinellidae). European Journal of Entomology 98:287-293

2015 Soares AO, Coderre D, Schanderl H (2003) Effect of temperature and intraspecific allometry on
2016 predation by two phenotypes of *Harmonia axyridis* Pallas (Coleoptera : Coccinellidae).
2017 Environmental Entomology 32:939-944

2018 Soares AO, Coderre D, Schanderl H (2004) Dietary self-selection behaviour by the adults of the
2019 aphidophagous ladybeetle *Harmonia axyridis* (Coleoptera : Coccinellidae). Journal of Animal Ecology
2020 73:478-486

2021 Soares AO, Serpa A (2007) Interference competition between ladybird beetle adults (Coleoptera:
2022 Coccinellidae): effects on growth and reproductive capacity. Population ecology 49:37-43

2023 Solano Y, Arcaya E (2014) Primer registro de *Harmonia axyridis* (Pallas, 1773) (Coleoptera:
2024 Coccinellidae) en Venezuela. Entomotropica 29:57-61

2025 Solter LF, Kyei-Poku GK, Johny S (2013) Comment on “Invasive harlequin ladybird carries biological
2026 weapons against native competitors”. Science 341:1342-1342

2027 Špyňar P (2008) Faunistic records from the Czech Republic – 252. Klapalekiana 44:77-79
 2028 Stals R (2008) Invasive ladybird spreading in South Africa. Biocontrol News Inf 29:24N–26N
 2029 Stals R (2010) The establishment and rapid spread of an alien invasive lady beetle: *Harmonia axyridis*
 2030 (Coleoptera: Coccinellidae) in southern Africa. IOBC/wprs Bulletin 58:125-132
 2031 Stals R, Prinsloo G (2007) Discovery of an alien invasive, predatory insect in South Africa: the
 2032 Multicoloured Asian ladybird beetle, *Harmonia axyridis* (Pallas) (Coleoptera : Coccinellidae). S. Afr. J.
 2033 Sci. 103:123-126
 2034 Statens landbrukstilsyn (2001) Assessment of *Harmonia axyridis*. Internal document.
 2035 Staverløkk A (2006) Fremmede arter og andre uønskede blindpassasjerer i import av
 2036 grøntanleggsplanter. (Occurrence of alien species and other unwanted stowaways in imported
 2037 horticultural plants). 111 pp. M. Sc. thesis, University of Life Sciences (UMB), Ås, Norway.
 2038 Steenberg T, Harding S (2009a) Entomopathogenic fungi recorded from the Harlequin ladybird,
 2039 *Harmonia axyridis*. J. Invertebr. Pathol. 102:88-89
 2040 Steenberg T, Harding S (2009b) The harlequin ladybird (*Harmonia axyridis* Pallas) in Denmark: spread
 2041 and phenology during the initial phase of invasion. Entomologiske Meddelelser 77:27-39
 2042 Steenberg T, Harding S, Bønløkke J, et al. (2009) En ny art invaderer. Aktuell Naturvidenskab 3:4-7
 2043 Tan C-C, Li J-C (1934) Inheritance of the elytral color patterns of the lady-bird beetle, *Harmonia*
 2044 *axyridis* Pallas. The American Naturalist 68:252-265
 2045 Tan CC (1946) Mosaic dominance in the inheritance of color patterns in the lady-bird beetle,
 2046 *Harmonia axyridis*. Genetics 31:195
 2047 Tang B, Zhu J, Guo HS, et al. (2012) Studies of the diversity of multiple elytral color morphs of
 2048 *Harmonia axyridis* (Pallas) (Coleoptera: Coccinellidae). Journal of Hangzhou Normal University
 2049 11:132-136
 2050 Tang DL, Qiu BL, Ren SX (2014) A review of insecticide resistance in the natural enemies of pest
 2051 insects. Chinese Journal of Applied Entomology 51:13-25
 2052 Tayeh A, Estoup A, Hufbauer R, et al. (2013) Investigating the genetic load of an emblematic invasive
 2053 species: the case of the invasive harlequin ladybird *Harmonia axyridis*. Ecology and evolution 3:864-
 2054 871
 2055 Tayeh A, Estoup A, Laugier G, et al. (2012) Evolution in biocontrol strains: insight from the harlequin
 2056 ladybird *Harmonia axyridis*. Evolutionary applications 5:481-488
 2057 Tayeh A, Hufbauer RA, Estoup A, et al. (2015) Biological invasion and biological control select for
 2058 different life histories. Nature communications 6
 2059 Tedders WL, Schaefer PW (1994) Release and establishment of *Harmonia axyridis* (Coleoptera,
 2060 Coccinellidae) in the southeastern United States. Entomological News 105:228-243
 2061 Thomas AP, Trotman J, Wheatley A, et al. (2013) Predation of native coccinellids by the invasive alien
 2062 *Harmonia axyridis* (Coleoptera: Coccinellidae): detection in Britain by PCR-based gut analysis. Insect
 2063 conservation and diversity 6:20-27
 2064 Timuş A, Stahi N (2013) The acclimatization and trophic spectrum of *Harmonia axyridis* in the
 2065 Republic of Moldova. VIII international conference of zoologists. Actual problems of protection and
 2066 sustainable use of the animal world diversity (Chisinau, 10–12.10. 2013). Book of abstract. Academy
 2067 of Sciences of Moldova, Chisinau.
 2068 Toda Y, Sakuratani Y (2006) Expansion of the geographical distribution of an exotic ladybird beetle,
 2069 *Adalia bipunctata* (Coleoptera: Coccinellidae), and its interspecific relationships with native ladybird
 2070 beetles in Japan. Ecol. Res. 21:292-300
 2071 Tourniaire R, Ferran A, Giuge L, et al. (2000) A natural flightless mutation in the ladybird, *Harmonia*
 2072 *axyridis*. Entomologia Experimentalis et Applicata 96:33-38
 2073 Travis JM, Dytham C (2002) Dispersal evolution during invasions. Evolutionary Ecology Research
 2074 4:1119-1129
 2075 Turgeon J, Tayeh A, Facon B, et al. (2011) Experimental evidence for the phenotypic impact of
 2076 admixture between wild and biocontrol Asian ladybird (*Harmonia axyridis*) involved in the European
 2077 invasion. Journal of Evolutionary Biology 24:1044-1052

2078 Ukrainsky AS, Orlova-Bienkowskaja MJ (2014) Expansion of *Harmonia axyridis* Pallas (Coleoptera:
 2079 Coccinellidae) to European Russia and adjacent regions. *Biological invasions* 16:1003-1008
 2080 van Wielink P, Spijkers H (2013) Insects nightly attracted to light at a single site in De Kaaistoep, The
 2081 Netherlands. Orders, families and species identified in 1995–2011. *Entomol Ber (Amst)* 73:200-214
 2082 Vandereycken A, Brostaux Y, Joie E, et al. (2013) Occurrence of *Harmonia axyridis* (Coleoptera:
 2083 Coccinellidae) in field crops. *European Journal of Entomology* 110:285-292
 2084 Varenhorst AJ, O'Neal ME (2012) The Response of Natural Enemies to Selective Insecticides Applied
 2085 to Soybean. *Environmental entomology* 41:1565-1574
 2086 Veran S, Piry S, Ternois V, et al. (2015) Modeling spatial expansion of invasive alien species: relative
 2087 contributions of environmental and anthropogenic factors to the spreading of the harlequin ladybird
 2088 in France. *Ecography*
 2089 Verheggen FJ, Fagel Q, Heuskin S, et al. (2007) Electrophysiological and behavioral responses of the
 2090 Multicolored Asian Lady Beetle, *Harmonia axyridis* Pallas, to sesquiterpene semiochemicals. *Journal*
 2091 *of Chemical Ecology* 33:2148-2155
 2092 Verizhnikova I (2011) Invasion by *Harmonia axyridis* (Coleoptera: Coccinellidae): population growth
 2093 in Kyiv region. *Bull. Kharkiv Nat. Agr. Univ.(Phytopathol. Entomol.)* 9:23-26
 2094 Verizhnikova I, Shylova K (2013a) Consequences of entomophage *Harmonia axyridis* Pall. Coleoptera,
 2095 Coccinellidae) introduction and its predicted area of acclimatization. *Urgent Ecol Env Manag Probl*
 2096 *PFUR Moscow* 15:65-68
 2097 Verizhnikova I, Shylova K (2013b) Consequences of entomophage *Harmonia axyridis* Pallas
 2098 (Coleoptera, Coccinellidae) introduction and its predicted area of acclimatization. *Urgent Ecol Env*
 2099 *Manag Probl PFUR Moscow* 15:65-68
 2100 Veselý P, Ernestová B, Nedvěd O, et al. (in press) Topsy-turvy aposematism: sparrows eat warning
 2101 and distasteful prey. *Proceedings of the Royal Society B: Biological Sciences*
 2102 Vilcinskis A, Stoecker K, Schmidtberg H, et al. (2013) Invasive Harlequin Ladybird Carries Biological
 2103 Weapons Against Native Competitors. *Science* 340:862
 2104 Vorontsov N, Blekhan A (2001) Geographical range and intraspecific structure of the ladybird
 2105 *Harmonia axyridis* Pall., 1773 (Coleoptera, Coccinellidae). *Evolution, ecology and biodiversity.*
 2106 *Proceedings of the Conference in memory of Nikolai Nikolaevich Vorontsov (1934–2000). Uchebno-*
 2107 *nauchnyj tsent dovuzovskogo obrazovaniya, Moscow.* pp. 150-156
 2108 Wang S, Michaud J, Tan X, et al. (2011) The aggregation behavior of *Harmonia axyridis* in its native
 2109 range in Northeast China. *BioControl* 56:193-206
 2110 Wang S, Michaud JP, Zhang RZ, et al. (2009) Seasonal cycles of assortative mating and reproductive
 2111 behaviour in polymorphic populations of *Harmonia axyridis* in China. *Ecological Entomology* 34:483-
 2112 494
 2113 Wang S, Zhang RZ, Zhang F (2007a) Research progress of biology and ecology of *Harmonia axyridis*
 2114 (Pallas) (Coleoptera: Coccinellidae). *Chinese Journal of Applied Ecology* 18:2117-2126
 2115 Wang S, Zhang RZ, Zhang F (2007b) Research progress on biology and ecology of *Harmonia axyridis*
 2116 Pallas (Coleoptera: Coccinellidae). *Chinese Journal of Applied Ecology* 18:2117-2126
 2117 Ware RL, Majerus MEN (2008) Intraguild predation of immature stages of British and Japanese
 2118 coccinellids by the invasive ladybird *Harmonia axyridis*. *BioControl* 53:169-188
 2119 Ware RL, Yguel B, Majerus MEN (2009) Effects of competition, cannibalism and intra-guild predation
 2120 on larval development of the European coccinellid *Adalia bipunctata* and the invasive species
 2121 *Harmonia axyridis*. *Ecological Entomology* 34:12-19
 2122 Weissenberger A, Brun J, Piotte C, et al. (1999) Comparison between the wild type and flightless type
 2123 of the coccinellid *Harmonia axyridis* (Pallas) in control of the damson hop aphid. *Fifth International*
 2124 *Conference on Pests in Agriculture. Association Nationale pour la Protection des Plantes (ANPP).* pp.
 2125 727-734
 2126 Welch VL, Sloggett JJ, Webberley KM, et al. (2001) Short-range clinal variation in the prevalence of a
 2127 sexually transmitted fungus associated with urbanisation. *Ecological Entomology* 26:547-550

2128 Wells PM (2011) Intraguild predation by *Harmonia axyridis*: effects on native enemies and aphid
 2129 suppression. Department of Zoology. University of Cambridge,
 2130 Wheeler Jr A, Hoebeke E (1995) *Coccinella novemnotata* in northeastern North America: historical
 2131 occurrence and current status (Coleoptera: Coccinellidae). Entomological Society of Washington
 2132 (USA)
 2133 Woltz J, Landis D (2014) Comparison of sampling methods of *Aphis glycines* predators across the diel
 2134 cycle. Journal of applied entomology 138:475-484
 2135 Yang YW, Zhang QM, Du L, et al. (2006) Study on selective preferences of *Harmonia axyridis* (Pallas)
 2136 collected from different habitats to predating aphids. China Plant Protection 26:5-7
 2137 Yasuda H, Evans EW, Kajita Y, et al. (2004) Asymmetric larval interactions between introduced and
 2138 indigenous ladybirds in North America. Oecologia 141:722-731
 2139 Zakharov IA, Goryacheva II, Suvorov A (2011) Mitochondrial DNA polymorphism in invasive and
 2140 native populations of *Harmonia axyridis*. European Journal of Environmental Sciences 1
 2141 Zenni RD, Nuñez MA (2013) The elephant in the room: the role of failed invasions in understanding
 2142 invasion biology. Oikos 122:801-815

2143

2144

2145

2146 Figure 1 Global distribution of *Harmonia axyridis*.
2147 Figure 2 European distribution of *Harmonia axyridis*.
2148
2149
2150
2151
2152

Table 1: Invasion characteristics and ecological attributes of *Harmonia axyridis* documented from countries around the world. It is important to note that various approaches have been adopted for categorising the attributes and so direct comparison between countries is not always possible. Standard approaches to capturing this information would be useful to enable quantitative assessments on a global scale. Natural enemies have not been systematically surveyed in most countries and so the lists represent opportunistic observations. The origin is often inferred from geographic proximity to other countries but in some cases through molecular analysis. Spread has been measured as distance (km per year) for some countries. Broad habitat categories are provided but harmonization of terminology is recommended. Diet breadth has not been systematically analysed for each country and the lists represent opportunistic observations. Taxonomy for species is provided at first mention within the main text other than Aphididae: *Aphis craccivora* Koch, *Aphis fabae* Scopoli, *Chaitophorus leucomelas* Koch, *Chromaphis juglandicola* (Kaltenbach), *Eucallipterus tiliæ* (L.); Araneidae: *Araneus diadematus* Clerck (*Chrysoperla carnea* (Stephens) (Chrysopidae); *Homalotylus flaminus* (Dalman) (Encyrtidae); *Medina separata* (Meigen) (Tachinidae); *Metcalfa pruinosa* (Say) (Flatidae); *Oomyzus scaposus* (Thomson) (Eulophidae); *Passer montanus* (L.) (Passeridae)

	Origin	Establishment	Spread	Habitat breadth	Diet breadth	Natural enemies
China	Native	Native	Native	Wide Agricultural Grasslands Woodlands	Wide Aphids Mites Coccids Pollen Young plant tissues	Braconidae <i>Medina luctuosa</i> <i>Phalacrotophora philaxyridis</i> <i>Hesperomyces virescens</i> <i>Strongygaster triangulifera</i> Birds
Japan	Native	Native	Native	Wide Urban habitats Parks Backyards Agricultural Gardens (mainly on trees and shrubs)	Wide Aphids Ladybird larvae	<i>Dinocampus coccinellae</i> <i>Phalacrotophora</i> sp.

Siberia and the Far East	Native	Native	Native	Wide Deciduous Coniferous forests Gardens Urban habitats Grasslands Fields	Wide Aphids Coccids Psyllids Coccinellids	Birds <i>Dinocampus coccinellae</i>
European Russia	Multiple intentional releases in the Caucasus starting in 1930s (not established). First population seen in wild in 2010 in the west. Invasive population likely secondary spread from Poland	Established in the west and in the south. Individual specimens found in centre of European Russia	Range expansion in the south estimated at 300 km/year. Expansion to the central regions is not yet known.	Wide Urban habitats Forests Fields Gardens Floodplains	Wide Aphids Coccids Psyllids Coccinellids	Unknown
North America	Multiple intentional releases across North America, starting California in 1916. Introduced into North America throughout 1980s. First population seen in wild in 1988	Found across USA by mid-1990s. First seen in Canada in 1994.	Initial range expansion in southeastern USA estimated at 309 km/year	Wide Agricultural Woodlands Grasslands Residential and urban landscapes	Wide Many phytophagous Hemiptera (i.e. aphids, psyllids and coccids) Tetranychidae Immature stages of Lepidoptera and Coleoptera (i.e. Coccinellidae, Chrysomelidae, Curculionidae) Fruit crops Pollen Nectar	<i>Coccipolipus hippodamiae</i> <i>Hesperomyces virescens</i> <i>Strongygaster triangulifera</i> <i>Dinocampus coccinellae</i>

South America (Chile)	Eastern North USA Intentional introduction of flightless strain from France in 1998 but this did not establish	Wild population of flying strain discovered in 2003	Expanded from a range of 250 km (north to south) near Santiago, to 2600 km (north to south); approximately at a rate of 160 km per year, mostly towards south	Agricultural (particularly abundant in alfalfa during spring and autumn but also present in wheat) Houses (autumn and winter) Vineyards Orchards Gardens Native matorral	Wide Mostly seen feeding on aphids <i>Aphis craccivora</i> <i>Acyrtosiphon pisum</i> <i>Chromaphis juglandicola</i> <i>Chaitophorus leucomelas</i>	<i>Dinocampus coccinellae</i>
Belgium	Intentional introduction as biocontrol agent in 1997	First wild population found in 2001	Range expansion of 189% on average per year (2002 to 2006)	Agricultural (dominant in corn) Urban and anthropogenic habitats Less abundant in (semi-) natural areas Predominantly on trees and shrubs but also on herbs	Wide	<i>Dinocampus coccinellae</i> <i>Hesperomyces virescens</i> Araneidae
Czech Republic	Intentional introduction as a biocontrol agent in early 2000s, however this population did not establish Invasive population likely secondary spread from Germany	First establishment seen in 2006	First invasive population seen in 2006, by 2009 it could be found in most disturbed habitats across the country	Urban Abundant on deciduous trees and shrubs but also ruderal herbs	Wide Mostly aphids Many psyllids e.g. <i>Cacopsylla pyri</i>	<i>Hesperomyces virescens</i> <i>Passer montanus</i> Other parasites and parasitoids only occasionally

Denmark	Unintentional introduction Presumed to have arrived from Germany	First recorded in 2006	Expanded from Copenhagen east and west, however establishment is limited in the north of the country.	Urban Agricultural Parks and gardens	Wide Aphids (<i>Eucallipterus tiliae</i>) Hemiptera (<i>Anthocoris nemoralis</i>)	<i>Beauveria bassiana</i> <i>Isaria farinosa</i> <i>Lecanicillium</i> spp.
France	Intentional introduction in 1982. <i>H. axyridis</i> sampled across France all belong to a single genetic unit that has invaded the Western part of Europe. It bears traces of genetic admixture between an eastern North American wild source, and a biocontrol strain used in Europe	Populations started to spread in 2004	Dispersal was greatly aided by anthropogenic factors, especially the relative surface area of urban habitats, and average summer temperatures not being too high or low	Urban Parks and gardens Agricultural	Wide Most aphids Many psyllids	Not recorded
Germany	Intentional introduction during end of 1990s. Unclear, if invasive populations derives from intentional introductions or spread.	Considered established since 2002	First reports from Frankfurt City in 2000 followed by rapid spread across Germany.	Agricultural Gardens Vineyards Orchards	Wide Many aphid species, e.g. on cereals (wheat, corn), <i>Aphis fabae</i> other plant sucking insects <i>Eriosoma lanigerum</i> on apple Grape phylloxera <i>Daktulosphaira viticolae</i> Pollen Fruit sap (e.g. from	<i>Hesperomyces virescens</i> <i>Dinocampus coccinellae</i> Allantonematidae

					grape berries)	
Italy	<p>Intentional introductions from 1995 to 1999.</p> <p>Presumed that wild populations originated from biocontrol releases or secondary spread from France, Switzerland and other parts of Italy</p>	Population first recorded as in wild in 2006	Expanded from first wild recorded individuals in 2006 to cover 18 of 20 Italian regions in 2015	<p>Agricultural</p> <p>Gardens</p> <p>Vineyards</p> <p>Orchards</p>	<p>Wide</p> <p>Aphids</p> <p>Psyllids</p> <p>Coccids</p> <p><i>Metcalfa pruinosa</i></p> <p>Pollen</p>	<p><i>Dinocampus coccinellae</i></p> <p><i>Phalacrotophora fasciata</i></p>
The Netherlands	<p>Intentional introduction</p> <p>Presumed secondary spread from Belgium</p>	Found on outdoor crops in 1996 but first specimen (a pupa) in the wild recorded on 2002	Expanded from the first record in 2002 to spread rapidly from 2004 onwards, from south to north. By 2006 establishment across the entire country	<p>Agricultural (dominant in corn)</p> <p>Urban</p> <p>Predominantly on trees and shrubs but also on herbs in urban and anthropogenic habitats, less abundant in (semi-) natural areas</p>	<p>Wide</p> <p>Including cannibalism (observation of specimens egg-eating while other specimen was laying eggs)</p>	<p><i>Araneus diadematus</i> (autumn),</p> <p><i>Coccipolipus hippodamiae</i> (winter),</p> <p><i>Dinocampus coccinellae</i> (summer, winter)</p> <p><i>Hesperomyces virescens</i> (summer, winter)</p> <p><i>Parasitylenchus bifurcatus</i> (winter)</p> <p><i>Phalacrotophora fasciata</i> and</p> <p><i>Phalacrotophora</i> sp. (spring, summer)</p>

Norway	<p>Unintentional introduction</p> <p>Oslo in 2006, arrived as a stowaway on horticultural plants, <i>Thuja</i> sp. (Cupressaceae) imported from the Netherlands</p>	Late 2007 and throughout 2008 several adults were found indoors in the urban and suburban areas of Oslo	Limited to urban and suburban regions, and not recorded from commercial crops or in natural habitats	<p>Urban Gardens Parks</p> <p>Most records from the wider Oslo area. No reports from agricultural or horticultural crops</p>	Not measured	<i>Dinocampus coccinellae</i>
Slovakia	<p>Unintentional introduction.</p> <p>Presumed secondary spread from Austria, Poland and the Czech Republic</p>	First recorded in 2008	Expanded from the first record in 2008 across the entire country by 2009 at a rate of about 200km per year. Local topography played a role in the spread	<p>Urban Orchards Deciduous and lowland pine forests Vineyards Agricultural</p>	<p>Wide</p> <p>Hemiptera Larvae of Lepidoptera Dying flightless adults of Diptera</p>	<i>Chrysoperla carnea</i> <i>Passer montanus</i>
Switzerland	<p>Unintentional introduction</p> <p>Presumed secondary spread from Germany and France</p>	First recorded in 2004, with very large populations found in 2006	Expanded from the first record in 2004 to rapidly invade all areas of low and middle altitudes of the country by 2006	<p>Urban Woodland</p> <p>(particularly on broadleaved trees in woodlands and urban areas, but much less frequent on conifers, meadows or in agricultural habitats)</p>	Not measured	Not recorded

UK	<p>Unintentional introduction</p> <p>Presumed secondary spread from mainland Europe particularly Belgium and France</p>	First recorded in 2003 and as established in 2005	Expanded from first record in 2003 at more than 100km per year in the UK, both via flight and anthropogenic transport	<p>Urban Parks and gardens Woodlands (mainly deciduous) Grasslands Agricultural</p>	<p>Wide</p> <p>Hemiptera Lepidoptera Coleoptera</p>	<p><i>Dinocampus coccinellae</i> <i>Hesperomyces virescens</i> <i>Homalotylus flaminus</i> <i>Medina separata</i> <i>Oomyzus scaposus</i> <i>Phalacrotophora berlinensis</i> <i>Phalacrotophora fasciata</i> (Lower susceptibility of <i>H. axyridis</i> compared to other species evident from laboratory and field studies)</p>
South Africa	<p>Intentional introduction c. 1980: no establishment.</p> <p>Present invasion likely from unintentional introduction (unknown whether multiple).</p> <p>Eastern North America (genetic analysis of 4 South African populations)</p>	First records 2001 (museum records). Established by 2006	Spread rapidly, established largely in cooler and more mesic regions. Initial spread of up to 500 km per year likely human-assisted.	<p>Gardens (urban and rural Orchards (deciduous and subtropical fruit) Vineyards Field crops Pine plantations Indoors Untransformed fynbos, grassland, moist savanna</p> <p>Frequently on oak trees, ornamental conifers, garden roses</p>	<p>Wide</p> <p>Field observations include: Aphids Coccinellids</p>	<p><i>Dinocampus coccinellae</i> <i>Hesperomyces virescens</i></p>

Table 2: Life history traits of *Harmonia axyridis* documented from countries around the world. It is important to note that there are various measurements for some of the life history traits and so direct comparison between countries is not always possible. Standard approaches to capturing this information would be useful to enable quantitative assessments on a global scale. Body size has been measured either as length and/or width (mm) or volume (μ l) or mass (mg). Voltinism is given as number of generations per year. Fecundity has been measured as clutch size (number of eggs per cluster) and lifetime (number of eggs per lifetime). In some cases the maximum (max.) number of eggs within a cluster is given. Hatching success is given as % . Temperature tolerance is measured in various ways: development threshold ($^{\circ}$ C), accumulated temperature (degree-days), lower lethal temperature ($^{\circ}$ C), lower lethal time (weeks), development time (days), maximum and minimum critical temperature (CTmax and CTmin respectively) ($^{\circ}$ C). Unless otherwise stated mean \pm standard error are given for all traits.

	Body size	Colour form(s)	Voltinism	Fecundity	Hatching success	Temperature tolerance and development
China	Egg: length 1.0–1.5 mm 4th larval instar: length 10–15 mm Pupa: length 7 mm, width 4 mm Adult: length 5.4–8.0 mm, width 3.8–5.2 mm	> 200 colour forms including: <i>succinea</i> <i>spectabilis</i> <i>conspicua</i> <i>axyridis</i>	2–8	Clutch size: 29.95 \pm 0.46 (mean \pm SD) Lifetime: 1063.02 \pm 62.15 (mean \pm SD)	84.85 \pm 0.46%	Development threshold: 10.5 $^{\circ}$ C; cannot develop above 34 $^{\circ}$ C Accumulated temperature: 231.3 degree-days
Japan	Egg: volume 0.25 \pm 0.04 μ l (mean \pm SD) Adult female: size 6.73 \pm 0.53 mm (n=1106) (mean \pm SD) Adult male: size 6.16 \pm 0.43 mm (n=813)	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i> <i>axyridis</i>	2–3	Clutch size: 39.11 \pm 1.47 Lifetime: 455.4 \pm 163.46	65.84 \pm 4.34% (including individuals with male-killing endosymbiont)	Not measured
Siberia and the Far East	4th larval instar: length 8.2–11.2 mm Adult length: 4.9–8.2 mm	In population living west to Baikal lake almost all specimens are <i>axyridis</i> ; other forms are rare.	1–3	Clutch size: 11–52	Not measured	Mating at 16–18 $^{\circ}$ C. Optimal temperature for egg development: 17–26 $^{\circ}$ C; for larval development: 26–

		<p>In populations living east to Baikal lake 3 colour forms are common: <i>succinea</i> <i>conspicua</i> <i>spectabilis</i></p> <p>Other forms are rare.</p>				<p>30°C; for pupal development: 22–30°C.</p> <p>All stages can survive at temperature 40°C for several hours.</p>
European Russia	Not measured	<p>4 colour forms: <i>succinea</i> <i>Spectabilis</i> <i>Conspicua</i> <i>axyridis</i></p> <p>Populations in the vicinity of Sochi and Abkhazia are all <i>succinea</i></p>	At least 2	Not measured	Not measured	Not measured
North America (USA)	<p>Adult: length 4.8–7.47 mm, width 3.9–5.89 mm</p> <p>Egg: 0.25±0.0mg Adult male: 31.6±0.9mg Adult female: 53.6±2.2mg</p>	Mainly <i>succinea</i> ; melanic <i>H. axyridis</i> uncommon	2 plus a partial third	<p>Clutch size: single eggs to more than 20 per cluster (> 60% of 3,149 females observed had egg cluster with > than 20 eggs)</p> <p>37.3±4.3 eggs per day</p>	<p>93.6% non-cannibalising larvae and 94.1% of larvae that had engaged in sibling cannibalism survived to adulthood (laboratory).</p> <p>In wild-caught individuals 97±4% (n = 22 egg clutches)</p>	Development from egg to adult requires 231.3 degree days above a lower developmental threshold of 10.5°C.

South America (Chile)	<p>Larvae: length L1: 1.96±0.04 mm; L2: 4.6±0.08 mm; L3: 6.3±0.11 mm; L4: 8.3±0.17 mm</p> <p>Adults female: length 7.2±0.04 mm (range: 6–8.3mm); width 5.6±0.06 mm (range: 4.7 – 6.2 mm)</p> <p>Adult male: length 6.7±0.04mm (range: 5.5 – 7.7mm), width 5.3±0.05mm (range: 4.1 – 5.8mm)</p>	<i>succinea</i> with some variation in spot number and red intensity	2–3	Clutch size: 14.1±1.2 eggs per day at 20°C (fed with <i>A. pisum</i>)	Not measured	No study of tolerance but in relation to preference <i>H. axyridis</i> preferred temperatures between 15 and 20°C, with thermal between 8.6–27.2°C. No eggs hatched at 33°C
Belgium	Adult: length 5–9 mm (median =7)	<i>succinea spectabilis</i> <i>conspicua</i>	2 in (semi-) natural areas, one in field crops (second develops late in the season in other habitats)	Clutch size: 12-48 eggs (depending on food source and colour morph)	15–64% (depending on food source and colour morph)	Lower lethal temperature for outdoor overwintering: from –17.1 to –16.3°C. Lower lethal time of field populations at 0 and –5°C: 18–24 and 17–22 weeks respectively
Czech Republic	<p>Adult female: (reproduction period) 44.3±9.3 mg; (migrating) 36.1±5.7 mg</p> <p>Adult male: (reproduction period) 29.8±6.0 mg;</p>	<p>88% <i>succinea</i></p> <p>9% <i>spectabilis</i></p> <p>3% <i>conspicua</i></p> <p><i>axyridis</i> - rare</p> <p><i>equicolor</i> - extremely rare</p>	2–3	<p>Clutch size: 21 eggs per day (max. 86)</p> <p>Lifetime: 1,174 eggs (max. 2,497)</p>	82%	High egg and larval mortality at 33°C; LDT 11°C, SET 175 DD

	(migrating) males: 30.0±5.3 mg					
Denmark	Not measured	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i> <i>axyridis</i>	2 plus occasional partial third	Not measured	Not measured	Not measured
France	Adult female: length 7.19±0.18 mm; width 5.71±0.11 mm; mass 51.5±5.5 mg Adult male: length 6.55±0.11 mm; width 5.40±0.09 mm; mass 40.2±3.9 mg	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i>	2–3	Clutch size: 33.5±5 eggs (at 24°C, 60% RH, L:D 14:10; fed ad libitum with irradiated eggs of <i>E. kuehniella</i>)	79.5±12.8%	Not measured
Germany	Adult female: length 5.8 ± 0.4 mm; width 4.2 ± 0.3 mm Adult male: length 5.4 ± 0.4 mm; width 4.0 ± 0.3 mm	92 % <i>succinea</i> 7 % <i>spectabilis</i> 1 % <i>conspicua</i>	2	Not measured	Not measured	Not measured
Italy	Adult female: length 7.14±0.10 mm; width 5.54±0.07 mm (n=24) Adult male: length 6.48±0.08 mm; width 5.19±0.07 mm (n=20)	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i>	At least 2, potentially 4	Clutch size: 15.4±3.9 eggs Lifetime: 560.5 - 783.8 eggs 228.3±30.7 eggs laid in 10 days	58–100% (n=988) reared on <i>Aphis fabae</i> ; 53% reared on <i>M.</i> <i>persicae</i>	At 35°C: no egg hatching. At 30°C: 12.8±0.19 d preimaginal development time; 1.15±1.01 egg fertility (%) At 25°C: 13.84±0.09 d

						preimaginal development time; 21.63±5.12 egg fertility (%)
The Netherlands	Adult: length 5.4–7.8 mm (median = 6.8); H/L elytra 0.44–0.48 mm Adult female: 28.2–37.9 mg Adult male: 24.1–32.8 mg	81% <i>succinea</i> 16% <i>spectabilis</i> 4% <i>conspicua</i> <i>axyridis</i> (6 records) <i>equicolor</i> (6 records) very sporadically (less than 1:1000) <i>intermedia</i>	2–3	Clutch size: 20-30 eggs	Not measured	Nightly flight activity commences at 10.5°C
Norway	Not measured	<i>succinea</i>	Not measured, suggests 1-2	Not measured	Not measured	Not measured
Slovakia	Adult female: length 6.0–7.5 mm (median = 6.8 mm); width 4.7–5.9 mm (median 5.4 mm, n=38) Adult male: length 5.8–7.0 mm (median 6.5 mm); width 4.1–5.9 mm (median 5.1mm, n=34)	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i> <i>axyridis</i> <i>equicolor</i>	2	Not measured	Not measured	Not measured
Switzerland	Not measured	<i>succinea</i> <i>spectabilis</i> <i>conspicua</i>	2 (in Northern Switzerland, measured by regular sampling programmes)	Not measured	Not measured	Not measured

UK	Adult female: length 5.8–8.0 mm (median = 7.0); width 4.8–6.5 mm (median = 5.4) (n = 37) Adult male: length 5.4–7.5 mm (median = 6.7); width 4.6–5.9 mm (median = 5.1) (n = 39)	<i>succinea spectabilis conspicua</i>	2 (but with occasional partial third generation)	Not measured	Not measured	Not measured but UK distribution currently limited within Scotland and climatic factors (particular temperature) assumed to be important
South Africa	Adult female: length 6.9±0.1 mm (n=10); mass 35.3±1.0 mg (n=34) Adult male: length 6.3±0.1 mm (n=10); mass 29.1±0.9 mg (n=31)	<i>succinea</i> with variation in spot number and background colour	Exact number unknown but at least 2	Over 70 days: 245 ± 40 eggs Rearing conditions: 25/18°C (18h day/ 6h night cycle; 14hL/10hD photoperiod)	53±6.5% (n=650) Rearing conditions: 25/18 °C (18h day/ 6 night cycle; 14hL/10hD photoperiod)	Development time: 23.1±0.1 days (n=267); CTmax = 44.1±0.1°C (n=34); CTmin = 0.9±0.2°C (n=31), start temp 25°C, rate of change: 0.05°C/min; Rearing conditions: 25/18°C (18h day/ 6h night cycle; 14hL/10hD photoperiod)

Legend

Harmonia_Europe

No record of invasion

- 1991
- 1994
- 1998
- 1999
- 2001
- 2002
- 2003
- 2004
- 2005
- 2006
- 2007
- 2008
- 2009
- 2010
- 2011
- 2012
- 2013
- 2014



