

Cardoso DM & Rowlinson P (2010) Spectral upper bounds for the order of a  $k$ -regular induced subgraph, *Linear Algebra and Its Applications*, 433 (5), pp. 1031-1037.

**This is the peer reviewed version of this article**

*NOTICE: this is the author's version of a work that was accepted for publication in Linear Algebra and Its Applications. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Linear Algebra and Its Applications, [VOL 433, ISS 5 (2010)] DOI: <http://dx.doi.org/10.1016/j.laa.2010.04.029>*

# Spectral upper bounds for the order of a $k$ -regular induced subgraph

Domingos M. Cardoso<sup>1</sup>  
Departamento de Matemática  
Universidade de Aveiro  
3810-193 Aveiro  
Portugal  
dcardoso@ua.pt

Peter Rowlinson<sup>2</sup>  
Department of Computing Science and Mathematics  
University of Stirling  
Stirling FK9 4LA  
Scotland  
p.rowlinson@stirling.ac.uk

**Abstract.** Let  $G$  be a simple graph with least eigenvalue  $\lambda$ , and let  $S$  be a set of vertices in  $G$  which induce a subgraph with mean degree  $k$ . We use a quadratic programming technique in conjunction with the main angles of  $G$  to establish an upper bound of the form  $|S| \leq \inf\{(k+t)q_G(t): t > -\lambda\}$ , where  $q_G$  is a rational function determined by the spectra of  $G$  and its complement. In the case  $k = 0$  we obtain improved bounds for the independence number of various benchmark graphs.

Keywords: graph, main eigenvalue, independence number, clique number

AMS Classification: 05C50

---

<sup>1</sup>Research supported by the Centre for Research on Optimization and Control from the Fundação para a Ciência e Tecnologia, cofinanced by the European Community Fund FEDER/POCI 2010

<sup>2</sup>Corresponding author

# 1 Introduction

Let  $G$  be a simple graph of order  $n$  with  $(0, 1)$ -adjacency matrix  $A$  and characteristic polynomial  $P_G(x) = \det(xI - A)$ . The  $i$ -th largest eigenvalue of  $A$  is denoted by  $\lambda_i(G)$ , and we write  $\lambda_i = \lambda_i(G)$ ,  $\bar{\lambda}_i = \lambda_i(\bar{G})$ , where  $\bar{G}$  denotes the complement of  $G$ .

Let  $S$  be a set of vertices in  $G$  which induce a subgraph with mean degree  $k$ . We use a quadratic programming technique [2, 3] in conjunction with the main angles of  $G$  [8, Section 4.5] to prove that

$$|S| \leq \inf\{h_k^G(t) : t > -\lambda_n(G)\}, \quad (1)$$

where

$$h_k^G(t) = (k + t) \left\{ 1 - \frac{P_{\bar{G}}(t - 1)}{(-1)^n P_G(-t)} \right\}.$$

Thus if we write  $H_G(t)$  for the walk-generating function of  $G$  (see [4] or [14]) then

$$h_k^G(t) = \left(1 + \frac{k}{t}\right) H_G\left(-\frac{1}{t}\right).$$

We give computational results which demonstrate that the bound (1) is superior to previous bounds. We make use of the functions  $f_{k,t}^A : \mathbb{R}^n \rightarrow \mathbb{R}$  defined for  $t > 0$  by

$$f_{k,t}^A(\mathbf{x}) = 2\mathbf{j}^\top \mathbf{x} - \frac{1}{k+t} \mathbf{x}^\top (A + tI) \mathbf{x},$$

where  $\mathbf{j}$  denotes the all-1 vector in  $\mathbb{R}^n$ . These functions were constructed in [3] to determine upper bounds for the order of a  $k$ -regular induced subgraph in terms of eigenvalues. The problem of finding the largest order of such a subgraph is NP-complete [2, Section 2], whereas spectral upper bounds can be computed in polynomial time. We too state our results in terms of  $k$ -regular induced subgraphs, but they apply equally to induced graphs with mean degree  $k$  (for example, induced unicyclic graphs, with mean degree 2). When  $k = 0$  we obtain an upper bound for the independence number  $\alpha(G)$ ; a spectral lower bound for  $\alpha(G)$ , in terms of  $n$ ,  $\bar{\lambda}_n$  and the mean degree of  $G$ , is derived in [13].

We shall first summarize the basic argument in [3]. Recall that the eigenvalue  $\lambda$  of  $G$  is a *main* eigenvalue if the eigenspace  $\mathcal{E}_A(\lambda)$  is not orthogonal to  $\mathbf{j}$ . In particular,  $\lambda_1$  is a main eigenvalue because the Perron-Frobenius theory ensures that  $A$  has a corresponding eigenvector whose entries are all non-negative.

If  $t \geq -\lambda_n$  then  $f_{k,t}^A$  is concave, that is,

$$f_{k,t}^A(\theta \mathbf{x} + (1 - \theta) \mathbf{y}) \geq \theta f_{k,t}^A(\mathbf{x}) + (1 - \theta) f_{k,t}^A(\mathbf{y})$$

whenever  $0 \leq \theta \leq 1$ . (To see this, express  $\mathbf{x}, \mathbf{y}$  as sums of eigenvectors of  $A$ ; alternatively, note that the Hessian matrix of  $f_{k,t}^A(t)$  is  $\frac{-2}{k+t}(A + tI)$ , which is negative semi-definite when  $t \geq -\lambda_n$ .) Accordingly,  $f_{k,t}^A$  has a global maximum at  $\mathbf{x}^*$  if and only if  $\nabla f_{k,t}^A(\mathbf{x}^*) = \mathbf{0}$ , that is,

$$\mathbf{j} - \frac{1}{k+t}(A + tI)\mathbf{x}^* = \mathbf{0}.$$

Then  $f_{k,t}^A(\mathbf{x}^*) = \mathbf{j}^\top \mathbf{x}^*$ . If  $\mathbf{x}_S$  is the characteristic vector of  $S$  then  $\mathbf{x}_S^\top A \mathbf{x}_S = k|S|$  and so  $|S| = f_{k,t}^A(\mathbf{x}_S) \leq f_{k,t}^A(\mathbf{x}^*)$ . Note that  $f_{k,t}^A(\mathbf{x}_S) = f_{k,t}^A(\mathbf{x}^*)$  if and only if  $(A + tI)\mathbf{x}_S = (k + t)\mathbf{j}$ , equivalently  $S$  is a  $(k, k + t)$ -regular set (that is,  $S$  induces a  $k$ -regular subgraph, while each vertex outside  $S$  is adjacent to  $k + t$  vertices inside  $S$ ).

Let  $J$  denote an all-1 matrix. If  $G \neq \overline{K}_n$  and  $\bar{\lambda}$  is a main eigenvalue of  $\overline{G}$  such that  $\bar{\lambda} \geq -\lambda_n - 1$ , then we may take  $t = \bar{\lambda} + 1$  and

$$\mathbf{x}^* = \frac{k + t}{\mathbf{j}^\top \mathbf{u}} \mathbf{u},$$

where  $\mathbf{u}$  is an eigenvector of  $J - I - A$  corresponding to  $\bar{\lambda}$  such that  $\mathbf{j}^\top \mathbf{u} \neq 0$ . (Note that then  $(A + tI)\mathbf{u} = J\mathbf{u} = \mathbf{j} \mathbf{j}^\top \mathbf{u}$ .) The Courant - Weyl inequalities imply that

$$\lambda_2(\overline{G}) + \lambda_n(G) \leq \lambda_2(K_n) = -1 = \lambda_n(K_n) \leq \lambda_1(\overline{G}) + \lambda_n(G).$$

Thus we may always take  $\bar{\lambda} = \bar{\lambda}_1$ , and the remaining possibility is  $\bar{\lambda} = -\lambda_n - 1$  when  $-\lambda_n - 1$  is a main eigenvalue of  $\overline{G}$ . Since  $f_{k,t}^A(\mathbf{x}^*) = \bar{\lambda} + k + 1$ , we obtain:

**Theorem 1.1** (cf. [3, Section 3]). *Let  $G$  be a graph of order  $n$ , and let  $S$  be a set of vertices which induces a  $k$ -regular subgraph of  $G$  ( $0 \leq k \leq n - 1$ ). Then*

$$|S| \leq \bar{\lambda}_1 + k + 1. \quad (2)$$

*If  $-\lambda_n - 1$  is a main eigenvalue of  $\overline{G}$  then*

$$|S| \leq -\lambda_n + k \quad (3)$$

Two remarks are in order:

(i) When  $k = 0$  we obtain from (2) the well-known upper bound  $\bar{\lambda}_1 + 1$  for the independence number  $\alpha(G)$ . This bound is attained when, for example,  $G$  is a complete graph or a complete bipartite graph.

(ii) If  $-\lambda_n - 1$  is a main eigenvalue of  $\overline{G}$  then  $\lambda_n$  is a non-main eigenvalue of  $G$ , and  $-\lambda_n - 1$  is a multiple eigenvalue of  $\overline{G}$ . This is a particular case of the following observation, essentially Theorem 2.12 of [5], for which we give a direct proof.

**Proposition 1.2.** *If  $\lambda$  is an eigenvalue of  $G$  such that  $-\lambda - 1$  is a main eigenvalue of  $\overline{G}$ , then  $\lambda$  is a non-main eigenvalue of  $G$ ; moreover, if  $\lambda$  has multiplicity  $d$  as an eigenvalue of  $G$  then  $-\lambda - 1$  has multiplicity  $d + 1$  as an eigenvalue of  $\overline{G}$ .*

**Proof.** Let  $(J - I - A)\mathbf{y} = (-\lambda - 1)\mathbf{y}$ , where  $\mathbf{j}^\top \mathbf{y} \neq 0$ . Let  $\mathbf{x} \in \mathcal{E}_A(\lambda)$ . Then  $(J - A)\mathbf{y} = -\lambda\mathbf{y}$  and  $\mathbf{x}^\top A = \lambda\mathbf{x}^\top$ . Hence  $\mathbf{x}^\top (J - A)\mathbf{y} = -\lambda\mathbf{x}^\top \mathbf{y}$  and  $\mathbf{x}^\top A\mathbf{y} = \lambda\mathbf{x}^\top \mathbf{y}$ . Adding, we have  $\mathbf{x}^\top J\mathbf{y} = 0$ , that is,  $\mathbf{x}^\top \mathbf{j} \mathbf{j}^\top \mathbf{y} = 0$ . Hence  $\mathbf{x}^\top \mathbf{j} = 0$  for all  $\mathbf{x} \in \mathcal{E}_A(\lambda)$ ; in other words,  $\lambda$  is a non-main eigenvalue of  $G$ . Now  $\mathcal{E}_{J-I-A}(-\lambda - 1) \cap \mathbf{j}^\perp = \mathcal{E}_A(\lambda)$ , and the second assertion follows.  $\square$

## 2 Further bounds

Here we introduce improved bounds by involving the main angles of  $G$ . We write  $\mu_1, \dots, \mu_s$  for the main eigenvalues of  $G$  in decreasing order. Then  $\mathbf{j}$  is expressible as

$$\mathbf{j} = \mathbf{u}_1 + \dots + \mathbf{u}_s \quad (\mathbf{u}_i \in \mathcal{E}_A(\mu_i)).$$

Thus  $\mu_1 = \lambda_1$ , and the non-zero main angles of  $G$  are  $\beta_1, \dots, \beta_s$  where  $\sqrt{n}\beta_i = \|\mathbf{u}_i\|$  ( $i = 1, \dots, s$ ).

**Theorem 2.1.** *Let  $G$  be a graph of order  $n$ , and let  $S$  be a set of vertices which induces a  $k$ -regular subgraph of  $G$  ( $0 \leq k \leq n-1$ ). If  $t > -\lambda_n$  then*

$$|S| \leq n \sum_{i=1}^s \frac{t+k}{t+\mu_i} \beta_i^2; \quad (4)$$

equivalently,

$$|S| \leq h_k^G(t), \quad (5)$$

where

$$h_k^G(t) = (k+t) \left\{ 1 - \frac{P_G(t-1)}{(-1)^n P_G(-t)} \right\}. \quad (6)$$

**Proof.** If  $t > -\lambda_1$  then the function  $f_{k,t}^A$  is concave and attains its maximum at

$$\mathbf{x}^* = \sum_{i=1}^s \frac{k+t}{\mu_i+t} \mathbf{u}_i.$$

Hence

$$|S| = f_{k,t}^A(\mathbf{x}_S) \leq \mathbf{j}^\top \mathbf{x}^* = n \sum_{i=1}^s \frac{t+k}{t+\mu_i} \beta_i^2.$$

The equivalent bound (5) is obtained by setting  $x = t-1$  in the formula [7, p.90]

$$P_G(x) = (-1)^n P_G(-x-1) \left\{ 1 - \sum_{i=1}^s \frac{n\beta_i^2}{x+1+\mu_i} \right\}. \quad (7)$$

□

When  $\lambda_n$  is a main eigenvalue of  $G$ , the graph of  $y = h_k^G(t)$  has  $t = -\lambda_n$  as an asymptote, and so we state our main result as follows. Here the second assertion follows from our remarks in Section 1.

**Corollary 2.2.** *If  $S$  induces a  $k$ -regular subgraph of  $G$  then*

$$|S| \leq \inf \{ h_k^G(t) : t > -\lambda_n(G) \}.$$

We have  $|S| = h_k^G(t_0)$  if and only if  $S$  is a  $(k, k+t_0)$ -regular set.

When  $\lambda_n$  is a non-main eigenvalue of  $G$ , we have  $G \neq \overline{K_n}$  and we may take  $t = -\lambda_n$  to obtain the following reformulation of [3, Theorem 3.4]:

**Theorem 2.3.** *Let  $G$  be a graph of order  $n$ , and let  $S$  be a set of vertices which induces a  $k$ -regular subgraph of  $G$  ( $0 \leq k \leq n-1$ ). If  $\lambda_n$  is a non-main eigenvalue of  $G$  then*

$$|S| \leq n \sum_{i=1}^s \frac{-\lambda_n + k}{-\lambda_n + \mu_i} \beta_i^2. \quad (8)$$

In Equation (5) we should cancel factors common to  $P_{\overline{G}}(t-1)$  and  $P_G(-t)$ . To this end, let  $M_G(x) = (x - \mu_1) \cdots (x - \mu_s)$ , and  $M_{\overline{G}}(x) = (x - \overline{\mu}_1) \cdots (x - \overline{\mu}_s)$ , where  $\overline{\mu}_1, \dots, \overline{\mu}_s$  are the main eigenvalues of  $\overline{G}$  (cf. [14]). By Proposition 1.2 applied to  $G$  and  $\overline{G}$ , or by Equation (8) of [14], we have

$$\frac{P_{\overline{G}}(t-1)}{(-1)^n P_G(-t)} = \frac{M_{\overline{G}}(t-1)}{(-1)^s M_G(-t)}; \quad (9)$$

moreover,  $M_{\overline{G}}(t-1)$  and  $M_G(-t)$  have no common factors. Thus  $h_k^G(t) = k + t$  if and only if  $t - 1$  is a main eigenvalue of  $\overline{G}$ . In particular, we may take  $t = 1 + \overline{\lambda}_1$  to obtain the bound (1). In the case that  $-1 - \lambda_n$  is a main eigenvalue of  $\overline{G}$ , we take  $t = -\lambda_n$  in (4) and (6) to deduce:

**Proposition 2.4.** *When  $-\lambda_n - 1$  is a main eigenvalue of  $\overline{G}$ , the upper bounds (3) and (8) coincide.*

To discuss the improvements on (2) afforded by Corollary 2.2, we write  $h_k(t)$  for  $h_k^G(t)$ . If either

$$-\lambda_n < \overline{\lambda}_1 + 1 \text{ and } h'_k(1 + \overline{\lambda}_1) \neq 0$$

or

$$-\lambda_n = \overline{\lambda}_1 + 1 \text{ and } h'_k(1 + \overline{\lambda}_1) < 0,$$

then an improvement on (1) is assured in a neighbourhood of  $1 + \overline{\lambda}_1$ . We have

$$h'_k(1 + \overline{\lambda}_1) = 1 - (k + 1 + \overline{\lambda}_1)(-1)^s \left\{ \frac{M'_{\overline{G}}(\overline{\lambda}_1)}{M_G(-1 - \overline{\lambda}_1)} \right\},$$

but it is more revealing to inspect two small examples.

**Example 2.5.** Let  $G = 3K_1 \dot{\cup} K_2 \dot{\cup} K_3$ . Then  $P_G(x) = (x-2)(x-1)x^3(x+1)^3$ . Using the computer package GRAPH, we find that  $P_{\overline{G}}(x) = (x^3 - 2x^2 - 21x - 24)x^3(x+1)^2$ ; moreover,  $0$  ( $= -\lambda_8 - 1$ ) is not a main eigenvalue of  $\overline{G}$ . We have  $\overline{\lambda}_1 \approx 6.0930$ , and so the bound (1) yields  $|S| \leq 7$  when  $k = 0$ . Here  $\mu_s = 0 = k$  and  $y = h_0(t)$  does not have  $t = 0$  as an asymptote. We have

$$h_0(t) = \frac{2(2t+3)(2t+1)}{(t+1)(t+2)},$$

a function which increases monotonically on  $[-\lambda_8, \infty)$ . Whenever  $h_k(t)$  has this property, and  $\mu_s > \lambda_n$ , the best bound arises when  $t = -\lambda_n$ , giving a formula that coincides with (8). In this example, we obtain  $|S| \leq 5$  (a sharp upper bound since  $\alpha(G) = 5$ ).  $\square$

**Example 2.6.** Let  $G$  be the graph on 6 vertices numbered 50 in the table [6], where characteristic polynomials are listed and main angles are identified; the complement of  $G$  is numbered 100 in [6]. We have  $s = 4$ ,  $\mu_4 = \lambda_6 \approx -2.508$  and  $\overline{\lambda}_1 \approx 2.228$ . We take  $k = 0$  again, and then the upper bound (1) for  $|S|$  is 3.228. In this case  $y = h_0(t)$  has  $t = -\lambda_6$  as an asymptote. Explicitly,

$$h_0(t) = \frac{2t(3t^3 - 9t^2 + t + 7)}{t^4 - 9t^2 + 4t + 7}.$$

This function has a unique local minimum on  $(-\lambda_6, \infty)$ . Using Mathematica, we find that this minimum is 3.132 at  $t = 2.834$  (to three places of decimals).

This new upper bound is smaller, but of course both bounds yield  $|S| \leq 3$  (a sharp inequality since  $\alpha(G) = 3$ ).  $\square$

These examples are provided to illustrate differences in the behaviour of  $h_k$ . To demonstrate the superiority of the bound in Corollary 2.2, we should consider larger graphs, and this we do in the next section. Here we first discuss properties of  $h_k$  in the general case.

**Proposition 2.7** *The function  $h_k(t)$  has at most one local minimum in  $(-\mu_s, \infty)$ .*

**Proof.** The result is immediate if  $s = 1$  (that is, if  $G$  is regular), since then  $h_k(t)$  is monotonic. Accordingly we suppose that  $s > 1$ . We have

$$h_k(t) = n - \sum_{i=1}^s \frac{n(\mu_i - k)\beta_i^2}{t + \mu_i}. \quad (10)$$

Suppose first that  $k$  is not a main eigenvalue of  $G$ , so that the graph  $\mathcal{G}$  of  $y = h_k(t)$  has asymptotes  $t = -\mu_i$  ( $i = 1, \dots, s$ ). Note also that  $h_k(t) \rightarrow n$  as  $t \rightarrow \infty$  and as  $t \rightarrow -\infty$ .

If  $\mu_s < k$  then the line  $y = d$  cuts  $\mathcal{G}$  in (at least)  $s - 1$  points of  $(-\infty, -\mu_s)$  when  $d > n$ , and (at least)  $s - 2$  points of  $(-\infty, -\mu_s)$  when  $d < n$ . If  $\mu_s > k$  then the line  $y = d$  cuts  $\mathcal{G}$  in (at least)  $s$  points of  $(-\infty, -\mu_s)$  when  $d > n$ , and (at least)  $s - 1$  points of  $(-\infty, -\mu_s)$  when  $d < n$ .

Now suppose that  $h_k(t)$  has a local minimum at  $t_0 \in (-\mu_s, \infty)$ . Then  $h'_k(t) \geq 0$  for all  $t \geq t_0$ , for otherwise  $h_k(t)$  has a local maximum at some point  $t_1 \in (t_0, \infty)$ . If  $h_k(t_1) > n$  then for some  $d > n$ , the line  $y = d$  cuts  $\mathcal{G}$  in (at least) 3 points in  $(-\mu_s, \infty)$ . If  $h_k(t_1) \leq n$  then for some  $d < n$ , the line  $y = d$  cuts  $\mathcal{G}$  in (at least) 4 points in  $(-\mu_s, \infty)$ . In any case, the function  $h_k(t) - d$  has more than  $s$  zeros in  $\mathbb{R}$ . This is a contradiction because  $h_k(t) - d$  ( $d \neq n$ ) has the form  $p(t)/q(t)$ , where  $p(t), q(t)$  are polynomials of degree  $s$ .

If  $k$  is a main eigenvalue of  $G$ , then the same arguments apply to a graph with  $s - 1$  vertical asymptotes.

It follows that  $h_k(t)$  has no more than one local minimum in  $(-\mu_s, \infty)$ .  $\square$

**Corollary 2.8** *For a non-regular graph  $G$ , we have:*

- (i) *if  $\mu_s < 0$  then  $h_0^G(t)$  has a unique local minimum in  $(-\mu_s, \infty)$ ,*
- (ii) *if  $\mu_s = 0$  then  $h_0^G(t)$  is increasing on  $(-\mu_{s-1}, \infty)$ ,*
- (iii) *if  $\mu_s > 0$  then  $h_0^G(t)$  is increasing on  $(-\mu_s, \infty)$ .*

**Proof.** We have  $h_0(1 + \bar{\lambda}_1) = 1 + \bar{\lambda}_1 < n$  and  $1 + \bar{\lambda}_1 \in (-\mu_s, \infty)$ . Thus if  $\mu_s < 0$  then  $h_0(t)$  has a local minimum on  $(-\mu_s, \infty)$ , and this minimum is unique by Proposition 2.7. If  $\mu_s = 0$  then from (10) we see that  $h'_0(t) > 0$  for all  $t \in (-\mu_{s-1}, \infty)$ , and if  $\mu_s > 0$  then  $h'_0(t) > 0$  for all  $t \in (-\mu_s, \infty)$ .  $\square$

We conclude this section by deriving sharp upper bounds in two special cases. First, if  $G$  is  $r$ -regular, we may apply Theorem 2.3 to obtain

$$|S| \leq \frac{n(k - \lambda_n)}{r - \lambda_n}.$$

This bound, known as the Hoffman bound when  $k = 0$ , coincides with that obtained from interlacing (cf. [10, Lemma 9.6.2]). It is attained in some of

the regular graphs  $\overline{G}$  discussed in Section 3. Other generalizations of the Hoffman bound may be found in [1, Theorem 7] and [9, Corollary 3.2].

Secondly, consider a connected harmonic graph  $G$ , that is, a connected graph  $G$  for which  $A\mathbf{d} = \mu_1\mathbf{d}$ , where  $\mathbf{d}$  is the vector whose entries are the vertex degrees. We show that if  $G$  has  $e$  edges then

$$\alpha(G) \leq n - \frac{e}{\mu_1}. \quad (11)$$

The main eigenvalues of  $G$  are  $\mu_1$  and 0 [14, Proposition 3.3], and so

$$\alpha(G) \leq h_0(-\lambda_n) = n \left\{ 1 - \frac{\mu_1}{\mu_1 - \lambda_n} \beta_1^2 \right\} \leq n(1 - \tfrac{1}{2}\beta_1^2).$$

To determine  $\beta_1$  when  $G$  is connected, note that

$$\mathbf{u}_1 = \frac{1}{\|\mathbf{d}\|^2} (\mathbf{d}^\top \mathbf{j}) \mathbf{d}, \text{ whence } n\beta_1^2 = \frac{4e^2}{\|\mathbf{d}\|^2}.$$

Since  $\mathbf{d} - \mu_1\mathbf{j} \in \mathcal{E}_A(0) \subseteq \mathcal{E}_A(\mu_1)^\perp = \mathbf{d}^\perp$ , we have  $\|\mathbf{d}\|^2 = 2e\mu_1$ , and so  $n(1 - \frac{1}{2}\beta_1^2) = n - \frac{e}{\mu_1}$ , proving (11). We note that this bound is attained in all Grünwald trees [11, 14]: for such a tree  $T$  we have  $\lambda_n = -\mu_1$ ,  $e = n - 1 = \mu_1(\mu_1^2 - \mu_1 + 1)$  and  $\alpha(T) = (\mu_1 - 1)(\mu_1^2 - \mu_1 + 1) + 1$ .

### 3 Computational results

Here we apply our results to  $\overline{G}$  with  $k = 0$  to obtain bounds on the clique number  $\omega(G) = \alpha(\overline{G})$ . We compare old and new bounds for  $\omega(G)$  for graphs  $G$  from the Second DIMACS Implementation Challenge [12]: these are benchmark graphs used for testing algorithms that determine or estimate  $\omega(G)$ . The old bounds in the table are given by  $1 + \lambda_1(G)$ , while the new bounds  $h_0^{\overline{G}}(t^*)$  are calculated in accordance with Corollary 2.8: if  $\overline{\mu}_s \geq 0$  (in particular, if  $\overline{G}$  is regular) then  $t^* = -\overline{\lambda}_n$ ; otherwise  $h_0^{\overline{G}}(t^*)$  is the unique local minimum on  $(-\overline{\lambda}_n, \infty)$ . In practice,  $t^*$  is determined to within a computational error, and so

$$h_0^{\overline{G}}(t^*) \approx \inf\{h_0^{\overline{G}}(t) : t > -\lambda_n(\overline{G})\}.$$

Most of the graphs in the table have  $\lambda_n(\overline{G})$  as a main eigenvalue, with  $h_0'(1 + \lambda_1(G)) > 0$ , where  $h_0 = h_0^{\overline{G}}$ . Then  $\overline{\mu}_s < 0$  and we estimate  $t^*$  using successive bisections of intervals, starting with  $[-\lambda_n(\overline{G}) + 10^{-6}, \lambda_1(G) + 1]$ , where the value of  $h_0$  at the mid point is less than the value at each end point. For an interval  $[a, b]$  with mid-point  $c$ , let  $x, y$  be the mid points of  $[a, c], [c, b]$  respectively. If  $h_0(x)$  and  $h_0(y)$  are both greater than  $h_0(c)$  then we replace  $[a, b]$  with  $[x, y]$ . Otherwise,  $[a, b]$  is replaced with  $[a, c]$  if  $h_0(x) \leq h_0(c)$ , or with  $[c, b]$  if  $h_0(x) > h_0(c)$ . The process is repeated until we reach an interval where the values of  $h_0$  at the mid point and end points coincide to within four decimal places.

In the graph c-fat200-1.clq,  $-\lambda_n(\overline{G}) - 1$  is a main eigenvalue of  $G$  and  $h_0'(-\lambda_n(\overline{G})) > 0$ ; thus the best upper bound is that in (3), attained when  $t^* = h_0(t^*) = -\lambda_n(\overline{G}) = 17.2675$ .



$G$	$n$	$\omega(G)$	$\lambda_1(G) + 1$	$t^*$	$h_0^{\tilde{G}}(t^*)$	Notes
brock200-1.clq	200	21	149.5707	12.4952	43.3005	(a)
brock200-2.clq	200	12	100.1963	14.0483	26.4234	(a)
brock200-3.clq	200	15	121.8181	13.9645	32.0650	(a)
brock200-4.clq	200	17	132.2037	13.5104	35.3994	(a)
brock400-1.clq	400	27	299.8496	17.2781	62.8351	(a)
brock400-2.clq	400	29	300.1480	17.4017	62.8164	(a)
brock400-3.clq	400	31	299.6317	17.6204	63.9385	(a)
brock400-4.clq	400	33	300.0543	17.5317	63.3207	(a)
c-fat200-1.clq	200	12	17.8135	17.2675	17.2675	
c-fat200-2.clq	200	24	33.6036	32.7001	32.9611	(a)
c-fat200-5.clq	200	58	85.7778	64.7787	72.9051	
hamming6-2.clq	64	32	58		32	(b)
hamming6-4.clq	64	4	23		13.5385	(b)
hamming8-2.clq	256	128	248		128	(b)
hamming8-4.clq	256	16	164		72	(b)
johnson8-2-4.clq	28	4	16		4	(b)
johnson8-4-4.clq	70	14	54		14	(b)
johnson16-2-4.clq	120	8	92		8	(b)
johnson32-2-4.clq	496	16	436		16	(b)
MANN-a9.clq	45	16	41.8039	2.3885	19.7076	
MANN-a27.clq	378	126	374.3035	6.7405	278.9118	
p-hat300-1.clq	300	8	80.7579	16.6554	26.3647	(a)
p-hat300-2.clq	300	25	158.9345	30.3485	78.1328	(a)
p-hat300-3.clq	300	36	225.8307	19.3401	88.3742	(a)
keller4.clq	171	11	111.8552	17.7206	41.1585	
san200-0.7-1.clq	200	30	140.5107	51.6650	94.7681	(a)
san200-0.7-2.clq	200	18	143.5080	68.3020	117.1690	(a)
san200-0.9-1.clq	200	70	180.3256	22.8092	118.7377	(a)
san200-0.9-2.clq	200	60	180.1964	17.4725	98.3736	(a)
san200-0.9-3.clq	200	44	180.1697	14.1434	86.5558	(a)
san400-0.5-1.clq	400	13	202.9588	151.2577	179.3039	(a)
san400-0.7-1.clq	400	40	280.4968	102.2726	184.7757	(a)
san400-0.7-2.clq	400	30	280.5105	98.4703	182.4865	(a)
san400-0.7-3.clq	400	22	280.8343	93.7929	183.7393	(a)

Notes: (a)  $-\lambda_n(\overline{G})$  is a main eigenvalue, (b)  $G$  is regular.

## References

- [1] R. Andersen and S. M. Cioabă, Spectral densest subgraph and independence number of a graph, *J. Universal Computer Science* **13** (2007), 1501-1513.
- [2] D. M. Cardoso, M. Kamiński and V. Lozin, Maximum  $k$ -regular induced subgraphs, *J. Comb. Optim.* **14** (2007), 455-463.
- [3] D. M. Cardoso and S. J. Pinheiro, Spectral upper bounds on the size of  $k$ -regular induced subgraphs, *Electronic Notes in Discrete Mathematics* **32** (2009), 3-10.
- [4] D. Cvetković, The generating function for variations with restrictions and paths of the graph and self-complementary graphs, *Univ. Beograd, Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* Nos. 320-328 (1970), 27-34.
- [5] D. Cvetković, Graphs and their spectra, *Univ. Beograd, Publ. Elektrotehn. Fak. Ser. Mat. Fiz.* Nos. 354-356 (1971), 1-50.
- [6] D. Cvetković and M. Petrić, A table of connected graphs on six vertices, *Discrete Math.* **50** (1984), 37-49.
- [7] D. Cvetković and P. Rowlinson, Some results in the theory of graph spectra, in: *Topics in Algebraic Graph Theory* (eds. L. W. Beineke and R. J. Wilson), Cambridge University Press (Cambridge) 2004, pp.88-112.
- [8] D. Cvetković, P. Rowlinson and S. K. Simić, *Eigenspaces of Graphs*, Cambridge University Press, 1997.
- [9] C. D. Godsil and M. F. Newman, Eigenvalue bounds for independent sets, *J. Combin. Theory, Ser. B* **98** (2008), 721-734.
- [10] C. D. Godsil and G. Royle, *Algebraic Graph Theory*, Springer-Verlag, New York, 2001.
- [11] S. Grünewald, Harmonic trees, *Appl. Math. Lett.* **15** (2002), 1001-1004.
- [12] D. S. Johnson and M A. Trick (Eds.), Cliques, Coloring, and Satisfiability: Second DIMACS Challenge, *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, American Mathematical Society, Providence, 1996.
- [13] V. Nikiforov, More spectral bounds on the clique and independence numbers, *J. Combin. Theory, Ser. B* **99** (2009), 819-826.
- [14] P. Rowlinson, The main eigenvalues of a graph: a survey, *Appl. Anal. Discrete Math.* **1** (2007), 445-471.