

Digit ratio (2D:4D) predicts facial, but not voice or body odour, attractiveness in men

| | |
|-------------------------------|--|
| Journal: | <i>Proceedings B</i> |
| Manuscript ID: | RSPB-2011-0544.R1 |
| Article Type: | Research |
| Date Submitted by the Author: | n/a |
| Complete List of Authors: | Ferdenzi, Camille; University of Geneva, Swiss Center for Affective Sciences Lemaître, Jean-François; Université Lyon 1, Unité Mixte de Recherche 5558, Biométrie et Biologie Evolutive Leongómez, Juan; University of Stirling, Psychology Roberts, S.; University of Stirling, Psychology |
| Subject: | Behaviour < BIOLOGY, Evolution < BIOLOGY |
| Keywords: | Mate Choice, Finger Ratio, Testosterone, Face Symmetry, Masculinity |
| Proceedings B category: | Behavioural Ecology |
| | |

SCHOLARONE™
Manuscripts

Digit ratio predicts face attractiveness

Digit ratio (2D:4D) predicts facial, but not voice or body odour, attractiveness in men

Camille Ferdenzi¹, Jean-François Lemaître², Juan David Leongómez³
and S. Craig Roberts³

¹ Swiss Centre for Affective Sciences, University of Geneva, 7 rue des Batoirs, 1205 Geneva, Switzerland. camille.ferdenzi@unige.ch

² Unité Mixte de Recherche 5558, Biométrie et Biologie Evolutive, Université Lyon 1, 43 Boulevard du 11 novembre 1918, 69622 Villeurbanne Cedex, France. jean-francois.lemaitre@univ-lyon1.fr

³ University of Stirling, Stirling FK9 4LA, Scotland, United Kingdom. j.d.leongomezpena@stir.ac.uk and craig.roberts@stir.ac.uk

Abstract

There is growing evidence that human second-to-fourth digit ratio, or 2D:4D, is related to facial features involved in attractiveness, certainly mediated by *in utero* hormonal effects. The present study extends the investigation to other phenotypic, hormone-related, determinants of human attractiveness: voice and body odour. Pictures of faces with a neutral expression, recordings of voices pronouncing vowels, and axillary odour samples captured on cotton pads worn for 24 hours, were provided by 49 adult male donors. These stimuli were rated on attractiveness and masculinity scales by two groups of 49 and 35 females, approximately half of these in each sample using hormonal contraception. Multivariate regression analyses showed that males' lower (more masculine) right 2D:4D ratio and lower right minus left 2D:4D (Dr-l) were associated with a more attractive, and in some cases more symmetrical, but not more masculine face. However, 2D:4D and Dr-l did not predict voice and body odour masculinity or attractiveness. The results were interpreted in terms of differential effects of prenatal and circulating testosterone, male facial shape being supposedly more dependent on foetal levels (reflected by 2D:4D ratio), whereas body odour and vocal characteristics could be more dependent on variation in adult circulating testosterone levels.

Keywords: Mate choice; Finger Ratio; Testosterone; Face Symmetry; Masculinity.

1. INTRODUCTION

The relative length of the second (index) and fourth (ring) fingers, or 2D:4D ratio, is sexually dimorphic in several species, with lower 2D:4D ratios for males than females in mammals [1,2], while in birds it appears to be the reverse [3]. Although the precise genetic mechanism explaining this sexual dimorphism is still unclear, there is compelling evidence that *in utero* foetal testosterone and foetal estrogen influence 2D:4D ratio in humans [1,4-6]. For example, males suffering from congenital adrenal hyperplasia (CAH), an enzymatic deficiency that entails excessive levels of androgens during the foetal period, have particularly low 2D:4D ratio [7]. More generally, men exposed to high levels of prenatal androgens develop low 2D:4D ratio [4,7].

Androgens such as testosterone are also involved in the development and maintenance of secondary sexual characters and thereby in mate choice [8,9]. Because maintaining a high level of testosterone is costly for males [e.g., 10,11], those that display enhanced sexual characters without suffering too much from immunosuppression are considered as high quality males [12]. Therefore, women should ultimately increase their reproductive success by choosing mates displaying testosterone-dependent sexual traits [12,13]. In humans, men with higher levels of circulating testosterone have voices with lower fundamental frequency [14] and more masculine faces [15,16], two traits that are preferred by women when they become sexually mature (see [17] for voices, and [18] for faces).

Since the growth of the 4th finger is dependent on the level of prenatal androgen, and since some authors have hypothesized a positive correlation between prenatal and adult testosterone levels [1,19], 2D:4D ratios might correlate negatively with some other

Digit ratio predicts face attractiveness

testosterone-dependent traits [1]. If these traits such as voices and faces are sexually selected, then measures of 2D:4D should be a good predictor of men's attractiveness. To date, investigations of these putative relationships are scarce and remain principally focused on face and body masculinity of men (e.g. [20]) since this trait is testosterone-dependent [16] and preferred by women in a mate choice context [18]. Moreover, results from these studies are conflicting. For example, Neave *et al.* [20] found a negative correlation between 2D:4D ratios of the left and right hand and the female perception of male facial dominance and masculinity but Koehler *et al.* [21] failed to repeat these results and found no relationship between 2D:4D and body and face masculinity. Furthermore, some authors found a link between 2D:4D and attractiveness [22-24], whereas others did not [20].

To date, studies testing relationships between 2D:4D ratios and sexually selected traits are only focused on men's bodies and faces although there is evidence that women use multiple testosterone-dependent cues to select mates, such as voice [25] and body odour [26,27]. As for faces [15], voice frequency and thus attractiveness are related to the level of salivary testosterone [14]. Similarly, androgen level is likely to influence body odour since steroid compounds of axillary odour such as androstadienone are more present in males [28] and are products of testosterone transformation by underarm bacteria [29,30].

In this study, for the first time, we investigated in three sensory modalities involved in human mate choice (voice, body odour and face) whether second-to-fourth digit ratio of left and right hands, and digit ratio difference between the two hands (Dr-I, also related to prenatal testosterone sensitivity [31,32]), can predict men masculinity and attractiveness. We predicted that 2D:4D ratio and Dr-I would be negatively correlated with face, voice and odour masculinity and attractiveness, as evaluated by females. As voice frequency and face

Digit ratio predicts face attractiveness

symmetry influence women preference for men (deeper voices are preferred [33]; more symmetrical faces are more attractive [34,35]), we also measured these two factors and linked them to 2D:4D ratios. Finally, we controlled for the use of hormonal contraceptives by the female raters, since this could alter women preference for various male features such as body odour, face and voice [36,37].

2. METHODS

(a) *Participants*

Participants were 49 Caucasian male donors aged between 18 and 33 years old (Mean \pm SD = 22.3 \pm 4.0 yrs), recruited among students of the University of Liverpool. From these, we obtained 2D:4D measures, a voice sample and a facial photograph. Axillary odour samples were collected for 28 of them who were non-smokers, as is standard in odour rating research because of the influence of smoking on body odour quality [38,39].

Male axillary odour samples were evaluated by 49 Caucasian female students of the University of Liverpool, aged between 19 and 34 years old (Mean \pm SD = 21.8 \pm 3.2 yrs). Of these, 26 reported taking hormone-based contraception (hereafter named 'pill users') and 23 were not (hereafter named 'non-pill users'). Each odour sample was rated fresh by nine to ten women during one of five rating sessions at the University of Liverpool, between November 2007 and February 2008. Men's faces were judged by 27 of these women (Mean \pm SD = 21.8 \pm 3.4 yrs, 14 'pill users', 13 'non-pill users'). Due to experimental constraints, the voices were evaluated later (November and December 2010) by a separate group of female students of the University of Stirling (n = 35, Mean \pm SD = 20.1 \pm 3.5, range: 18 to 34 yrs, 20 'pill users', 15 'non-pill users', Caucasian). Although both groups of raters were similar in terms

of age, culture (British) and occupation (students), we controlled for consistency of their evaluations. Hence, we asked the voice raters to rate the men's faces, previously rated by the Liverpool group. Rating of both groups were highly consistent for face short-term attractiveness (Intra-class Correlation Coefficient: $ICC = 0.944, p < 0.001$), long-term attractiveness ($ICC = 0.942, p < 0.001$), masculinity ($ICC = 0.923, p < 0.001$) and symmetry ($ICC = 0.889, p < 0.001$). Therefore, both groups were considered as equivalent. All participants gave their written informed consent, and the study was approved by the Committee on Research Ethics of the University of Liverpool and of the University of Stirling.

(b) Voice samples

Participants' voice was recorded on a digital recorder (M-Audio Microtrack 24/96) with a cardioid condenser microphone (Technica ATR55 Telemike Shotgun), in a quiet room at about 15 cm from the microphone. Participants were required to recite two sentences of *the rainbow passage* [40] and the monophthong vowels "eh", "ee", "ah", "oh" and "oo". This sequence was then repeated once. Female ratings and measure of voice frequency were performed on the second repetition (when participants are more relaxed) and on the three vowels in middle ("ee", "ah", "oh"; see Supplementary Material 1) to limit intonation variations. Voice frequency F_0 was measured with Praat 4.6 (www.praat.org). Voice attractiveness and masculinity ratings were collected on 1-to-7 scales with E-Prime Software (2.0, Psychology Software Tools), after equalizing the samples in intensity (in Matlab 7.10) and length (2 sec, in Praat).

Digit ratio predicts face attractiveness

143 **(c) Axillary odour samples**

144 Axillary odour samples were collected on cotton pads (9.5 x 6.5 cm, Boots UK Ltd)
145 fastened onto both axillae for 24 hours. Participants were instructed to refrain from eating
146 strong food, drinking alcohol, smoking, doing sport and having sexual intercourse, 2 days
147 before and during odour collection. They were also required to shower with a non-perfumed
148 soap before fastening the pads, and not to use any scented products such as perfume or
149 deodorant. Samples were presented fresh to female raters a few hours (range: 2-8 hours) after
150 pads were removed from the armpits. Odour samples were placed in glass flasks, presented in
151 a random order to the raters, and evaluated for attractiveness and masculinity on 9-point
152 scales. The variable used in this study was the average ratings of the right and left side. For
153 more details about the procedure of odour collection and rating, see [38].

154

155 **(d) Face samples**

156 Full face pictures of the male participants were taken in standardized conditions of light
157 with a Canon Powershot camera. Participants were asked to have a neutral expression and to
158 look at the camera without any vertical or horizontal tilt of the head. Distance to the camera
159 was constant and participants wore a dark gown. Images were resampled to 400x480 pixels
160 with resolution 72 dpi. Using Psychomorph 8.4 (Perrett & Tiddeman, University of St
161 Andrews, UK), faces were normalized according to pupils and mouth position, and face
162 symmetry was computed using 7 bilateral points (pupils, outermost and innermost eye
163 corners, leftmost and rightmost points of the nose, mouth corners, cheekbones and jaws;
164 Supplementary Material 2). The asymmetry index was the sum of the vertical and horizontal
165 asymmetry indices. Vertical and horizontal asymmetries were respectively the sum of
166 differences in vertical and horizontal locations of each of the seven facial features (see details
167 in [41]). Placement of the points and computation of the asymmetry index were performed

Digit ratio predicts face attractiveness

twice, and averaged since the two asymmetry indices were highly consistent ($ICC = 0.876$, $p < 0.001$). Men's faces were presented in random order with a java applet. Female participants were asked to rate the faces for short-term attractiveness (i.e., considering the person as a dinner date or holiday romance), long-term attractiveness (i.e., considering the person as a long-term partner), masculinity and symmetry of the faces on 1-to-7 scales. They were asked to skip the ratings of the men they knew.

(e) Measures of 2D:4D

The length of index and ring fingers of the male participants was measured to the nearest 0.1 mm using Vernier callipers, directly on fingers (more reliable than indirect measures performed on a photocopy of the hands [42]). Measurement was taken from the most proximal ventral crease of the digit to the tip of the finger. To limit measurement errors, the procedure was repeated three times, and as the measures were highly correlated ($ICC = 0.986$, $p < 0.001$) they were averaged. The index-to-ring ratio (2D:4D) for the left and right hand separately were then computed, as well as the difference between right and left 2D:4D (Dr-l).

(f) Data analysis

All variables had normal distributions (assessed by Kolmogorov-Smirnov tests) and parametric statistics were thus used. In addition, no extreme values were to be removed before performing analyzes. Tests were two-tailed and were conducted using Statistica 9.0 and SPSS 18.0. The link between 2D:4D and visual, auditory, and olfactory stimuli was investigated using multivariate simple regressions, with face, voice, and odour characteristics (masculinity, attractiveness, etc.) as dependent variables and 2D:4D as predictor. The difference between 'pill users' and 'non-pill users' was tested with paired t -tests and the

relation between masculinity, attractiveness and other dimensions was assessed with Pearson correlation coefficients.

3. RESULTS

(a) *Voice*

First, correlations between male voice frequency and both rated attractiveness and rated masculinity were significantly negative ($r = -0.69$ and $r = -0.63$, respectively, $n = 48$, $p < 0.001$). Attractiveness and masculinity correlated positively ($r = 0.77$, $n = 48$, $p < 0.001$). ‘Pill users’ gave slightly higher attractiveness ratings than the ‘non-pill users’ ($t_{47} = 2.14$, $p = 0.038$), but the two groups did not differ on the masculinity ratings ($t_{47} = 0.95$, $p = 0.35$) (Supplementary Material 3).

Multivariate simple regressions were performed to determine whether 2D:4D ratio of right hand, 2D:4D of left hand, and difference between right and left 2D:4D (Dr-l), were significant predictors of voice frequency, and rated attractiveness and masculinity. Voice frequency and voice attractiveness were predicted neither by the right 2D:4D ratio, Dr-l (table 1) nor left 2D:4D (Supplementary Material 4). Voice masculinity was predicted only by left hand 2D:4D when ‘non-pill users’ were taken into account (Supplementary Material 4).

(b) *Body odour*

The correlation between masculinity and attractiveness of males’ body odours was significantly negative ($r = -0.54$, $n = 28$, $p = 0.003$). Average ratings of the ‘pill users’ and ‘non-pill users’ did not differ (attractiveness: $t_{27} = 0.44$, $p = 0.66$; masculinity: $t_{27} = 0.01$, $p = 0.99$; Supplementary Material 3).

Digit ratio predicts face attractiveness

As for voice ratings, multivariate simple regressions were performed to determine whether 2D:4D ratio of right and left hand were significant predictors of body odour attractiveness and masculinity. There were significant effects for 2D:4D of the right hand only. Although masculinity was not predicted by 2D:4D (right, left, Dr-l), attractiveness was (by right 2D:4D) when only ‘non-pill users’ were taken into account (table 1, Supplementary Material 4).

(c) Face

First, masculinity was correlated neither with attractiveness (short-term attractiveness: $r = 0.15$, $p = 0.30$; long-term attractiveness: 0.24 , $p = 0.10$; $n = 47$) nor with face symmetry (perceived by females: $r = 0.20$, $p = 0.17$; measured in Psychomorph: $r = 0.05$, $p = 0.73$; $n = 47$). Long-term and short-term attractiveness were highly correlated ($r = 0.96$, $n = 47$, $p < 0.001$), and symmetry rated and perceived were correlated too ($r = 0.39$, $n = 47$, $p < 0.01$). Attractiveness was correlated with perceived face symmetry (short-term attractiveness: $r = 0.67$, $p < 0.001$; long-term attractiveness: 0.66 , $p < 0.001$; $n = 47$), but not or only marginally with measured face symmetry (short-term attractiveness: $r = -0.22$, $p = 0.14$; and long-term attractiveness: $r = -0.28$, $p = 0.06$; $n = 47$). Contrary to odours, mean face ratings of the ‘pill users’ and ‘non-pill users’ significantly differed. Compared to ‘pill users’, ‘non-pill users’ gave higher attractiveness (short-term: $t_{46} = 1.99$, $p = 0.052$; long-term: $t_{27} = 6.06$, $p < 0.001$), higher masculinity ($t_{27} = 4.59$, $p < 0.001$) and higher symmetry ratings ($t_{27} = 8.03$, $p < 0.001$) (Supplementary Material 3).

As for voice and odour ratings, multivariate simple regressions were performed to determine whether right 2D:4D, left 2D:4D, and Dr-l were significant predictors of face attractiveness and masculinity. There were significant effects for 2D:4D of the right hand and for the right-left difference Dr-l (table 1). Long-term and short-term attractiveness were

Digit ratio predicts face attractiveness

significantly predicted by right 2D:4D and by Dr-I. Perceived symmetry was predicted by right 2D:4D only, and face masculinity was not predicted by any of the 2D:4D variables. These results remained unchanged when ‘pill users’ and ‘non-pill users’ are analyzed separately.

4. DISCUSSION

In this study, we tested whether second-to-fourth digit ratio (2D:4D) of left hand, right hand and right minus left hand Dr-I, can predict men masculinity and attractiveness for three sensory modalities involved in human mate choice: voice, body odour and face. Our main finding is that right hand 2D:4D and Dr-I are significant predictors of attractiveness but not masculinity of male faces, whether they are considered as short-term or long-term potential partners. Right hand 2D:4D also predicts perceived facial symmetry. The link between 2D:4D and facial attractiveness is consistent with previous studies investigating either self-evaluated attractiveness [22,23] or men’s attractiveness rated by women [22,24]. This illustrates a female preference for low 2D:4D men, possibly driven by the fact that these have more symmetrical faces. Such a preference might have evolved because it increases females’ reproductive success by gaining benefits from partners who are physically more robust [1] and who have more fertile ejaculates [43,44].

Our results differ from other studies that found significant relationship between dominant/masculine personality traits and 2D:4D [45], and from a study by Neave *et al.* [20] who report a similar negative association between 2D:4D ratio (of both hands) and masculinity, but not attractiveness. However, ours and Neave *et al.*’s results are not necessarily contradictory. Indeed, men who are able to pay costs of high levels of testosterone (see [12]), will consequently develop masculine phenotypes [15]. At the same time,

Digit ratio predicts face attractiveness

symmetric faces are likely to be found in individuals who have a better developmental stability [13], which reflects a better resistance to parasites and poor environmental conditions [46]. Therefore, some particularly good quality males should express simultaneously masculine and symmetric faces. Thus, one could expect 2D:4D to predict both face masculinity and symmetry, and the fact that only one of these predictions were found in our and Neave *et al.*'s study might be an effect of sampling or of differences in 2D:4D measurement procedure. These effects are likely to be subtle since other studies failed, as we did, to find a link between 2D:4D ratio and masculinity features [21].

Replication of our findings, the direction of which contradicts another study on 2D:4D and facial symmetry [47], would thus be worthwhile. Furthermore, future research will be necessary to better understand the relationship between prenatal androgen exposure and adult face attractiveness. Our study cannot directly address the mechanism underlying this relationship, and the present results provide no evidence that prenatal testosterone is the causal factor of both low 2D:4D and high attractiveness (via face symmetry). Indeed, it may be possible that the causal factor that explains the relationship between 2D:4D and attractiveness is situated at another level. For example, a high level of parental attractiveness, because it reflects genetic quality, might provide the foetus both with 'good genes' (high level of testosterone) and a healthy prenatal environment allowing high developmental stability.

The significant link between male facial attributes and 2D:4D ratio we found was observed for the right hand only, which has a more male-like ratio than the left hand (right: $M = 0.97$, $SD = 0.03$; left: $M = 0.98$, $SD = 0.02$; $t_{48} = 3.57$, $p < 0.001$). This result supports the assumption of Tanner (1990, cited in [1]), according to which "sexually dimorphic traits tend to be expressed in the male form more strongly on the right side of the body". This side-

Digit ratio predicts face attractiveness

related difference is also supported by Manning *et al.* [43], who found stronger association between testosterone levels and 2D:4D ratios on the right hand compared to the left hand, as well as by other authors [4]. The authors hypothesize a stronger action of androgens on the digits of the right hand (see [48] for a meta-analysis).

Surprisingly, we found a positive relationship between 2D:4D ratios of men and the evaluation of their body odour attractiveness and voice masculinity by the ‘non-pill users’. This result is contrary to our predictions and deserves further investigation, especially taking into account the impact of cycle stage on this kind of evaluations. In this respect, we found that spontaneously ovulating women gave higher facial attractiveness, masculinity and symmetry ratings than ‘pill users’, which is concordant with previous studies showing that fertile women prefer less feminized [49] and more symmetrical male faces ([50] but see [51]). However, this result was not confirmed in the second group of females who evaluated the faces, which might be due to the proportion of women being in their fertile phase during data collection, a factor that we did not control.

The fact that 2D:4D does not reliably predict voice and body odour attractiveness or masculinity is not due to the fact that different groups of females rated faces and voices. Indeed, not only both groups were highly correlated, but we also performed the regressions of table 1 and Supplementary Material 4 again with the data of the ‘voice raters’ group and the results were replicated (the only difference being a lower level of significance for the effects in pill users, detail of the results not presented here). Rather, this absence of relationship between 2D:4D ratio and vocal and olfactory traits might stem from the fact that voice and body odour are by nature more variable than facial shape and more related to current circulating levels of testosterone in the adult individual (but see [52]). Consistent with this

hypothesis, Evans *et al.* [14] found that voice frequency is related to the level of circulating testosterone but not to the indicator of prenatal testosterone level 2D:4D (see also [53,54]), whereas the reverse was found for faces ([20], but see [15]). All together, these results raise the question of the relative dissociation between organisational and activational effects of testosterone [8], organisational effects being irreversible and occurring during sensitive periods of early development, and activational ones being impermanent and occurring in adulthood. Foetal testosterone might serve to organize certain features of the face like bones (jaws and cheek bones) that will subsequently be activated during puberty and remain relatively stable thereafter. On the contrary, voice is produced by the larynx that is made of muscles and cartilage, and body odour consists in the degradation of products of the metabolism by skin bacteria: both of them are likely to change at anytime under the influence of circulating hormones. Indeed, voice quality significantly changes with therapeutic administration of testosterone (e.g. [55]) or more subtly with normal daily variations of testosterone concentration [14], and some major compounds of axillary odours are by-products of androgen substances [29].

5. CONCLUSION

Our study suggests that both right hand 2D:4D and right-minus-left 2D:4D (Dr-l) are good predictors of facial attractiveness in men, but not of their voice or body odour attractiveness. We showed, for the first time, that this effect might be supported by the link between 2D:4D and face symmetry, one indicator of male quality. Physical features closely linked to foetal levels of testosterone, such as face shape, are more likely to be correlated with second-to-fourth digit ratios than traits believed to be directly controlled by circulating

Digit ratio predicts face attractiveness

level of testosterone later in life history (voice and body odour). We advocate that more research is needed to investigate the action of both early and adult testosterone levels on the development of sexually dimorphic traits involved in human attractiveness, including those we have examined here. The present study suggests that masculine and attractive features of voice and body odour might not be shaped *in utero* but later during life history: the timing (and even the existence, for body odour) of an action of testosterone on these two modalities remain to be elucidated in the future.

Acknowledgments

The authors wish to thank the participants, Kirsty Elliott and Marta Vondrova for their help in data collection, Rob Burriss, Hannah Rowland and Tamsin Saxton for technical advice, and Bernard Tiddeman and David Perrett for use of the Psychomorph software. One anonymous reviewer and Markus Rantala are also thanked for their constructive comments on the manuscript. This research was supported by Fyssen Foundation, Paris.

REFERENCES

1. Manning, J. T. 2002 *Digit ratio: A pointer to fertility, behavior, and health*. New Brunswick, NJ: Rutgers University Press.
2. Nelson, E. & Shultz, S. 2010 Finger length ratios (2D:4D) in anthropoids implicate reduced prenatal androgens in social bonding. *Am. J. Phys. Anthropol.* **141**, 395-405. (doi:10.1002/ajpa.21157).
3. Burley, N. T. & Foster, V. S. 2004 Digit ratio varies with sex, egg order and strength of mate preference in zebra finches. *Proc. R. Soc. B* **271**, 239-244. (doi:10.1098/rspb.2003.2562).
4. Lutchmaya, S., Baron-Cohen, S., Raggatt, P., Knickmeyer, R. & Manning, J. T. 2004 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Hum. Dev.* **77**, 23-28.

Digit ratio predicts face attractiveness

(doi:10.1016/j.earlhumdev.2003.12.002).

5. Breedlove, S. M. 2010 Minireview: Organizational hypothesis: instances of the fingerpost. *Endocrinology* **151**, 4116-4122. (doi:10.1210/en.2010-0041).

6. van Honk, J., Schutter, D. J., Bos, P. A., Kruijt, A., Lentjes, E. G. & Baron-Cohen, S. In press. Testosterone administration impairs cognitive empathy in women depending on second-to-fourth digit ratio. *PNAS*. (doi:10.1073/pnas.1011891108).

7. Okten, A., Kalyoncu, M. & Yariş, N. 2002 The ratio of second- and fourth-digit lengths and congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Early Hum. Dev.* **70**, 47-54. (doi:10.1016/S0378-3782(02)00073-7).

8. Neave, N. 2008 *Hormones and Behaviour: A Psychological Approach*. Cambridge: Cambridge University Press.

9. Andersson, M. B. 1994 *Sexual Selection*. Princeton, NJ: Princeton University Press.

10. Reed, W. L., Clark, M. E., Parker, P. G., Raouf, S. A., Arguedas, N., Monk, D. S., Snajdr, E., Nolan, V. & Ketterson, E. D. 2006 Physiological effects on demography: a long-term experimental study of testosterone's effects on fitness. *Am. Nat.* **167**, 667-683. (doi:10.1086/503054).

11. Mills, S. C., Grapputo, A., Jokinen, I., Koskela, E., Mappes, T. & Poikonen, T. 2010 Fitness trade-offs mediated by immunosuppression costs in a small mammal. *Evolution* **64**, 166-179. (doi:10.1111/j.1558-5646.2009.00820.x).

12. Folstad, I. & Karter, A. J. 1992 Parasites, bright males, and the immunocompetence handicap. *Am. Nat.* **139**, 603-622. (doi:10.1086/281919).

13. Thornhill, R. & Gangestad, S. W. 1999 The scent of symmetry: a human sex pheromone that signals fitness? *Evol. Hum. Behav.* **20**, 175-201. (doi:10.1016/S1090-5138(99)00005-7).

14. Evans, S., Neave, N., Wakelin, D. & Hamilton, C. 2008 The relationship between testosterone and vocal frequencies in human males. *Physiol. Behav.* **93**, 783-788. (doi:10.1016/j.physbeh.2007.11.033).

15. Penton-Voak, I. S. & Chen, J. Y. 2004 High salivary testosterone is linked to masculine male facial appearance in humans. *Evol. Hum. Behav.* **25**, 229-241. (doi:10.1016/j.evolhumbehav.2004.04.003).

16. Pound, N., Penton-Voak, I. S. & Surridge, A. K. 2009 Testosterone responses to competition in men are related to facial masculinity. *Proc. R. Soc. B* **276**, 153-159. (doi:10.1098/rspb.2008.0990).

17. Saxton, T. K., Debruine, L. M., Jones, B. C., Little, A. C. & Roberts, S. C. 2009 Face and voice attractiveness judgments change during adolescence. *Evol. Hum. Behav.* **30**, 398-408. (doi:10.1016/j.evolhumbehav.2009.06.004).

18. Little, A. C., Saxton, T. K., Roberts, S. C., Jones, B. C., Debruine, L. M., Vukovic, J.,

Digit ratio predicts face attractiveness

- 424 Perrett, D. I., Feinberg, D. R. & Chenore, T. 2010 Women's preferences for masculinity in
 425 male faces are highest during reproductive age range and lower around puberty and post-
 426 menopause. *Psychoneuroendocrinology* **35**, 912-920. (doi:10.1016/j.psyneuen.2009.12.006).
 427
- 428 19. Jamison, S. C., Meier, R. J. & Campbell, B. C. 1993 Dermatoglyphic asymmetry and
 429 testosterone levels in normal males. *Am. J. Phys. Anthropol.* **90**, 185-198.
 430 (doi:10.1002/ajpa.1330900205).
 431
- 432 20. Neave, N., Laing, S., Fink, B. & Manning, J. T. 2003 Second to fourth digit ratio,
 433 testosterone and perceived male dominance. *Proc. R. Soc. B* **270**, 2167-2172.
 434 (doi:10.1098/rspb.2003.2502).
 435
- 436 21. Koehler, N., Simmons, L. W. & Rhodes, G. 2004 How well does second-to-fourth-digit
 437 ratio in hands correlate with other indications of masculinity in males? *Proc. R. Soc. B* **271**,
 438 S296-298. (doi:10.1098/rsbl.2004.0163).
 439
- 440 22. Bogaert, A. F., Fawcett, C. C. & Jamieson, L. K. 2009 Attractiveness, body size,
 441 masculine sex roles and 2D:4D ratios in men. *Pers. Individ. Differ.* **47**, 273-278.
 442 (doi:10.1016/j.paid.2009.03.011).
 443
- 444 23. Manning, J. T. & Quinton, S. 2007 Association of digit ratio (2D:4D) with self-reported
 445 attractiveness in men and women: evidence from the BBC internet survey. *J. Individ. Diff.*
 446 **28**, 73-77. (doi:10.1027/1614-0001.28.2.73).
 447
- 448 24. Roney, J. R. & Maestripieri, D. 2004 Relative digit lengths predict men's behavior and
 449 attractiveness during social interactions with women. *Hum. Nature* **15**, 271-282.
 450 (doi:10.1007/s12110-004-1009-5).
 451
- 452 25. Collins, S. 2000 Men's voices and women's choices. *Anim. Behav.* **60**, 773-780.
 453 (doi:10.1006/anbe.2000.1523).
 454
- 455 26. Havlicek, J., Saxton, T. K., Roberts, S. C., Jozifkova, E., Lhota, S., Valentova, J. & Flegr,
 456 J. 2008 He sees, she smells? Male and female reports of sensory reliance in mate choice and
 457 non-mate choice contexts. *Pers. Individ. Differ.* **45**, 565-570.
 458 (doi:doi:10.1016/j.paid.2008.06.019).
 459
- 460 27. Wedekind, C., Seebeck, T., Bettens, F. & Paepke, A. J. 1995 MHC-dependent mate
 461 preferences in humans. *Proc. R. Soc. B* **260**, 245-249. (doi:10.1098/rspb.1995.0087).
 462
- 463 28. Bird, S. & Gower, D. B. 1981 The validation and use of a radioimmunoassay for 5 alpha-
 464 androst-16-en-3-one in human axillary collections. *J. Steroid Biochem.* **14**, 213-219.
 465 (doi:10.1016/0022-4731(81)90176-X).
 466
- 467 29. Rennie, P. J., Holland, K. T., Mallet, A. I., Watkins, W. J. & Gower, D. B. 1989
 468 Testosterone metabolism by human axillary bacteria. *Biochem. Soc. Trans.* **17**, 1017-1018.
 469 (doi:10.1042/bst0171017).
 470
- 471 30. Havlicek, J., Murray, A. K., Saxton, T. K. & Roberts, S. C. 2010 Current issues in the
 472 study of androstenes in human chemosignalling. *Vitam. Horm.* **83**, 47-81.
 473 (doi:10.1016/S0083-6729(10)83003-1).

Digit ratio predicts face attractiveness

31. Manning, J. T., Bundred, P. E., Newton, D. J. & Flanagan, B. F. 2003 The second to fourth digit ratio and variation in the androgen receptor gene. *Evol. Hum. Behav.* **24**, 399-405. (doi:10.1016/S1090-5138(03)00052-7).
32. Hurd, P. L., Vaillancourt, K. L. & Dinsdale, N. L. In press. Aggression, digit ratio and variation in androgen receptor and monoamine oxidase A genes in men. *Behav. Genet.* (doi:10.1007/s10519-010-9404-7).
33. Feinberg, D. R., Jones, B. C., Little, A. C., Burt, D. M. & Perrett, D. I. 2005 Manipulations of fundamental and formant frequencies influence the attractiveness of human male voices. *Anim. Behav.* **69**, 561-568. (doi:10.1016/j.anbehav.2004.06.012).
34. Grammer, K. & Thornhill, R. 1994 Human (*Homo sapiens*) facial attractiveness and sexual selection: the role of symmetry and averageness. *J. Comp. Psychol.* **108**, 233-242. (doi:10.1037/0735-7036.108.3.233).
35. Perrett, D. I., Burt, D. M., Penton-Voak, I. S., Lee, K. J., Rowland, D. A. & Edwards, R. 1999 Symmetry and human facial attractiveness. *Evol. Hum. Behav.* **20**, 295-307. (doi:10.1016/S1090-5138(99)00014-8).
36. Alvergne, A. & Lummaa, V. 2010 Does the contraceptive pill alter mate choice in humans? *Trends Ecol. Evol.* **25**, 171-179. (doi:10.1016/j.tree.2009.08.003).
37. Roberts, S. C., Gosling, L. M., Carter, V. & Petrie, M. 2008 MHC-correlated odour preferences in humans and the use of oral contraceptives. *Proc. R. Soc. B* **275**, 2715-2722. (doi:10.1098/rspb.2008.0825).
38. Ferdenzi, C., Schaal, B. & Roberts, S. C. 2009 Human axillary odor: are there side-related perceptual differences? *Chem. Senses* **34**, 565-571. (doi:10.1093/chemse/bjp037).
39. Havlicek, J. & Lenochova, P. 2006 The effect of meat consumption on body odor attractiveness. *Chem. Senses* **31**, 747-752. (doi:10.1093/chemse/bjl017).
40. Fairbanks, G. 1960 *Voice and Articulation Drillbook*. New York: Harper & Row.
41. Scheib, J. E., Gangestad, S. W. & Thornhill, R. 1999 Facial attractiveness, symmetry and cues of good genes. *Proc. R. Soc. B* **266**, 1913-1917. (doi:10.1098/rspb.1999.0866).
42. Manning, J. T., Baron-Cohen, S., Wheelwright, S. & Fink, B. 2010 Is digit ratio (2D:4D) related to systemizing and empathizing? Evidence from direct finger measurements reported in the BBC internet survey. *Pers. Individ. Differ.* **48**, 767-771. (doi:10.1016/j.paid.2010.01.030).
43. Manning, J. T., Scutt, D., Wilson, J. & Lewis-Jones, D. I. 1998 The ratio of 2nd to 4th digit length: a predictor of sperm numbers and concentrations of testosterone, luteinizing hormone and oestrogen. *Hum. Reprod.* **13**, 3000-3004. (doi:10.1093/humrep/13.11.3000).
44. Auger, J. & Eustache, F. In press. Second to fourth digit ratios, male genital development and reproductive health: a clinical study among fertile men and testis cancer patients. *Int. J.*

Digit ratio predicts face attractiveness

- 524 *Androl.* (doi:10.1111/j.1365-2605.2010.01124.x).
- 525
- 526 45. Manning, J. T. & Fink, B. 2008 Digit ratio (2D:4D), dominance, reproductive success,
 527 asymmetry, and sociosexuality in the BBC Internet Study. *Am. J. Hum. Biol.* **20**, 451-461.
 528 (doi:10.1002/ajhb.20767).
- 529
- 530 46. Rhodes, G. 2006 The evolutionary psychology of facial beauty. *Annu. Rev. Psychol.* **57**,
 531 199-226. (doi:10.1146/annurev.psych.57.102904.190208).
- 532
- 533 47. Fink, B., Manning, J. T., Neave, N. & Grammer, K. 2004 Second to fourth digit ratio and
 534 facial asymmetry. *Evol. Hum. Behav.* **25**, 125-132. (doi:10.1016/S1090-5138(03)00084-9).
- 535
- 536 48. Hönekopp, J. & Watson, S. 2010 Meta-analysis of digit ratio 2D:4D shows greater sex
 537 difference in the right hand. *Am. J. Hum. Biol.* **22**, 619-630. (doi:10.1002/ajhb.21054).
- 538
- 539 49. Penton-Voak, I. S., Perrett, D. I., Castles, D. L., Kobayashi, T., Burt, D. M., Murray, L.
 540 K. & Minamisawa, R. 1999 Menstrual cycle alters face preference. *Nature* **399**, 741-742.
 541 (doi:10.1038/21557).
- 542
- 543 50. Little, A. C., Jones, B. C., Burt, D. M. & Perrett, D. I. 2007 Preferences for symmetry in
 544 faces change across the menstrual cycle. *Biol. Psychol.* **76**, 209-216.
 545 (doi:10.1016/j.biopsycho.2007.08.003).
- 546
- 547 51. Peters, M., Simmons, L. W. & Rhodes, G. 2009 Preferences across the menstrual cycle
 548 for masculinity and symmetry in photographs of male faces and bodies. *PLoS ONE* **4**, e4138.
 549 (doi:10.1371/journal.pone.0004138).
- 550
- 551 52. Rantala, M. J., Eriksson, C. J. P., Vainikka, A. & Kortet, R. 2006 Male steroid hormones
 552 and female preference for male body odor. *Evol. Hum. Behav.* **27**, 259-269.
 553 (doi:10.1016/j.evolhumbehav.2005.11.002).
- 554
- 555 53. Hughes, S. M., Harrison, M. A. & Gallup, G. G. 2002 The sound of symmetry: voice as a
 556 marker of developmental instability. *Evol. Hum. Behav.* **23**, 173-180. (doi:10.1016/S1090-
 557 5138(01)00099-X).
- 558
- 559 54. Puts, D. A., Gaulin, S. J. C., Sporter, R. J. & McBurney, D. H. 2004 Sex hormones and
 560 finger length. What does 2D:4D indicate? *Evol. Hum. Behav.* **25**, 182-199.
 561 (doi:10.1016/j.evolhumbehav.2004.03.005).
- 562
- 563 55. Akcam, T., Bolu, E., Merati, A. L., Durmus, C., Gerek, M. & Ozkaptan, Y. 2004 Voice
 564 changes after androgen therapy for hypogonadotrophic hypogonadism. *Laryngoscope* **114**,
 565 1587-1591. (doi:10.1097/00005537-200409000-00016).

566 **TABLE CAPTION**

567

568 **Table 1** - Link between digit ratio (2D:4D) of the right hand and right hand minus left hand
569 2D:4D (Dr-l), and voice, odour and face characteristics of 49 men, determined by a simple
570 linear regression (voice pitch, R^2) and multivariate linear regressions (other measures, β). df:
571 degrees of freedom; *: $p < 0.05$; **: $p < 0.01$. A negative β value indicates an inverse
572 relationship between 2D:4D ratio and the dependent variable. Results for left 2D:4D are
573 presented in Supplementary Material 4.

Digit ratio predicts face attractiveness

| Modality | Raters | Dimension | Right 2D:4D | | | | Right minus Left 2D:4D (Dr-l) | | | |
|----------|------------------|--------------------------------|-----------------|-------------|--------------|----------|-------------------------------|-------------|-------------|----------|
| | | | Wilks λ | R^2/β | $F(df)$ | P | Wilks λ | R^2/β | $F(df)$ | P |
| Voice | n/a | Frequency | | 0.02 | (1,46) 0.88 | 0.353 | | 0.01 | (1,46) 0.00 | 0.949 |
| | | Attractiveness | | 0.19 | (1,46) 1.78 | 0.198 | | 0.05 | (1,46) 0.10 | 0.749 |
| | Total | Masculinity | 0.96 | 0.17 | (1,46) 1.43 | 0.237 | 0.98 | -0.04 | (1,46) 0.09 | 0.771 |
| | 'Pill users' | Attractiveness | | 0.18 | (1,46) 1.58 | 0.215 | | 0.01 | (1,46) 0.01 | 0.924 |
| | | Masculinity | 0.97 | 0.16 | (1,46) 1.16 | 0.287 | 1.00 | -0.02 | (1,46) 0.02 | 0.881 |
| | 'Non-pill users' | Attractiveness | | 0.20 | (1,46) 1.82 | 0.183 | | 0.09 | (1,46) 0.37 | 0.547 |
| | | Masculinity | 0.95 | 0.19 | (1,46) 1.77 | 0.190 | 0.96 | -0.07 | (1,46) 0.24 | 0.625 |
| | | | | | | | | | | |
| Odour | Total | Attractiveness | | 0.25 | (1,26) 1.78 | 0.193 | | 0.09 | (1,26) 0.24 | 0.632 |
| | | Masculinity | 0.89 | -0.32 | (1,26) 2.98 | 0.096 | 0.93 | -0.27 | (1,26) 2.00 | 0.169 |
| | 'Pill users' | Attractiveness | | 0.05 | (1,26) 0.06 | 0.811 | | -0.04 | (1,26) 0.05 | 0.831 |
| | | Masculinity | 0.90 | -0.31 | (1,26) 2.81 | 0.105 | 0.99 | -0.06 | (1,26) 0.11 | 0.745 |
| | 'Non-pill users' | Attractiveness | | 0.38 | (1,26) 4.38 | 0.046 * | | 0.36 | (1,26) 3.96 | 0.057 |
| | | Masculinity | 0.85 | -0.10 | (1,26) 0.25 | 0.618 | 0.86 | -0.23 | (1,26) 1.45 | 0.239 |
| | | | | | | | | | | |
| | | | | | | | | | | |
| Face | Total | Attractiveness ST ^a | | -0.42 | (1,45) 9.59 | 0.003 ** | | -0.33 | (1,45) 5.51 | 0.023 * |
| | | Attractiveness LT ^b | | -0.43 | (1,45) 10.49 | 0.002 ** | | -0.35 | (1,45) 6.48 | 0.014 * |
| | | Symmetry perceived | 0.76 * | -0.41 | (1,45) 9.17 | 0.004 ** | 0.86 | -0.25 | (1,45) 3.07 | 0.086 |
| | | Asymmetry measured | | 0.29 | (1,45) 4.05 | 0.050 | | 0.20 | (1,45) 1.78 | 0.188 |
| | | Masculinity | | -0.20 | (1,45) 1.79 | 0.187 | | -0.02 | (1,45) 0.02 | 0.903 |
| | 'Pill users' | Attractiveness ST | | -0.41 | (1,45) 8.86 | 0.005 ** | | -0.30 | (1,45) 4.61 | 0.037 * |
| | | Attractiveness LT | | -0.41 | (1,45) 9.26 | 0.004 ** | | -0.38 | (1,45) 7.66 | 0.008 ** |
| | | Symmetry perceived | 0.79 * | -0.40 | (1,45) 8.45 | 0.006 ** | 0.81 | -0.25 | (1,45) 3.00 | 0.090 |
| | | Masculinity | | -0.21 | (1,45) 2.17 | 0.148 | | -0.01 | (1,45) 0.01 | 0.921 |
| | | | | | | | | | | |
| | 'Non-pill users' | Attractiveness ST | | -0.41 | (1,45) 8.83 | 0.005 ** | | -0.34 | (1,45) 5.94 | 0.019 * |
| | | Attractiveness LT | | -0.44 | (1,45) 10.89 | 0.002 ** | | -0.32 | (1,45) 5.16 | 0.028 * |
| | | Symmetry perceived | 0.78 * | -0.38 | (1,45) 7.51 | 0.009 ** | 0.88 | -0.23 | (1,45) 2.47 | 0.123 |
| | | Masculinity | | -0.19 | (1,45) 1.71 | 0.198 | | -0.05 | (1,45) 0.11 | 0.737 |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |

^a ST: for a Short-Term partner; ^b LT: for a Long-Term partner