

The Journal of Agricultural Science

<http://journals.cambridge.org/AGS>

Additional services for *The Journal of Agricultural Science*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Application of genetics and genomics to aquaculture development: current and future directions

B. McANDREW and J. NAPIER

The Journal of Agricultural Science / Volume 149 / Supplement S1 / February 2011, pp 143 - 151

DOI: 10.1017/S0021859610001152, Published online: 22 December 2010

Link to this article: http://journals.cambridge.org/abstract_S0021859610001152

How to cite this article:

B. McANDREW and J. NAPIER (2011). Application of genetics and genomics to aquaculture development: current and future directions. The Journal of Agricultural Science, 149, pp 143-151
doi:10.1017/S0021859610001152

Request Permissions : [Click here](#)

FORESIGHT PROJECT ON GLOBAL FOOD AND FARMING FUTURES

Application of genetics and genomics to aquaculture development: current and future directions

B. McANDREW^{1*} AND J. NAPIER²

¹ Genetics and Reproduction Research Group, Institute of Aquaculture, University of Stirling,
Stirling FK9 4LA, UK

² Department of Biological Chemistry, Rothamsted Research, Harpenden, Herts AL5 2JQ, UK

(Revised MS received 6 October 2010; Accepted 6 October 2010; First published online 22 December 2010)

SUMMARY

Global aquaculture production continues to grow rapidly yet a small proportion of the animals and plants being used come from managed breeding and improvement programmes. The biology of aquatic organisms offer many opportunities for rapid genetic gains as new genetic and genomic techniques make the management of improvement programmes feasible in a wider range of species. The current paper describes the application of a wide range of techniques, many unique to aquatic organisms, and their potential to secure aquaculture production in the future.

INTRODUCTION

Aquaculture production has grown from less than 1 million tonnes (t) in the 1950s to nearly 60 million t in 2008 (FAO 2009). However, production of many species still depends on wild-caught fry or broodstock: 0.86 of the 69 farmed Chinese freshwater species (Honglang 2007) come from wild-caught fish. For domesticated stocks, there is growing evidence that traditional breeding approaches seriously degrade the genetic quality without continuous replacement by wild or managed fish (Mair *et al.* 2007). Less than 0.05 of production was estimated to come from scientifically managed breeding programmes (Gjedrem 2005).

The history of fish genetics has been reviewed by Dunham (2004) and Gjedrem (2005). A range of attributes has been targeted, including growth, maturation, environmental tolerance and disease resistance. Advantages for genetic gain include external fertilization and high fecundity, allowing a large number of gametes to be collected and fertilized under controlled conditions. Breeding designs with

large family sizes can be constructed, allowing better estimates of genetic parameters and high-selection pressures, and improvements can be passed on quickly to industry. Many species are still close to being wild organisms and display high levels of genetic variation; heritabilities are medium to high across a wide range of traits. Aquatic organisms are susceptible to environmental manipulation and a single genotype can potentially have many phenotypes. Single-sex populations can dramatically boost production. The phenotypic sex of many species can be changed by administering sex hormones during the sexually labile period of development, single-sex stock being generated directly or through controlled breeding of sex-reversed individuals. Other major traits such as the onset of sexual maturity can be advanced or delayed by the manipulation of daylength.

The new ‘-omic’ (e.g. genomic and proteomic) tools have also been applied in a wide range of species, though with few practical applications as yet. More widely in the development of productive and cost-effective aquaculture, genetic applications for developing high-quality feed inputs have great significance, progress in which is described in the final section of the present review.

* To whom all correspondence should be addressed.
Email: b.j.mcandrew@stir.ac.uk

SELECTIVE BREEDING

The species that have seen the most improvement are common carp, rainbow trout, Atlantic salmon and tilapia, channel catfish and ornamental fishes. Modern programmes for carp were established in Central and Eastern Europe (Kirpichnikov 1981), and improved strains contribute to the bulk of European and Asian production, with programmes to further develop these in Vietnam and India (Penman *et al.* 2005). Recent studies have shown that cross breeding and pedigree-based selection can still produce improvements (Vandeputte 2009).

Rainbow trout has also had a long history of domestication, in the USA (Gall & Huang 1988), Norway (selected for seawater production) (Gjedrem 1992), Finland (Kause *et al.* 2003) and Denmark (Henryon *et al.* 2002). But the uptake of selected strains by the industry has been slow; for instance, the largest commercial US egg supplier, Trout Lodge, only started a family-based breeding programme within the last 5 years. Short-term improvements such as all-female production or all-female triploids have been adopted but most eggs still come from mass-selected domesticated strains.

Norwegian Atlantic salmon programmes were initially government funded, but are now run by private companies (AquaGen and Salmobreed; Gjedrem 2005). Individual and family performance is assessed for a range of commercial traits (growth, sexual maturation and body conformation) and, more recently, disease resistance (infectious pancreatic necrosis virus (IPNV), *Aeromonas*, infectious salmon anaemia (ISA)). Estimated gains are between 8 and 10% per generation (Gjoen & Bentsen 1997). Similar programmes have also been developed by Landcatch Natural Selection in Scotland and Chile, Stofnfiskur in Iceland and Aquachile in Chile. Most farmed eggs now come from scientific breeding programmes, with significantly improved performance. Genomic technologies, including quantitative trait loci (QTL) and marker-assisted selection (MAS) approaches, are already benefiting improved virus resistance (Houston *et al.* 2008; Moen *et al.* 2009).

Tilapia have become the second most important group of farmed fish in the world (>2.7 million t in 2008) with *Oreochromis niloticus* becoming the dominant species in fresh water. Early selection work was hindered by the degraded nature of many of the commercial strains being used. The Genetically Improved Farm Tilapia (GIFT) programme, starting in 1987 in the Philippines, systematically compared wild and commercial strains in a variety of aquaculture environments and began family-based selection for growth. The work was further developed by Worldfish (CGIAR, see Acosta & Gupta 2010). Other programmes have either used the GIFT strain as a starting point (Genomar 2008) or have constructed

new synthetic populations using a number of selected lines (Rutten *et al.* 2005). The bulk of tilapia production, however, does not come from improved sources and producers use direct hormone sex-reversal of mixed sex fry to produce phenotypic all-male populations to overcome problems of excessive fry production during growout (see Beveridge & McAndrew 2000).

Selective breeding programmes have been more recently applied to other marine high-value species: European sea bass, Gilthead sea bream and Turbot. By 2005 it was assumed that as much as 0.80 of the European sea bass production already came from commercial populations which had undergone some level of genetic improvement and at least one major Greek company has initiated (in 2002) a large-scale family-based selective breeding programme on sea bream. Programmes on other high-value species, such as cod, halibut and tuna, are also under way.

HYBRIDIZATION

The ease of gamete collection in fish means that there are many references to inter-specific and inter-generic hybridization and the potential for using hybrids in aquaculture (Bartley *et al.* 2001). Hybrids have become important at the national level in several countries, e.g. *O. niloticus* × *Oreochromis aureus* in Israel because of the skewed male sex ratio and cold tolerance of the hybrid (Hulata *et al.* 1993). In the USA, the sunshine bass (*Morone chrysops* × *Morone saxatilis*) grows faster and has better overall culture characteristics than either parental species under commercial culture conditions (Smith 1988). However, despite the large numbers of reported hybrids few have been successfully cultured for extended periods because of the added complexity of production or because of introgression of the hybrid back into the parental species and loss of beneficial characteristics.

CHROMOSOME SET MANIPULATION

A range of genetic and environmental manipulations can be applied to aquatic species. If normally fertilized eggs are heat or pressure shocked at the 2nd meiotic division, a triploid (3n) embryo, containing three chromosome sets, is produced. Protocols are now available for over 30 different fish and shellfish species (Dunham 2004). Triploids are effectively sterile and are used in the production of larger rainbow trout, usually from all-female strains (Bye & Lincoln 1986), channel catfish (Wolters *et al.* 1991), common carp (Basavaraju *et al.* 2002) and in other species whose maturation slows growth. They are also used to improve growth rate and flesh quality in oysters (Guo *et al.* 1996). Triploids can be used to reduce the risk of stocked or escaped farmed strains interbreeding with native populations (Kozfkay *et al.* 2006) or exotic

species, such as grass carp for aquatic weed control, becoming established (Wattendorf 1986). Triploid sterility is also proposed to stop impacts from escaped transgenic fish (Mair *et al.* 2007). A shock at first mitotic division will produce a tetraploid embryo with four sets of chromosomes (4n). Male 4ns produce diploid sperm and can be crossed with normal fish to produce 3n directly avoiding having to shock large numbers of eggs, e.g. for rainbow trout (Myers *et al.* 1986).

Techniques for parthenogenesis can derive offspring from wholly maternal or paternal origins (Komen & Thorgaard 2007). These use gamma or ultraviolet radiation to destroy the nuclear DNA in eggs or sperm; the treated gamete is then fused with an untreated sperm or egg to produce a haploid embryo, which can be made diploid by inhibiting the second meiotic (meiotic gynogenetic) or first mitotic division (mitotic gynogenetic or androgenetic). Although still mainly used as research tools they can generate unique genotypes for analysing complex traits or generating new strains. Containing sufficient DNA, haploid embryos can be used for gene-mapping (Kocher *et al.* 1998), generating unusual genotypes such as YY male tilapia broodstock used to produce all-male XY tilapia offspring by crossing with normal XX females (Myers *et al.* 1986). Meiotic gynogenetic silver barb (*Puntius gonionotus*) are all-female, can be produced in large numbers and can be sex-reversed to generate neo-males without the need for progeny testing; they are now used in large numbers as broodstock for commercial production of all-female fry (Pongthana *et al.* 1999). Gynogenesis was similarly used in the development of all-female turbot (Cal *et al.* 2006).

Double haploid individuals from either a mitotic gynogenetic or androgenetic background can also be used to generate clonal or isogenic lines after a second round of parthenogenesis. Apart from model species such as zebrafish (Streisinger *et al.* 1981), clonal lines have been produced in tilapia (Müller-Belecke & Hörstegen Schwark 1995; Hussain *et al.* 1998; Sarder *et al.* 1999) rainbow trout (Scheerer *et al.* 1986; Thorgaard *et al.* 1990; Quillet *et al.* 2007) and common carp (Bongers *et al.* 1997). These are useful for disease and vaccine studies and an important resource for whole genome sequencing.

GENE TRANSFER TECHNOLOGIES

Transgenic fish containing the human growth hormone gene (hGHg) have been produced in goldfish (Zhu *et al.* 1985), rainbow trout (Penman *et al.* 1990), channel catfish (Dunham *et al.* 1987) and Nile tilapia (Brem *et al.* 1988). However, less than 0.05 integrated a copy of the construct into the host genome, and a number of other problems arose. Improved techniques resulted in growth enhancement in rainbow trout

(Penman *et al.* 1991), common carp (Zhang *et al.* 1990), channel catfish (Dunham *et al.* 1992) and tilapia (Rahman & Maclean 1999). The most dramatic results were with total piscine constructs using either an ocean pout antifreeze promoter (ocAFP) controlling a chinook salmon growth hormone (GH) cDNA or a sockeye salmon metallothionein (MT) promoter controlling a full-length sockeye GH gene. Transgenics showed a 5–30-fold increase in growth up to 1 year of age (Devlin *et al.* 1995) and individuals successfully passed this performance onto their offspring. The same construct had similar effects when used in other salmonid species (Du *et al.* 1992; Devlin *et al.* 1995; Devlin 1997; Cook *et al.* 2000).

Salmonids have a very positive response to GH (Dunham & Devlin 1998), while warm water species such as tilapia and carp, growing rapidly throughout the year, are probably not as reliant on GH regulation. Effects are also greater on wild fish than selected strains; Devlin *et al.* (2001) found a 17-fold improvement of transgenics over their wild sibs in rainbow trout, but only 4.4% gain with selected domestic strains.

Quite apart from the difficult issue of public acceptability and environmental risk, a range of practical constraints means that transgenic fish are unlikely to become a commercial reality in the immediate future (Mair *et al.* 2007). However, functional genomic work is already identifying candidate transgenic genes in the area of improved disease resistance.

MOLECULAR TECHNIQUES

Genetic markers

Liu & Cordes (2004) and Liu (2009) provide reviews on molecular markers in aquatic organisms. Markers are defined as Type 1 or actual genes of known function and Type 2 or anonymous DNA segments. To date, Type 2 and in particular microsatellite markers (Tautz 1989) have had a notable impact and large numbers of microsatellite loci (small tandem repeat DNA sequences) have been generated in salmonids, tilapia, sea bass, sea bream and carp.

Microsatellites have immediate application in defining stock parentage (Norris *et al.* 2000), particularly for marine species where it is difficult to develop single family rearing. Studies in mass spawning species such as sea bream (Brown *et al.* 2005) and cod (Herlin *et al.* 2008) and in manually spawned halibut (Jackson *et al.* 2003) have shown that parentage assignment is critical to avoid the offspring of a few individuals dominating replacement broodstocks. This technology also removes the need for physical tags, increasing potential family numbers in commercial environments and avoiding the risk of tags entering the human food chain, and can also reduce the cost of maintaining breeding nucleus backup populations.

Genetic markers have many other potential uses in managing farmed and wild stocks, for example identifying origin of escaped farm stock in Atlantic salmon, rainbow trout and Atlantic cod (Glover 2010) and defining potential introgression of genes from farm escapes into wild populations of salmon (Skaala *et al.* 2006).

Genetic mapping

Genetic linkage maps (Danzmann & Gharbi 2007) enable identification of QTL or genome sections containing genes influencing important traits. The QTL data for species such as Atlantic salmon, tilapia and common carp are steadily increasing (Korol *et al.* 2007), though most maps require further precision. However, using the large differences in recombination rate between sexes (Hayes *et al.* 2006), it has been possible to speed up QTL identification for resistance to infectious pancreatic necrosis (IPN) and incorporate the QTL into a commercial Atlantic salmon breeding programme (Houston *et al.* 2008).

The majority of QTL described are for easily measured traits. Future programmes will benefit by identifying QTL or more focused MAS for traits that are more difficult to measure, e.g. those with low heritability, identifying candidates early before maturity, traits only observed in one of the sexes and a range of post-harvest traits such as flesh quality (e.g. colour, fillet conformation and adiposity; Lande & Thompson 1990). The use of MAS technology will increase in aquaculture, and De Santis & Jerry (2007) have listed a number of candidate genes derived from what is known in terrestrial livestock.

Rapid developments in identifying expressed sequence tags (EST) are opening prospects for mapping Type 1 markers. For stock in different developmental stages or environments it will be possible to see which genes are expressed and how a given gene or subset of genes is affected. Mapping of ESTs is easier if a radiation hybrid (RH) mapping panel is available. The first panel was constructed in a commercial species, gilthead sea bream *Sparus auratus* (Senger *et al.* 2006), and work is in progress for species such as sea bass. Type 1 markers allow comparative use of genomic approaches to help identify possible loci for less well-studied commercial species Sarropoulou *et al.* (2008).

This area is developing very rapidly (Rexroad 2007), with work on single nucleotide polymorphisms (SNP) (Liu 2009) and bacterial artificial chromosome (BAC) libraries to more densely map species (He *et al.* 2007). Several BACs are now available: for Atlantic salmon (Davidson, 2007), tilapia (Katagiri *et al.* 2001), sea bass (Whitaker *et al.* 2006) and Pacific oyster (*Crassostrea gigas*; Cunningham *et al.* 2006). Physical mapping projects based on BAC are under way in Atlantic salmon (Ng *et al.* 2005), tilapia

(Katagiri *et al.* 2005) and Channel catfish (Xu *et al.* 2007). New generation sequencing (NGS), particularly restriction site-associated DNA (RAD tagging; Miller *et al.* 2007) offers the opportunity to rapidly and cost-effectively identify and analyse thousands of SNPs and should speed up the discovery rate of QTL, particularly in species with poor genomic resources.

Functional genomics

The construction of microarray chips, containing thousands of ESTs derived from the whole animal or more focused subsets of genes specific to a tissue or a biological function, allows more specific assessment of characteristics. Work in salmonids has progressed the furthest (Rise *et al.* 2007), studying gene expression for a range of traits including disease-related responses, such as for *Piscirickettsia salmonis* (Rise *et al.* 2004), *Aeromonas salmonicida* (Ewart *et al.* 2005), Ameobic gill disease (Morrison *et al.* 2006), immune response to a lipopolysaccharide challenge (MacKenzie *et al.* 2006), live bacterial vaccines (Martin *et al.* 2006), DNA vaccination (Purcell *et al.* 2006) and *Gyrodactylus* species (Fast *et al.* 2006; Lindenstrom *et al.* 2006). Other traits include response to growth in transgenic salmon (Rise *et al.* 2007) and stress associated with handling (Krasnov *et al.* 2005), temperature (Vornanen *et al.* 2005) and highly unsaturated fatty acid lipid metabolism (Taggart *et al.* 2008).

With large numbers of genes being monitored for expression under a range of different conditions, it is likely that integrating QTL mapping with global gene expression may well identify patterns correlating with differences in key traits (Haley & de Koning 2006).

THE POTENTIAL FOR ENGINEERING PLANT SUBSTRATES FOR OMEGA-3 STOCKS IN FISH FEEDS

A further area of genetic exploration in aquaculture is that of providing feeding sources, particularly to replace or supplement high-quality inputs currently derived from fishmeal and oil, increasingly seen as a limitation for future growth in aquaculture production. Sourcing of key components, particularly omega-3 fatty acids in plant substrates, could offer great advantages of supply and cost. The reproductive tissues of higher plants store significant amounts of neutral lipids (predominantly as triacylglycerols), providing a convenient and renewable source of useful fatty acids. Over 400 different fatty acids have been identified in seed oils although, remarkably, none have been found to contain the very long chain omega-3 polyunsaturated fatty acids (LC-PUFAs), particularly eicosapentanoic acid (EPA) and docosa-hexanoic acid (DHA).

Thus, there exists the possibility of genetically engineering plants to modify seed oil composition to include omega-3 PUFAs. Its feasibility has already been demonstrated with the 'stacking' of multiple genes in a single transgenic plant, and also the identification from marine microbes of the biosynthetic genes which direct the synthesis of EPA and DHA. As outlined below, considerable progress has been made in the last 5 years towards producing an alternative, sustainable source of fish oils from transgenic plants.

The first landmark 'proof of concept' demonstration of the accumulation of the C20 omega-3 LC-PUFA EPA in transgenic plants was achieved in 2004 by several groups, using genes from PUFA-synthesizing micro-organisms to direct the synthesis of this fatty acid in either leaves or seeds of different plant species (*Arabidopsis*, Qi *et al.* 2004; tobacco and linseed, Abbadi *et al.* 2004). Both studies reported low but significant levels of EPA (0.01–0.03 of total fatty acids). Building on these foundations, further work was carried out in *Brassica juncea* (Indian mustard) in which up to nine algal and moss genes were expressed, resulting in the accumulation of 0.15 EPA but <0.01 DHA in seed oils (Wu *et al.* 2005). Similar attempts to produce DHA in transgenic *Arabidopsis* seeds resulted in <0.01 DHA in total fatty acids (Robert *et al.* 2005), whereas further fine-tuning resulted in accumulation of up to 0.25 EPAs in *Brassica carinata* (Cheng *et al.* 2010). However, while transgenic accumulation of the C20 omega-3 PUFA EPA is achievable at levels similar or greater to those found in marine organisms (fish, algae or diatoms), the synthesis of DHA is still a considerable challenge, with no current demonstration of accumulation above 0.03–0.04 (Venegas-Calerón *et al.* 2010). However, since DHA is a direct metabolite of EPA, it might be expected that the recently reported high levels of EPA will serve as a superior platform with which to increase transgenic DHA levels.

In addition to EPA and DHA, efforts have been focused at transgenic production of the omega-3 surrogate stearidonic acid (SDA): SDA is not a *bona fide* omega-3 LC-PUFA, being only 18 carbons long, but has been demonstrated to undergo *in vivo* conversion to EPA in animals and could serve as an alternative source of fish oils. Only a very few plant species accumulate SDA, such as *Echium*, a non-agronomically adapted and low-yielding species. Transgenic soybeans accumulating significant SDA (up to 0.29 of total seed fatty acids) have been reported (Eckert *et al.* 2006), and has transgenic linseed enriched with up to 0.13 SDA (Ruiz-Lopez *et al.* 2009). Interestingly, although SDA levels in transgenic linseed were lower than that reported for soybean, the C18 omega-3 content was superior, as was the absence of the omega-6 fatty acid dihomo-gamma-linoleic acid (GLA; Ruiz-Lopez *et al.* 2009).

Stearidonic acid has been reported to be effective as a fish oil replacement in aquaculture, though this has been disputed.

Collectively, these data confirm the promise of transgenic plants as sources of omega-3 LC-PUFAs. However, while levels of EPA (and SDA) achieved are equivalent to marine sources, DHA still represents a challenge (most likely due to its more complicated biosynthetic pathway and possibly also its reduced oxidative stability). However, given recent progress in elevating EPA it seems likely that higher DHA levels will be achievable, most likely some 0.05–0.10 of total seed oil. Such levels, especially in conjunction with moderate (0.10–0.15) levels of EPA would represent a very significant and useful source of omega-3 LC-PUFAs for aquaculture. Alternatively, an EPA-containing plant oil (lacking DHA) could provide a convenient feedstock, particularly as such material has already been produced. It will also be important to evaluate the performance of different farmed fish species when provided with these novel sources of omega-3 LC-PUFAs.

For applications in aquaculture, several factors need to be highlighted. Firstly, these oils should be considered as 'enhanced' with omega-3 LC-PUFAs, rather than a direct replacement, as plant oils contain significant oleic, linoleic and linolenic acids not usually found in marine oils. Thus, they might require blending to ensure optimal composition. Cultivation and use of genetically modified (GM)-derived products are also widely regulated and positive consumer responses would also need to be assured. Another consideration is the volume of GM crops to be grown – aquaculture demands could require considerable area just to provide sufficient omega-3 LC-PUFA oils to substitute for the many thousands of tonnes of fish oils consumed in the UK.

CONCLUSIONS

The current review has outlined genetic applications related to aquaculture and associated activities, ranging from immediately practical, commercially developed approaches already delivering tangible improvements to emerging techniques which will inform potential for a range of stock, environmental and husbandry interactions. There is substantial scope for using existing practical techniques; widespread adoption of broodstock management and improvement programmes would have significant impacts on sector performance, and routine application of genetic principles to hatchery-based fishery stocking programmes, and to defining and managing the aquaculture impact on biodiversity would bring about important resource and environmental gains. With a mix of strategies based on the more experimental techniques, and with more quantitative rigour the

costs and benefits to the sector and to the wider environment and resource base can be better defined.

While the application of GM technologies to stocks themselves remains highly controversial, other areas may have more immediate potential. The possibility of using transgenic plants to synthesize omega-3

LC-PUFAs is within sight of providing a terrestrial feedstock unconstrained by current concerns for marine sources (Napier & Graham 2010). With suitable controls and consumer trust, genetically modified oilseed crops could become more widely grown, providing further potential for a wide range of aquaculture production.

REFERENCES

- ABBADI, A., DOMERGUE, F., BAUER, J., NAPIER, J. A., WELTI, R., ZÄHRINGER, U., CIRPUS, P. & HEINZ, E. (2004). Biosynthesis of very-long-chain polyunsaturated fatty acids in transgenic oilseeds: constraints on their accumulation. *Plant Cell* **16**, 2734–2748.
- ACOSTA, B. O. & GUPTA, M. V. (2010). The genetic improvement of farmed tilapias project: impact and lessons learned. In *Success Stories in Asian Aquaculture* (Eds S. S. De Silva & F. B. Davy), pp. 149–171. Dordrecht, The Netherlands: Springer.
- BARTLEY, D. M., RANA, K. & IMMINK, A. J. (2001). The use of inter-specific hybrids in aquaculture and fisheries. *Reviews in Fish Biology and Fisheries* **10**, 325–337.
- BASAVARAJU, Y., MAIR, G. C., MOHAN KUMAR, H. M., PRADEEP KUMAR, S., KESHAVAPPA, G. Y. & PENMAN, D. J. (2002). An evaluation of triploidy as a potential solution to the problem of precocious sexual maturation in common carp *Cyprinus carpio*, in Karnataka, India. *Aquaculture* **204**, 407–418.
- BEVERIDGE, M. C. M. & McANDREW, B. J. (2000). *Tilapia: Biology and Exploitation*. Dordrecht, The Netherlands: Kluwer Academic Publishers.
- BONGERS, A. B. J., BOVENHUIS, H., VAN STOKKOM, A. C., WIEGERTJES, G. F., ZANDIEH-DOULABI, B., KOMEN, J. & RICHTER, C. J. J. (1997). Distribution of genetic variance in gynogenetic or androgenetic families. *Aquaculture* **153**, 225–238.
- BREM, G., BRENIG, B., HORSTGEN-SCHWARK, G. & WINNACKER, E. L. (1988). Gene transfer in tilapia (*Oreochromis niloticus*). *Aquaculture* **68**, 209–219.
- BROWN, R. C., WOOLLIAMS, J. A. & McANDREW, B. J. (2005). Factors influencing effective population size in commercial populations of gilthead seabream, *Sparus aurata*. *Aquaculture* **247**, 219–225.
- BYE, V. J. & LINCOLN, R. F. (1986). Commercial methods for the control of sexual maturation in rainbow trout (*Salmo gairdneri* R.). *Aquaculture* **57**, 299–309.
- CAL, R. M., VIDAL, S., MARTINEZ, P., ALVAREZ-BLAZQUEZ, B., GOMEZ, C. & PIFERRER, F. (2006). Growth and gonadal development of gynogenetic diploid *Scophthalmus maximus*. *Journal of Fish Biology*, **68**, 401–413.
- CHENG, B., WU, G., VRINTEN, P., FALK, K., BAUER, J. & QIU, X. (2010). Towards the production of high levels of eicosapentaenoic acid in transgenic plants: the effects of different host species, genes and promoters. *Transgenic Research* **19**, 221–229.
- COOK, J. T., McNIVEN, M. A., RICHARDSON, G. F. & SUTTERLIN, A. M. (2000). Growth rate, body composition and feed digestibility/conversion of growth enhanced transgenic Atlantic salmon (*Salmo salar*). *Aquaculture* **188**, 15–32.
- CUNNINGHAM, C., HIKIMA, J. I., JENNY, M. J., CHAPMAN, R. W., FANG, G. C., SASKI, C., LUNDQVIST, M. L., WING, R. A., CUPIT, P. M., GROSS, P. S., WARR, G. W. & TOMKINS, J. P. (2006). New resources for marine genomics: Bacterial artificial chromosome libraries for the eastern and pacific oysters (*Crassostrea virginica* and *C. gigas*). *Marine Biotechnology* **8**, 521–533.
- DANZMANN, R. G. & GHARBI, K. (2007). Linkage mapping in aquaculture species. In *Aquaculture Genome Technologies* (Ed. Z. J. Liu), pp. 139–167. Oxford, UK: Blackwell Publishing.
- DAVIDSON, W. S. (2007). Bacterial Artificial Chromosome Libraries and BAC based physical mapping of aquaculture genomes. In *Aquaculture Genome Technologies* (Ed. Z. J. Liu), pp. 245–259. Oxford, UK: Blackwell Publishing.
- DE-SANTIS, C. & JERRY, D. R. (2007). Candidate growth genes in finfish – where should we be looking? *Aquaculture* **272**, 22–38.
- DEVLIN, R. H. (1997). Transgenic salmonids. In *Transgenic Animals: Generation and Use* (Ed. L. M. Houdebine), pp. 105–117. Amsterdam, The Netherlands Harwood Academic Publishers.
- DEVLIN, R. H., BIAGI, C. A., YESAKI, T. Y., SMAILUS, D. E. & BYATT, J. C. (2001). Growth of domesticated transgenic fish. *Nature* **409**, 781–782.
- DEVLIN, R. H., YESAKI, T. Y., DONALDSON, E. M., DU, S.-J. & HEW, C. L. (1995). Production of germline transgenic Pacific salmonids with dramatically increased growth performance. *Canadian Journal of Fisheries and Aquatic Sciences* **52**, 1376–1384.
- DU, S. J., GONG, Z., FLETCHER, G. L., SHEARS, M. A., KING, M. J., IDLER, D. R. & HEW, C. L. (1992). Growth enhancement in transgenic Atlantic salmon by the use of an ‘all fish’ chimeric growth hormone gene construct. *Nature Biotechnology* **10**, 176–181.
- DUNHAM, R. A. (2004). *Aquaculture and Fisheries Biotechnology: Genetic Approaches*. Wallingford, UK: CABI.
- DUNHAM, R. A. & DEVLIN, R. (1998). Comparison of traditional breeding and transgenesis in farmed fish with implications for growth enhancement and fitness. In *Transgenic Animals in Agriculture* (Ed. J. D. Murray, G. B. Anderson, A. M. Oberbauer & M. M. McGloughlin), pp. 209–229. Wallingford UK: CABI.
- DUNHAM, R. A., EASH, J., ASKINS, J. & TOWNES, T. M. (1987). Transfer of metallothionein-human growth hormone fusion gene into channel catfish. *Transactions of the American Fisheries Society* **116**, 87–91.
- DUNHAM, R. A., RAMBOUX, A. C., DUNCAN, P. L., HAYAT, M., CHEN, T. T., LIN, C. M.

- GONZALEZ-VILLASENOR, L. I. & POWERS, D. A. (1992). Transfer, expression and inheritance of salmonid growth hormone in channel catfish, *Ictalurus punctatus*, and effects on performance traits. *Molecular Marine Biology and Biotechnology* **1**, 380–389.
- ECKERT, H., LA VALLEE, B., SCHWEIGER, B. J., KINNEY, A. J., CAHOON, E. B. & CLEMENTE, T. (2006). Co-expression of the borage Delta 6 desaturase and the Arabidopsis Delta 15 desaturase results in high accumulation of stearidonic acid in the seeds of transgenic soybean. *Planta* **224**, 1050–1057.
- EWART, K. V., BELANGER, J. C., WILLIAMS, J., KARAKACH, T., PENNY, S., TSOI, S. C. M., RICHARDS, R. C. & DOUGLAS, S. E. (2005). Identification of genes differentially expressed in Atlantic salmon (*Salmo salar*) in response to infection by *Aeromonas salmonicida* using cDNA microarray technology. *Developmental and Comparative Immunology* **29**, 333–347.
- FAO (2009). Fisheries Statistics – Aquaculture Production. Available online at <http://www.fao.org/fi/statist/FISOFT/FISHPLUS.asp> (verified 15 October 2010).
- FAST, M. D., ROSS, N. W., MUISE, D. M. & JOHNSON, S. C. (2006). Differential Gene Expression in Atlantic Salmon infected with *Lepeophtheirus salmonis*. *Journal of Aquatic Animal Health* **18**, 116–127.
- GALL, G. A. E. & HUANG, N. (1988). Heritability and selection schemes for rainbow trout: Female reproductive performance. *Aquaculture* **73**, 57–66.
- GENOMAR. (2008). Trapia: Breeding Nucleus. Available online at: <http://www.genomar.no/?did=9078117> (verified 15 October 2010).
- GJEDREM, T. (1992). Breeding plans for rainbow trout. *Aquaculture* **100**, 73–83.
- GJEDREM, T. (2005). Selection and Breeding Programs in Aquaculture. Berlin: Springer. ISBN-10 1-4020-3341-9. 364p.
- GJOEN, H. M. & BENTSEN, H. B. (1997). Past, present and future of genetic improvement in salmon aquaculture. *ICES Journal of Marine Science* **54**, 1009–1014.
- GLOVER, K. A. (2010). Forensic identification of fish farm escapees: the Norwegian experience. *Aquaculture Environment Interactions* **1**, 1–10.
- GUO, X., DEBROSSE, G. A. & ALLEN, S. K. JR. (1996). All triploid Pacific oysters (*Crassostrea gigas* Thunberg) produced by mating tetraploids and diploids. *Aquaculture* **142**, 149–161.
- HALEY, C. & DE KONING, D. J. (2006). Genetical genomics in livestock: potentials and pitfalls. *Animal Genetics* **37**, 10–12.
- HAYES, B. J., GJUVSLAND, A. & OMHOLT, S. (2006). Power of QTL mapping experiments in commercial Atlantic salmon populations, exploiting linkage and linkage disequilibrium and effect of limited recombination in males. *Heredity* **97**, 19–26.
- HE, L., DU, C., LI, Y., SCHEURING, C. & ZHANG, H.-B. (2007). Construction of large-insert Bacterial Clone Libraries and their applications. In *Aquaculture Genome Technologies* (Ed. L. J. Liu), pp. 215–244. Oxford, UK: Blackwell Publishing.
- HENRYON, M., JOKUMSEN, A., BERG, P., LUND, I., PEDERSEN, P. B., OLESEN, N. J. & SLIERENDRECHT, W. J. (2002). Genetic variation for growth rate, feed conversion efficiency, and disease resistance exists within a farmed population of rainbow trout. *Aquaculture* **209**, 59–76.
- HERLIN, M., DELGHANDI, M., WESMAJERVI, M., TAGGART, J. B., MCANDREW, B. J. & PENMAN, D. J. (2008). Analysis of the parental contribution to a group of fry from a single day of spawning from a commercial Atlantic cod (*Gadus morhua*) breeding tank. *Aquaculture* **274**, 218–224.
- HONGLANG, H. (2007). Freshwater fish seed resources in China. In *Assessment of Freshwater Fish Seed Resources for Sustainable Aquaculture* (Ed. M. G. Bondad-Reantaso), pp. 185–199. FAO Fisheries Technical Paper No. 501. Rome: FAO.
- HOUSTON, R. D., HALEY, C. S., HAMILTON, A., GUY, D. R., TINCH, A. E., TAGGART, J. B., MCANDREW, B. J. & BISHOP, S. C. (2008). Major quantitative trait loci affect resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*). *Genetics* **178**, 1109–1115.
- HULATA, G., WOHLFARTH, G. W., KARPLUS, I., SCHROEDER, G. L., HARPAZ, S., HALEVY, A., ROTHBARD, S., COHEN, S., ISRAEL, I. & KAVESSA, M. (1993). Evaluation of *Oreochromis niloticus* × *O. aeneus* hybrid progeny of different geographical isolates, reared under varying management regimes. *Aquaculture* **115**, 253–271.
- HUSSAIN, M. G., PENMAN, D. J. & MCANDREW, B. J. (1998). Production of heterozygous and homozygous clones in Nile tilapia. *Aquaculture International* **6**, 197–205.
- JACKSON, T. R., MARTIN-ROBICHAUD, D. J. & REITH, M. E. (2003). Application of DNA markers to the management of Atlantic halibut (*Hippoglossus hippoglossus*) broodstock. *Aquaculture* **220**, 245–259.
- KATAGIRI, T., ASAKAWA, S., MINAGAWA, S., SHIMIXU, N., HIRONO, I. & AOKI, T. (2001). Construction and characterization of BAC libraries for three fish species; rainbow trout, carp and tilapia. *Animal Genetics* **32**, 200–204.
- KATAGIRI, T., KIDD, C., TOMASINO, E., DAVIS, J. T., WISHON, C., STERN, J. E., CALRETON, K. L., HOWE, A. E. & KOCHER, T. D. (2005). A BAC based physical map of the Nile tilapia genome *BMC Genomics* **6**, doi:10.1186/1471-2164-6-89.
- KAUSE, A., RITOLA, O., PAANANEN, T., MÄNTYSAARI, E. & ESKELINEN, U. (2003). Selection against early maturity in large rainbow trout *Oncorhynchus mykiss*: the quantitative genetics of sexual dimorphism and genotype-by-environment interactions. *Aquaculture* **228**, 53–68.
- KIRPICHNIKOV, V. S. (1981). *Genetic Bases of Fish Selection*. Berlin: Springer-Verlag.
- KOCHER, T. D., LEE, W.-J., SOBOLEWSKA, H., PENMAN, D. J. & MCANDREW, B. J. (1998). A genetic linkage map of a cichlid fish, the tilapia (*Oreochromis niloticus*). *Genetics* **148**, 1225–1232.
- KOMEN, H. & THORGAARD, G. H. (2007). Androgenesis, gynogenesis and the production of clones in fishes: a review. *Aquaculture* **269**, 150–173.
- KOROL, A., SHIRAK, A., CNAANI, A. & HALLERMAN, E. M. (2007). Detection and analysis of quantitative trait loci (QTL) for economic traits in Aquatic species. In *Aquaculture Genome Technologies* (Ed. L. J. Liu), pp. 169–187. Oxford, UK: Blackwell Publishing.
- KOZFKAY, J. R., DILLON, J. C. & SCHILL, D. J. (2006). Routine use of sterile fish in salmonid sport fisheries: are we there yet? *Fisheries* **31**, 392–401.
- KRASNOV, A., KOSKINEN, H., PEHKONEN, P., REXROAD, C. E. III, AFANASYEV, S. & MOLSA, H. (2005). Gene expression in the brain and kidney of rainbow trout in response to

- handling stress. *BMC Genomics* **6**, 3. doi: 10.1186/1471-2164-6-3.
- LANDE, R. & THOMPSON, R. (1990). Efficiency of marker assisted selection in improvement of quantitative traits. *Genetics* **124**, 743–756.
- LINDENSTROM, T., SIGH, J., DALGAARD, M. B. & BUCHMANN, K. (2006). Skin expression of IL-1 beta in East Atlantic salmon, *Salmo salar* L., highly susceptible to *Gyrodactylus salaris* infection is enhanced compared to a low susceptibility Baltic stock. *Journal of Fish Diseases* **29**, 123–128.
- LIU, Z. J. (2009). *Aquaculture Genome Technologies*. Oxford, UK: Blackwell Publishing.
- LIU, Z. J. & CORDES, J. (2004). DNA marker technology and their applications in aquaculture genetics. *Aquaculture* **238**, 1–37.
- MAIR, G. C., NAM, Y. K. & SOLAR, I. I. (2007). Risk management: Reducing risk through confinement of transgenic fish. In *Environmental Risk Assessment of Genetically Modified Organisms. Vol. 3. Methodologies for Transgenic Fish* (Eds A.R. Kapuscinski K. R. Hayes, S. Li & G. Dana), pp. 209–238. Wallingford, UK: CABI.
- MACKENZIE, S., ILIEV, D., LIARTE, C., KOSKINEN, H., PLANAS, J. V., GOETZ, F. W., MOLSA, H., KRASNOV, A. & TORT, L. (2006). Transcriptional analysis of LPS-stimulated activation of trout (*Oncorhynchus mykiss*) monocyte/macrophage cells in primary culture treated with cortisol. *Molecular Immunology* **43**, 1340–1348.
- MARTIN, S. A. M., BLANEY, S. C., HOULIHAN, D. F. & SECOMBES, C. J. (2006). Transcriptome response following administration of a live bacterial vaccine in Atlantic salmon (*Salmo salar*). *Molecular Immunology* **43**, 1900–1911.
- MILLER, M. R., DUNHAM, J. P., AMORES, A., CRESKO, W. A. & JOHNSON, E. A. (2007). Rapid and cost-effective polymorphism identification and genotyping using restriction site associated DNA (RAD) markers. *Genome Research* **17**, 240–248.
- MOEN, T., BARANSKI, M., SONESSON, A. K. & KJØGLUM, S. (2009). Confirmation and fine-mapping of a major QTL for resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*): population-level associations between markers and trait. *BMC Genomics* **10**, 368. doi:10.1186/1471-2164-10-368.
- MORRISON, R. N., COOPER, G. A., KOOP, B. F., RISE, M. L., BRIDLE, A. R., ADAMS, M. B. & NOWAK, B. F. (2006). Transcriptome profiling of the gills of amoebic gill disease (AGD)-affected Atlantic salmon (*Salmo salar*): a role for the tumor suppressor protein p52 in AGD – pathogenesis? *Physiological Genomics* **26**, 15–34.
- MÜLLER-BELECKE, A. & HÖRSTEGEN-SCHWARK, G. (1995). Sex determination in tilapia (*Oreochromis niloticus*) sex ratios in homozygous gynogenetic progeny and their offspring. *Aquaculture* **137**, 57–65.
- MYERS, J. M., HERSHBERGER, W. K. & IWAMOTO, R. N. (1986). The induction of tetraploidy in salmonids. *Journal of the World Aquaculture Society* **17**, 1–7.
- NAPIER, J. A. & GRAHAM, I. A. (2010). Tailoring plant lipid composition: designer oilseeds come of age. *Current Opinion in Plant Biology* **13**, 330–337.
- NG, S. H., ARTIERI, C. G., BOSDET, I. E., CHIU, R., DANZMANN, R. G., DAVIDSON, W. S., FERGUSON, M. M., FJELL, C. D., HOYHEIM, B., JONES, S. J., DE JONG, P. J., KOOP, B. F., KRZYWINSKI, M. I., LUBIENIECKI, K., MARRA, M. A., MITCHELL, L. A., MATHEWSON, C., OSEGAWA, K., PARISOTTO, S. E., PHILLIPS, R. B., RISE, M. L., VON SCHALBURG, K. R., SCHEIN, J. E., SHIN, H., SIDDIQUI, A., THORSEN, J., WYE, N., YANG, G. & ZHU, B. (2005). A physical map of the genome of Atlantic salmon (*Salmo salar*). *Genomics* **86**, 396–404.
- NORRIS, A. T., BRADLEY, D. G. & CUNNINGHAM, E. P. (2000). Parentage and relatedness determination in farmed Atlantic salmon (*Salmo salar*) using microsatellite markers. *Aquaculture* **182**, 73–83.
- PENMAN, D. J., BEECHING, A. J., PENN, S. & MACLEAN, N. (1990). Factors affecting survival and integration following microinjection of novel DNA into rainbow trout eggs. *Aquaculture* **85**, 35–50.
- PENMAN, D. J., BEECHING, A. J., PENN, S., RAHMAN, A., SULAIMAN, Z. & MACLEAN, N. (1991). Patterns of transgene inheritance in rainbow trout (*Oncorhynchus mykiss*). *Molecular Reproduction and Development* **30**, 201–206.
- PENMAN, D. J., GUPTA, M. V. & DEY, M. M. (2005). (eds). *Carp Genetic Resources for Aquaculture in Asia*. WorldFish Center Technical Report 65. Penang, Malaysia: WorldFish Center.
- PONGTHANA, N., PENMAN, D. J., BAOPRASERTKUL, P., HUSSAIN, M. G., ISLAM, M. S., POWELL, S. F. & McANDREW, B. J. (1999). Monosex female production in the silver barb (*Puntius gonionotus* Bleeker). *Aquaculture* **173**, 246–256.
- PURCELL, M. K., NICHOLS, K. M., WINTON, J. R., KURATH, G., THORGAARD, G. H., WHEELER, P., HANSEN, J. D., HERWIG, R. P. & PARK, L. K. (2006). Comprehensive gene expression profiling following DNA vaccination of rainbow trout against infectious hematopoietic necrosis virus. *Molecular Immunology* **43**, 2089–2106.
- QI, B., FRASER, T., MUGFORD, S., DOBSON, G., SAYANOVA, O., BUTLER, J., NAPIER, J. A., STOBART, A. K. & LAZARUS, C. M. (2004). Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants. *Nature Biotechnology* **22**, 739–745.
- QUILLET, E., DORSON, M., LE GUILLON, S., BENMANSOUR, A. & BOUDINOT, P. (2007). Wide range of susceptibility to rhabdoviruses in homozygous clones of rainbow trout. *Fish and Shellfish Immunology* **22**, 510–519.
- RAHMAN, M. A. & MACLEAN, N. (1999). Growth performance in transgenic tilapia containing an exogenous piscine growth hormone gene. *Aquaculture* **173**, 333–346.
- REXROAD, C. E. (2007). Radiation hybrid mapping in aquatic species. In *Aquaculture Genome Technologies* (Ed. Z. J. Liu), pp. 313–322. Oxford, UK: Blackwell Publishing.
- RISE, M. L., JONES, S. R., BROWN, G. D., VON SCHALBURG, K. R., DAVIDSON, W. S. & KOOP, B. F. (2004). Microarray analyses identify molecular biomarkers of Atlantic salmon macrophage and hematopoietic kidney response to *Piscirickettsia salmonis* infection. *Physiological Genomics* **20**, 21–35.
- RISE, M. L., VON SCHALBURG, K. R., COOPER, G. A. & KOOP, B. F. (2007). Salmonid DNA microarrays and other tools for functional genomics research In *Aquaculture Genome Technologies* (Ed. Z. J. Liu), pp. 369–411. Oxford, UK: Blackwell Publishing.
- ROBERT, S. S., SINGH, S. P., ZHOU, X.-R., PETRIE, J. R., BLACKBURN, S. I., MANSOUR, P. M., NICHOLS, P. D., LIU, Q. & GREEN, A. G. (2005). Metabolic engineering of

- Arabidopsis* to produce nutritionally important DHA in seed oil. *Functional Plant Biology* **32**, 473–479.
- RUIZ-LÓPEZ, N., HASLAM, R. P., VENEGAS-CALERÓN, M., LARSON, T. R., GRAHAM, I. A., NAPIER, J. A. & SAYANOVA, O. (2009). The synthesis and accumulation of stearidonic acid in transgenic plants: a novel source of 'heart-healthy' omega-3 fatty acids. *Plant Biotechnology Journal* **7**, 704–716.
- RUTTEN, M. J. M., KOMEN, H. & BOVENHUIS, H. (2005). Longitudinal genetic analysis of Nile tilapia (*Oreochromis niloticus* L.) body weight using a random regression model. *Aquaculture* **246**, 101–113.
- SARDER, M. R. I., PENMAN, D. J., MYERS, J. M. & MCANDREW, B. J. (1999). Production and propagation of fully inbred clonal lines in the Nile tilapia (*Oreochromis niloticus* L.). *Journal of Experimental Zoology* **284**, 675–685.
- SARROPOULOU, E., NOUSDILI, D., MAGOULAS, A. & KOTOULAS, G. (2008). Linking the genomes of nonmodel teleosts through comparative genomics. *Marine Biotechnology* **10**, 227–233.
- SCHEEER, P. D., THORGAARD, G. H., ALLENDORF, F. W. & KNUDSEN, K. L. (1986). Androgenetic rainbow trout produced from inbred and outbred sperm sources show similar survival. *Aquaculture* **57**, 289–298.
- SENGER, F., PRIAT, C., HITTE, C., SARROPOULOU, E., FRANCH, R., GEISLER, R., BARGELLONI, L., POWER, D. & GAILIBERT, F. (2006). The first radiation hybrid map of a perch like fish: the Gilthead seabream (*Sparus aurata* L.). *Genomics* **87**, 793–800.
- SKAALA, Ø., WENNEVIK, V. & GLOVER, K. A. (2006). Evidence of temporal genetic change in wild Atlantic salmon, *Salmo salar* L., populations affected by farm escapees. *ICES Journal of Marine Science* **63**, 1224–1233.
- SMITH, T. I. J. (1988). Aquaculture of striped bass and its hybrids in North America. *Aquaculture Magazine* **14**, 40–49.
- STREISINGER, G., WALKER, C., DOWER, N., KNAUBER, D. & SINGER, F. (1981). Production of clones of homozygous diploid zebra fish (*Brachydanio rerio*). *Nature* **291**, 293–296.
- TAGGART, J. B., BRON, J. E., MARTIN, S. A. M., SEEAR, P. J., HØYHEIM, B., TALBOT, R., CARMICHAEL, S. N., VILLENEUVE, L. A. N., SWEENEY, G. E., HOULIHAN, D. F., SECOMBES, C. J., TOCHER, D. R. & TEALE, A. J. (2008). A description of the origins, design and performance of the TRAITS-SGP Atlantic salmon *Salmo salar* L. cDNA microarray. *Journal of Fish Biology* **72**, 2071–2094.
- TAUTZ, D. (1989). Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Research* **17**, 6463–6471.
- THORGAARD, G. H., SCHEEER, P. D., HERSHBERGER, W. K. & MYERS, J. M. (1990). Androgenetic rainbow trout produced using sperm from tetraploid males show improved survival. *Aquaculture* **85**, 215–221.
- VANDEPUTTE, M. (2009). Genetic improvement of common carp (*Cyprinus carpio* L.). *Cahiers Agricultures* **18**, 256–261.
- VENEGAS-CALERÓN, M., SAYANOVA, O. & NAPIER, J. A. (2010). An alternative to fish oils: metabolic engineering of oil-seed crops to produce omega-3 long chain polyunsaturated fatty acids. *Progress in Lipid Research* **49**, 108–119.
- VORNANEN, M., HASSINEN, M., KOSKINEN, H. & KRASNOV, A. (2005). Steady state effects of temperature acclimation on the transcriptome of the rainbow trout heart. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology* **289**, 1177–1184.
- WATTERDORF, R. J. (1986). Rapid identification of triploid grass carp with Coulter counter and channelizer. *Progressive Fish Culturist* **48**, 125–132.
- WHITAKER, H. A., MCANDREW, B. J. & TAGGART, J. B. (2006). Construction and characterization of a BAC library for the European sea bass *Dicentrarchus labrax*. *Animal Genetics* **37**, 526.
- WOLTERS, W. R., LILYESTROM, C. G. & CRAIG, R. J. (1991). Growth, yield and dress-out percentage of diploid and triploid channel catfish in earthen ponds. *The Progressive Fish Culturist* **53**, 33–36.
- WU, G., TRUKSA, M., DATLA, N., VRINTEN, P., BAUER, J., ZANK, T., CIRPUS, P., HEINZ, E. & QIU, X. (2005). Stepwise engineering to produce high yields of very long-chain polyunsaturated fatty acids in plants. *Nature Biotechnology* **23**, 1013–1017.
- XU, P., WANG, S., LIU, L., THORSEN, J., KUCUKTAS, H. & LIU, Z. J. (2007). A BAC based physical map of the channel catfish genome. *Genomics* **90**, 380–388.
- ZHANG, P. J., HAYAT, M., JOYCE, C., GONZALEZ VILLASENOR, L. I., LIN, C. M., DUNHAM, R. A., CHEN, T. T. & POWERS, D. A. (1990). Gene-transfer, expression and inheritance of prsv-rainbow trout-gh cDNA in the common carp, *Cyprinus carpio* (Linnaeus). *Molecular Reproduction and Development* **25**, 3–13.
- ZHU, Z., LI, G., HE, L. & CHEN, S. (1985). Novel gene transfer into the fertilised eggs of goldfish (*Carassius auratus* 1758). *Journal of Applied Ichthyology* **1**, 31–33.