

1 **AMPHIPOD INTERSEX, METALS AND LATITUDE: A PERSPECTIVE**

2 M. Ramiro Pastorinho^{1*}, Trevor C. Telfer² and Amadeu M.V.M. Soares¹

3 *1-Center for Environmental and Marine Studies (CESAM) and Department of Biology,*
4 *University of Aveiro, Portugal*

5 *2-Environmental Group, Institute of Aquaculture, University of Stirling, United Kingdom*

7 **ABSTRACT**

8 Intersexuality has been widely reported in crustaceans with several mechanisms being
9 directly or indirectly held responsible for its occurrence, amongst which pollution. No
10 mechanistic relationship between metals and intersex has ever been established. Also the
11 incidence of intersex in populations of the same invertebrate species in a latitudinal gradient
12 has never been studied so far. Three populations (Iceland, Scotland and Portugal) of the
13 amphipod *Echinogammarus marinus* were scrutinized. Intersex females from Iceland
14 registered the highest fecundity loss. Only in Scottish samples females with two genital
15 papillae and males with only one genital papillae were observed. Nevertheless, water, biota
16 and sediment samples pointed to equivalent metal levels, and in consonance the prevalence
17 of intersex was not significantly different between locations. An unequivocal relationship

* Corresponding Author: University of Aveiro, Department of Biology, Campus de Santiago,
3810-193 Aveiro, Portugal. Fax: +351 234 372 587; e-mail: rpastorinho@ua.pt

between metal presence and intersex induction cannot be presented, but our results advocate the potential role of metals as a direct cause of intersexuality in *E. marinus*.

KEYWORDS: *Echinogammarus marinus*; intersex; heavy metals; Iceland; Scotland; Portugal.

1. INTRODUCTION

Intersexuality, the condition whereby otherwise normal gonochoristic species possess both male and female characteristics, has been reported throughout the animal kingdom (Reinboth, 1975). This phenomenon has been widely identified in crustaceans, among which a large number of amphipod species (*e.g.* Sars, 1895; Sexton and Huxley, 1921; Dunn et al., 1990; Ladewig et al., 2003; Ford and Fernandes 2005).

Several mechanisms are directly or indirectly held responsible for its occurrence including parasitism (Bulnheim, 1977); bacterial infection (Rigaud and Juchault, 1998); genetic control (Lebederf, 1939); the ablation or implantation of androgenic glands, from/into males and females, respectively (Charniaux-Cotton, 1958); environmental sex determination (ESD) (Dunn et al., 1993) and pollution (Moore and Stevenson, 1991; Ford et al., 2006). Recently it has been demonstrated to be induced chemically by juvenile hormone mimics (Olmstead and LeBlanc, 2007).

The full significance and costs of intersexuality for amphipods are still unknown. Some studies report lower fecundity and fertility, reduced pairing success and delayed maturation in *Echinogammarus marinus* (Ford et al., 2003; Ford and Fernandes, 2005). Dunn et al. (1993) also reported reduced pairing success in *Gammarus duebeni* and Kelly et al. (2004) found

lower fecundity in intersex females of the same species. An investigation by Plaistow et al. (2003) on the costs of intersex upon precopulatory guarding in *Gammarus pulex*, indicated that these were correlated with female size. Consequently, pair forming with larger intersex females (which grow larger than the normal counterpart) will lead not only to large energy costs, but also reduced reproductive success. Barbeau and Grecian (2003) noted reduced fertility in intersex male *Corophium volutator*, a finding confirmed for the same species by McCurdy et al. (2004) that registered smaller broods issuing from the pairing of intersex males with normal females compared to matings with normal males.

A model simulation for an *E. marinus* population run by Ford et al. (2007) to evaluate the reproductive costs associated with intersexuality revealed that intersexes can have dramatic effects: population extinction within approximately 6 years if 5% of normal females are replaced by intersex females. Nevertheless, in behavioral terms no differences appear to occur with intersex females and intersex males acting as do normal females and males, respectively (Ford et al., 2007).

Besides the classical case study of organotin-induced imposex and intersex in gastropods (Matthiessen and Gibbs, 1998), continued studies of metals endocrine disrupting effects in invertebrates are scarce, leading to inconsistent knowledge on the subject. More recently a growing body of work has been trying to bridge this gap of knowledge (e.g. Medesani et al., 2004; Bondgaard and Bjerregaard, 2005; Corrêa et al., 2005; Ford et al., 2006), as the study of metals as endocrine disruptors can be considered highly relevant bearing in mind their

ubiquitous presence in the environment (from both natural and anthropogenic sources) and in metabolic processes.

Echinogammarus marinus (Leach, 1815) (= *Chaetogammarus*; = *Marinogammarus*) is a highly abundant amphipod species of the marine and estuarine intertidal north-eastern Atlantic (Lincoln, 1979), with documented intersex prevalence (Ford et al., 2003). This species occurs in close association with *Fucus spp.* (Phaeophyta, Fucales), which is used by the amphipod as source of food and shelter to live and breed (Schreider et al., 2003).

To the author's best knowledge, no mechanistic relationship between endocrine disruptive capabilities of metals and the occurrence of intersex in crustaceans has ever been established. The incidence of intersex in populations of an invertebrate species in a latitudinal gradient is also unknown. In this context, this study aimed to ascertain the incidence of intersexuality in independent populations of *E. marinus* (encompassing the entire breadth of its distribution), while at the same time attempted to unveil possible correlations with the presence of heavy metals.

2. MATERIALS and METHODS

2.1. Study sites

The organisms used in this study to evaluate intersex variation in latitudinal terms were collected in Iceland, Scotland and Portugal (Figure 1). A literature survey was performed to choose specific sites where *Echinogammarus marinus* occurs and metal levels were of approximately similar magnitude.

Icelandic individuals were captured in Reikjanes Peninsula (64°02'N, 22°42'W). Despite these shores being considered pristine environments (Sarà et al., 2007), there is indication of heavy metal background values being higher than in equally remote areas. Volcanic activity is regarded as the probable cause (IME, 2001), mainly because eruptions substantially increase metals concentration in Icelandic rivers. The redistribution of these metals is efficiently performed by the unique hydrography of the Icelandic coastal current, which begins in the northeast, moves clockwise and ends at the mid-north coastal areas (Egilsson et al., 1999). The closest human settling from the collection site is the small town of Sandgerði (1700 inhabitants), which is one of the most important fishing harbours in Iceland, where fish processing facilities are installed. Olafsson (1986) used *Mytilus edulis* to run a “mussel watch”-type survey on the south-western coast to identify anthropogenically enhanced metal concentrations. The conclusion was that localized spots existed, being the Sandgerði area one of them. This verdict was confirmed by Leung et al. (2005) who allude to Sandgerði as a “polluted site”.

Loch Fyne is the longest, deepest fiordic sea loch of Scotland, being part of the Firth of Clyde watershed (one of the country's most contaminated). In spite of this, it was expected to be relatively free from metal contamination due to serving mainly forestry, extensive grazing, aquaculture and recreation purposes (SEPA, 2005). Leung et al. (2001) detected higher metal contents in tissues of *Nucella lapillus* from this area than in individuals collected in areas traditionally described as metal polluted. The organisms for the present study were collected

in the vicinity of Strachur (56°10'N, 5°05'W), located a few km to the south of a "Marine Consultation Area" (NCC, 1986).

The Mondego Estuary is located in the Portuguese Atlantic coast. It comprises two arms (north and south) with very different characteristics. North arm is highly hydrodynamic, possesses navigational capabilities and it is pointed as having pristine conditions in terms of heavy metals (Vale et al., 2002). South arm dynamics are much weaker, depending partly of the tidal excursion and of the artificially controlled discharges (according to the necessities of the extensive lower valley rice crops) of a small river – the Pranto. This leads to extended residence times and fine particles deposition. The Mondego possesses a sizeable watershed (6670 km²) upon which the impact of urban areas summing up to 695.000 inhabitants is felt (INAG, 2004). Few industries are to be noted, but deactivated uranium mines are located approximately 100km from the estuary and increases of metals, particularly Zn, Mn, Fe, U, Pb and Ra (Pinto et al., 2004) are registered in the areas directly receiving the mine effluents. The organisms were collected in the south arm (40°07'N, 8°49'W) in the confluence with Pranto river. Castro et al. (2007) noted an increase of Pb in sediment profiles from this area since the 1960's, but emphasized that the Mondego was not heavily contaminated compared to other European estuaries. Pereira et al. (2005) indicated enrichment in Hg, Cu, Cd, Cr, Mg, Zn and Fe of the fine fraction of the south arm, compared to the remainder of the estuary, pointing the Pranto river as the presumable source.

2.2. Sampling

E. marinus specimens were collected by gently scraping the surface of intertidal rocks with attached algae and were readily preserved in 75% ethanol. Simultaneously, samples of brown algae (*Fucus spp.*), water and sediment from the collection sites were collected for the analysis of metal contents. Sediment was collected with a Van Veen grab, and each sample consisted of the top 2 cm of three grabs. No sediment was collected in Iceland due to the rocky nature of the substrate at the sampling site.

2.3. Intersex analysis

Seven hundred and five adult specimens were collected (Iceland = 230, Scotland = 254, and Portugal = 221), sexed (females: presence of four pairs of brood plates – oostegites – in the thoracic region; males: presence of two genital papillae (GP) between the last pair of pleopods), observed for external intersex characteristics (presence of both structures) and measured (distance between the base of the first antenna to the base of the telson) (Ford et al., 2003). In ovigerous females, eggs were removed from the brood pouch (formed by the oostegites) and counted.

2.4. Metal analysis

2.4.1. Sediment

Extraction of metals from sediments was carried out using a nitric acid–hydrogen peroxide digestion. Approximately 2g wet weight (ww) of sediment per sample site was dried at 110°C for 24 hours and ground to a fine powder. Three 500mg sub-samples were added to 20ml

137 Teflon screw top digestion vessels. 5ml of concentrated nitric acid (69%, Aristar, BDH, 106
138 U.K.) was added and the sample was heated to 110°C for 24-hours. Once cooled, 3ml of
139 hydrogen peroxide (Aristar, BDH, U.K.) was added in 1 ml steps until the sample became
140 totally clear and ceased effervescing. Samples were re-heated at 110°C for a further 2 hours,
141 allowed to cool, made up to 15ml with distilled water and centrifuged at 2000 rpm for 15
142 minutes. Quantification (cadmium, copper, nickel and zinc) was performed using a THERMO™
143 ICP - Mass Spectrophotometer.

144 **2.4.2 *Fucus spp* and *E. marinus* tissues**

145 The algae were washed with deionised water to remove adhering sediment and the bladders
146 and tips were removed by tearing (Dobson, 2000). Tissues of amphipods and algae were
147 digested and subsequently analyzed using the methods for metals analysis previously
148 described (sub-section 2.4.1).

149 **2.5. Statistical analysis**

150 One-way Analysis of Variance (ANOVA) was used to evaluate the existence of significant
151 differences between groups of the different stations and between sampled compartments
152 among stations. When data proved to have a non-normal distribution (by application of the
153 Kolmogorov-Smirnov test) the Kruskal-Wallis test (ANOVA on ranks) was used. Tukey Test or
154 Dunn's Method multiple comparison procedures (normal or non-normal distributed data,
155 respectively) helped elucidate further the differences within groups of the same station. All
156 statistical analysis was performed using SigmaStat (Version 3.1) statistical software.

3. RESULTS

3.1. Organism size

The size of the organisms was variable in all three stations (Figure 1) and within each station decreased in the following order: intersex males, normal males, intersex females and normal females. All groups were significantly different ($p < 0.001$), except normal males and intersex females ($p > 0.05$). Intersex specimens were consistently bigger than the normal counterparts (Figure 2).

3.2. Intersex incidence

In general terms, intersex incidence was higher in Scotland (18.5%), followed by Portugal (14.3%) and Iceland (10.0%). Females registered higher intersex rates, once again with Scotland leading with 13.0%, followed by Portugal (9.5%) and Iceland (6.7%) (Table 1). Only in Scottish samples was possible to find females with two genital papillae (3.7% of the individuals analysed, Table 1).

Regarding males, the general pattern was once again repeated, though with more discreet differences, namely with 5.6%, 4.8% and 3.3% for Scotland, Portugal and Iceland, respectively (Table 1). Males with only one genital papillae were only observed in samples from Scotland (1.9%, Table 1).

3.3. Female fecundity

Normal ovigerous female's average fecundity (assessed by the number of eggs present in the brood cavity) showed little variation between the three locations (15.0, 16.7 and 15.3 for Scotland, Portugal and Iceland, respectively) (Table 1). Intersex specimens registered a generalized decline in these values: 13.2 for Scotland, 13.0 for Portugal and a pronounced decrease for Icelandic organisms, with 7.9 eggs/female (Table 1).

3.4. Metal Analysis

Figure 3 presents data on the analysis of water, sediment (except Iceland), *Fucus spp.* tissue and *E. Marinus* tissue of samples collected at the three sites, respectively. Significant differences (evaluated by pairwise comparison methods – Tukey Test for water, *Fucus spp* tissue and *E. marinus* tissue; Dunn's Method for sediment) showed no consistent patterns among the three sites for any of the analysed compartments. Nevertheless, Scotland systematically presented the most elevated metal concentrations for all compartments (with the exception of water samples from Iceland).

4. DISCUSSION

The relationship between pollution and the occurrence of intersex in crustaceans was never irrefutably established. The available literature provides several examples of studies that failed to prove this connection (despite some of them produced evidence that other endocrine disruptive effects were occurring). Marine copepod communities (*Paramphiascella hyperborean*, *Stenhelia gibba* and *Halectinosoma sp.*) associated with sewage outfalls

194 registered elevated numbers of intersex individuals (Moore and Stevenson, 1991). The
195 simultaneous occurrence of this phenomenon in several species indicated the involvement of
196 a common environmental factor. However, no clear relationship could be established
197 between exposure to sewage effluent and incidence of intersex (Moore and Stevenson,
198 1991). Gross et al. (2001) surveyed *Gammarus pulex* populations from an area where sewage
199 effluent was being discharged, but failed to reveal any relationship between intersex
200 incidence and the exposure, in spite of a highly significant number of females having oocytes
201 containing a reduced number of yolk bodies and lipid globules (suggesting reduced
202 vitellogenesis). In a study using populations of *Gammarus fossarum* (Jungmann et al., 2004),
203 the incidence of intersex increased among gammarids transplanted from a location with a
204 low incidence of intersex to a location having a high incidence of intersex, suggesting that
205 some environmental factor was responsible for the high incidence of intersex at some
206 localities. Sites considered unaffected by this factor had incidences of intersex lower than
207 1%. Yet, no discernible relationship between intersex incidence and pollution could be
208 established (Jungmann et al., 2004).

209 The interpretation of the results from the present study could point in a different direction.
210 The differences obtained among locations for all the four sampled compartments (water,
211 sediment, *Fucus spp.* tissues and *E. marinus* tissues) were in its majority non-significant. This
212 means that, from a metal contamination point of view, the initial literature-based selection
213 of sampling sites, which aimed to select locations with a profile of limited disturbance and a
214 moderate presence of metals above natural backgrounds, was proper. A similar setting for all

three sampled populations was thus confirmed. Coincidentally, the incidence of intersex in the three sampling sites was also not significantly different (one way ANOVA, $p>0.05$). Seemingly, the statistics-based conclusion from this study is that when exposed to approximately the same environmental metal concentrations, populations of *E. marinus* show similar incidences of intersex. However, we are aware that despite being statistic inference a fundamental tool, it should not be the sole contributor to the interpretation of a data set. Discreet differences can be otherwise detected, allowing the observer to retain palpable information regarding the capability of a certain location to exert a more marked influence upon organisms than others judged (e.g. by statistical analysis). Nevertheless, an “influence ranking” of the three sites is hard to establish. Chemical stressors, like heavy metals, tend to be discreetly present in the water column both in time and space (Hall et al., 1998). Periodical disturbances may occur and these pulse events (of natural and anthropogenic origin) when ended can leave the physical and chemical environment similar to pre-disturbance conditions (Bender et al., 1984). If sampling occurs in these intermediate periods, erroneous conclusions could be drawn in terms of frequency and levels of perturbations.

Icelandic organisms are likely to be more frequently influenced by natural pulse-like perturbation, whilst Scottish and Portuguese organisms more liable to frequently receive anthropogenic influences. These, due to their diverse and complex nature (e.g. accidental spills, urban runoff, licensed periodic discharges) represent added stress upon these organisms (atop natural occurring events). In the present study, the differences in intersex female fecundity between sites could represent a sign of this influence. Charnov (1982)

236 predicted that intermediates (=intersexes) should incur in reduced fitness. Our data
237 confirmed that overall prediction, but Icelandic intersex females had a two-fold decrease in
238 fecundity when compared to Portuguese individuals (48% versus 22%) and four-fold when
239 compared to Scottish individuals (48% versus 12%). *E. marinus* eggs are relatively large for
240 amphipods (Clarke et al., 1985) and as expected in the arcto-boreal zone they become even
241 larger (Skadsheim, 1984). This means that intersex females from Iceland will have more
242 difficulties in maintaining their brood inside the marsupium, especially if we consider that
243 one of the effects of intersex is the parcial or total loss of the setae closing the brood pouch
244 (Ladewig et al., 2003).

245 In a scenario of pollution-caused intersex, intermediates from populations that have been
246 inhabiting for numerous generations environments where recurrent episodes of stress occur,
247 hence subjected to selection mechanisms (Luoma, 1977), would be more prone to cope with
248 that fecundity loss, compared to individuals living in areas were those episodes are more
249 sparse. This idea is reinforced by given proof of a genetic basis for metal tolerance in
250 crustaceans (Barata et al., 1998; Ross et al., 2002). Besides this gradation in fecundity,
251 another fact reinforcing a discrepancy between locations is that females with two genital
252 papillae and males with only one genital papillae were only present in Scottish samples.
253 Despite a certain similarity in the observed conditions, this occurrence allows a
254 differentiation between the intensity of the intersex mechanisms affecting the Scottish
255 population and the ones affecting the Portuguese population.

Recent work by Ford et al. (2006) confirmed a consistently higher level of intersexuality in *E. marinus* throughout the year at sites receiving industrial contaminants (metal rich pulp mill effluent) compared to reference sites. In this context, one can envisage environmental constraints modulating pollution-derived intersex (possibly metal pollution), keeping the former an open door to make intermediates viable. Furthermore, one should expect that this situation would only occur in sites with marginal and not heavy pollution. A plausible scenario for *E. marinus* intersex, as it was observed in the present study.

In our study, specific focus was put upon the environmental occurrence of metals and their possible relationship with intersexuality. An unequivocal relationship between the two factors, such as for instance the case of organotin-caused intersex in gastropods (Matthiessen and Gibbs, 1998) cannot be presented, but our results advocate the potential role of metals as a direct cause of intersexuality in *E. marinus*.

Acknowledgements

We are very grateful to Dr. Gudmundur V. Helgason and Dr. Halldór P. Halldórson for arrangements and hospitality at the Sandgerði Marine Centre, Iceland (SMC). Work at the SMC was supported by the programme “Improving Human Potential: Access to Research Infrastructures” (IHP/ARI) provided by the European Community under Marie Curie Actions - EC-IHP Transnational Access Grant. This research was funded through a PhD Grant by FCT –

275 Fundação para a Ciência e Tecnologia (SFRH/BD/4778/2001). We are also extremely grateful
276 to the two anonymous reviewers whose comments helped improve this manuscript.

277

278 REFERENCES

279 Barata, C., Baird, D.J. & Markich, S.J. 1998. Influence of genetic and environmental factors on
280 the tolerance of *Daphnia magna* Straus to essential and non-essential metals. Aquatic
281 Toxicology 42, 115-137.

282 Barbeau, M.A. & Grecian, L.A. 2003. Occurrence of intersexuality in the amphipod *Corophium*
283 *volutator* (Pallas) in the upper Bay of Fundy, Canada. Crustaceana 76, 665-679.

284 Bender, E.A., Case, T.J. & Gilpin, M.E. 1984. Perturbation experiments in community ecology:
285 Theory and practice. Ecology 65, 1–13.

286 Bondgaard, M. & Bjerregaard, P. 2005. Association between cadmium and calcium uptake
287 and distribution during the moult cycle of female shore crabs, *Carcinus maenas*: an in
288 vivo study. Aquatic Toxicology 72, 17–28.

289 Bulnheim, H.-P. 1977. Sexual transformation in *Gammarus duebeni* (Crustacea, Amphipoda)
290 under the influence of hormonal and parasitic factors. Biologisches Zentralblatt 96, 61-
291 78.

292 Castro, P., Valiela, I. & Freitas, H. 2007. The use of sedimentary %C, %N, $\delta^{15}\text{N}$, and Pb
293 concentrations to assess historical changes in anthropogenic influence on Portuguese
294 estuaries. *Environmental Pollution* 147, 706-712.

295 Charniaux-Cotton, H. 1958. La gland androgène de quelques Crustacés Décapodes et
296 particulièrement de *Lysemata seticaudata*, espèce à hermaphrodisme protérandrique
297 fonctionnel. *Cahiers de Recherche de L'Académie de Sciences de Paris* 246, 2814–2817.

298 Charnov, E.L. 1982. The theory of sex allocation. *Monographs in Population Biology* 18, 1-
299 355.

300 Clarke, A., Skadsheim, A. & Holmes, L.J. 1985. Lipid biochemistry and reproductive biology in
301 two species of Gammaridae (Crustacea: Amphipoda). *Marine Biology* 88, 247-263.

302 Corrêa Jr., J.D., Silva, M.R., Silva, A.C., Lima, S.M., Malm, O. & Allodi, S. 2005. Tissue
303 distribution, subcellular localization and endocrine disruption patterns induced by Cr
304 and Mn in the crab *Ucides cordatus*. *Aquatic Toxicology* 73, 139-154.

305 Dobson, J. 2000. Long term trends in trace metals in biota in the Forth Estuary, Scotland,
306 1981–1999. *Marine Pollution Bulletin* 40, 1214–1220.

307 Dunn, A.M., Adams, J. & Smith, J.E. 1990. Intersexes in a shrimp: a possible disadvantage of
308 environmental sex determination. *Evolution* 44, 1875-1878.

309 Dunn A.M., Adams, J. & Smith, J.E. 1993. Is intersexuality a cost of environmental sex
310 determination in *Gammarus duebeni*? *Journal of the Zoological Society of London* 231,
311 383-389.

- 312 Egilsson, D., Ólafsdóttir, E.D., Ingvadóttir, E., Halldórsdóttir, H., Sigurðsson, F.H., Jónsson,
313 G.S., Jensson, H., Gunnarsson, K., Thráinsson, S.A., Stefànsson, A., Indriðason, H.D.,
314 Hjartarson, H., Thorlacius, J., Ólafsdóttir, K., Gíslason, S.R. & Svavarsson, J. 1999.
315 Mælingar á mengandi efnum á og við Ísland. Niðurstöður vöktunarmælinga.
316 Starfshópur um mengunarmælinga, Umhverfissráðuneytið (Measurements of
317 contaminants in and near Iceland, Results from monitoring studies, Working group of
318 monitoring, Ministry for the Environment), Reykjavík, 138 pp.
- 319 Ford, A.T. & Fernandes, T.F. 2005. Notes on the occurrence of intersex in amphipods.
320 Hydrobiologia 548, 313-318.
- 321 Ford, A.T., Fernandes, T.F., Rider, S.A., Read, P.A., Robinson, C.D. & Davies, I.M. 2003.
322 Reproduction in the amphipod, *Echinogammarus marinus*: a comparison between
323 normal and intersex specimens. Journal of the Marine Biological Association of the U.K.
324 83, 937-940.
- 325 Ford, A.T., Fernandes T.F., Robinson C.D., Davies I.A. & Read P.A. 2006. Can industrial
326 pollution cause intersexuality in the amphipod, *Echinogammarus marinus*? Marine
327 Pollution Bulletin 53, 100-106.
- 328 Ford, A.T., Martins, I. & Fernandes, T.F. (2007) Population level effects of intersexuality in the
329 marine environment. Science of the Total Environment 374, 102–111.

330 Gross, M.Y., Naycock, D.S., Thorndyke, M.C., Morritt, D. & Crane, M. 2001. Abnormalities in
 331 sexual development of the amphipod *Gammarus pulex* (L.) found below sewage
 332 treatment works. *Environmental Toxicology and Chemistry* 20, 1792–1797.

333 Hall Jr., L.W., Scott, M.C. & Killen, W.D. 1998. Ecological risk assessment of copper and
 334 cadmium in surface waters of Chesapeake Bay watershed. *Environmental Toxicology*
 335 *and Chemistry* 17, 1172–1189.

336 IME 2001. Iceland's National Program of Action for the protection of the marine environment
 337 from land-based activities. Iceland Ministry of Environment, Gudjonø, Reykjavic.

338 INAG 2004. (Portuguese Institute for the Water) Plano Nacional da Água, Lisboa.

339 Jungmann, D., Ladewig, V., Ludwichowski, K.U., Petzsch, P. & Nagel, R. (2004) Intersexuality
 340 in *Gammarus fossarum* KOCH—a common inducible phenomenon? *Archives of*
 341 *Hydrobiology* 159, 511–529.

342 Kelly, A., Hatcher, M. J. & Dunn, A.M. 2004. Intersexuality in the amphipod *Gammarus*
 343 *duebeni* results from incomplete feminisation by the vertically transmitted parasitic sex
 344 ratio distorter *Nosema granulosa*. *Evolutionary Ecology* 18, 121–132.

345 Landewig, V., Jungmann, D., Koehler, A., Schirling, M., Triebkorn, R. & Nagel, R. 2003.
 346 Intersexuality in *Gammarus fossarum* koch, 1835 (amphipoda). *Crustaceana* 75, 1289-
 347 1299.

348 Lebederf, G.A. 1939. A study of intersexuality in *Drosophila virilis*. *Genetics* 24, 553-586.

349 Leung, K.M.Y., Dewhurst, R.W., Halldórsson, H.P. & Svavarsson, J. 2005. Metallothioneins and
350 trace metals in the dogwhelk *Nucella lapillus* (L.) collected from Icelandic coasts.
351 Marine Pollution Bulletin 51, 729–737.

352 Leung, K.M.Y., Morgan, I.J., Wu, R.S., Lau, T.C., Svavarsson, J. & Furness, R.W. 2001. Growth
353 rate as a factor confounding the use of the dogwhelk *Nucella lapillus* as biomonitor of
354 heavy metal contamination. Marine Ecology Progress Series 221, 145-159.

355 Lincoln, R.J. 1979. British marine Amphipoda: *Gammaridea*, British Museum (Natural
356 History), London, pp. 658.

357 Luoma, S.N. 1977. Detection of trace contaminant effects in aquatic ecosystems. Journal of
358 the Fisheries Research Board of Canada 34, 436–439.

359 Matthiessen, P. & Gibbs, P.E. 1998. Critical appraisal of the evidence for tributyltin-mediated
360 endocrine disruption in mollusks. Environmental Toxicology and Chemistry 17(1), 37–
361 43.

362 McCurdy, D.G., Forbes, M.R., Logan, S.P., Kopec, M. & Mautner, S. 2004. The functional
363 significance of intersexes in the intertidal amphipod, *Corophium volutator*. Journal of
364 Crustacean Biology 24, 261–265.

365 Medesani, D.A., López Greco, L.S. & Rodríguez, E.M. 2004. Interference of cadmium and
366 copper with the endocrine control of ovarian growth, in the estuarine crab
367 *Chasmagnathus granulata*. Aquatic Toxicology 69, 165–174.

368 Moore, C.G. & Stevenson, J.M. 1991. The occurrence of intersexuality in harpacticoid
369 copepods and its relationship with pollution. *Marine Pollution Bulletin* 22, 72-74.

370 NCC 1986. Nature Conservancy Council, Marine Consultation Areas, NCC, Peterborough.

371 Olafsson, J. 1986. Trace metals in mussels (*Mytilus edulis*) from Southwest Iceland. *Marine*
372 *Biology* 90, 223-229.

373 Olmstead, A.W. & Leblanc, G.A. 2007. The environmental-endocrine basis of
374 gynandromorphism (intersex) in a crustacean. *International Journal of Biological*
375 *Sciences* 3, 77–84.

376 Pereira, P., Vale, C., Ferreira, A.M., Pereira, E., Pardal, M.A. & Marques, J.C. 2005. Seasonal
377 variation on surface sediments composition in Mondego River Estuary. *Journal of*
378 *Environmental Science and Health A40*, 317-329.

379 Pinto, M.M.S.C., Silva, M.M.V.G. & Neiva, A.M.R. 2004. Pollution of water and stream
380 sediments associated with the Vale de Abrutiga uranium mine, central Portugal. *Mine*
381 *Water and the Environment* 23, 66–75.

382 Plaistow, J., Bollache, L. & Cézilly, F. 2003. Energetically costly precopulatory mate guarding
383 in the amphipod *Gammarus pulex*: causes and consequences. *Animal Behavior* 65, 683–
384 691.

385 Reinboth, E. 1975. Intersexuality in the animal kingdom. Springer-Verlag, Berlin.

386 Rigaud, T. and Juchault, P. 1998. Sterile intersexuality in an isopod induced by the interaction
387 between a bacterium (wolbachia) and the environment. Canadian Journal of Zoology
388 76, 493-499.

389 Ross, K., Cooper, N., Bidwell, J.R. & Elder J. 2002. Genetic diversity and metal tolerance of
390 two marine species: a comparison between populations from contaminated and
391 reference sites. Marine Pollution Bulletin 44, 671–679.

392 Sarà, G., De Pirro, M., Romano, C., Rumolo, P., Sprovieri, M. & Mazzola, A. 2007. Sources of
393 organic matter for intertidal consumers on Ascophyllum-shores (SW Iceland): a multi-
394 stable isotope approach. Helgoland Marine Research 61, 297-302.

395 Sars, G.O. 1895. Amphipoda. An account of the crustacean of Norway with short descriptions
396 and figures of all the species, Vol.1. Cammermeyers Forlag: Christiania Copenhagen,
397 pp. 711.

398 Schreider, M.J., Glasby, T.M. & Underwood A.J. 2003. Effects of height on the shore and
399 complexity of habitat on abundances of amphipods on rocky shores in New South
400 Wales. Journal of Experimental Marine Biology and Ecology 293, 57–71.

401 Scottish Environmental protection Agency (SEPA) (2005) - Loch Fyne Coastal Strip Report -
402 Report # 9.

403 Sexton, E.W. & Huxley, J.S. 1921. Intersexes in *Gammarus cheureuxi* and related forms.
404 Journal of the Marine Biological Association of the U.K. 12, 506-556.

Skadsheim, A, 1984. Coexistence and reproductive adaptations of amphipods: the role of environmental heterogeneity. *Oikos* 43, 94-103 .

Vale, C., Ferreira, A., Caetano, M. & Brito, P. 2002. Elemental composition and contaminants in surface sediments of the Mondego river estuary. In M.A. Pardal, J.C. Marques, M.A. Graça, Aquatic Ecology of the Mondego River Basin. Global Importance of Local Experience, pp. 243–256. Imprensa da Universidade de Coimbra, Coimbra.

FIGURE CAPTIONS

Figure 1 - Geographic location of the three sampling sites.

Figure 2 - Organism length in the three locations. Error bars denote standard deviations.

Figure 3 - Metal concentrations in samples of water [A], sediments [B], *Fucus spp.* tissue [C] and *E. marinus* tissue [D] collected from the sites in Iceland, Scotland and Portugal (in ug/l). Error bars denote standard deviations. Letters (a, b) denote statistically significant differences ($p<0.05$) between locations for each metal analysed. Nickel results were not obtained for Portuguese sediments.

TABLE CAPTIONS

423 **Table 1-** Incidence of intersex (%) for males and females in the three stations and average
424 number of eggs inside females brood pouch. GP- genital papillae.



Figure 1




Figure 2




Figure 3

	Females			Males		
	Iceland	Scotland	Portugal	Iceland	Scotland	Portugal
Normal	33.3	29.6	33.3	56.7	51.9	52.4
Avg egg	15.3	15.0	16.7	-	-	-
Intersex	6.7	13.0	9.5	3.3	5.6	4.8
Avg egg	7.9	13.2	13.0	-	-	-
1 GP	6.7	9.3	9.5	0	1.9	0
2 GP	0	3.7	0	-	-	-

Table 1