

1 Effects of dietary protein and fat level and rapeseed oil on growth and
2 tissue fatty acid composition and metabolism in Atlantic salmon (*Salmo*
3 *salar* L.) reared at low water temperatures

4
5 **V. KARALAZOS¹, E.Å. BENDIKSEN², J. R. DICK¹, J.G. BELL¹**

6
7 ¹*Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK, ²BioMar AS,*
8 *N-7484, Trondheim, Norway*

9
10 *Correspondence: Vasileios Karalazos, Institute of Aquaculture, University of Stirling, Stirling*
11 *FK9 4LA, Scotland. Tel.: +44-1786-467993; Fax.: +44-1786-472133. E-mail:*
12 *vasileios.karalazos@stir.ac.uk*

13
14
15
16 *Running title: Protein/fat level and rapeseed oil in Atlantic salmon diets*

17

18

19

20

21

22

23

24

25

Abstract

A 12 week feeding trial was conducted to elucidate the interactive effects of dietary fat and protein contents and oil source on growth, fatty acid composition, protein retention efficiency (PRE) and β -oxidation activity of muscle and liver in Atlantic salmon (*Salmo salar* L.) at low water temperatures (4.2 °C). Triplicate groups of Atlantic salmon (initial weight 1168 g) were fed six isoenergetic diets formulated to provide either 390 g kg⁻¹ protein and 320 g kg⁻¹ fat (high protein (HP) diets) or 340 g kg⁻¹ protein and 360 g kg⁻¹ fat (low protein (LP) diets); within each dietary protein/fat level crude RO comprised 0, 30 or 60% (R0, R30, R60, respectively) of the added oil. After 12 weeks the overall growth and FCR were very good for all treatments (TGC; 4.76 (\pm 0.23), FCR; 0.85 (\pm 0.02)). Significant effects were shown due to oil source on SGR and TGC only. The liver and muscle FA compositions were highly affected by the graded inclusion of RO. The PRE was significantly affected by the dietary protein level, while no significant effects were shown in total β -oxidation capacity of liver and muscle. The results of this study suggest that more sustainable, lower protein diets with moderate RO inclusion can be used in Atlantic salmon culture at low water temperatures with no negative effects on growth and feed conversion, no major detrimental effects on lipid and fatty acid metabolism and a positive effect on protein sparing.

KEYWORDS: Rapeseed oil; Dietary protein / lipid ratio; Polyunsaturated fatty acids (PUFA); β -Oxidation; Protein sparing effect; Atlantic salmon

50 **Introduction**

51 Traditionally, marine fish meal (FM) and fish oil (FO) have been the major
52 protein and oil sources in aquafeeds, especially for carnivorous species such as
53 Atlantic salmon, mainly due to their high nutritional value (Sargent & Tacon, 1999).
54 The aquaculture sector is at present the biggest consumer of FM and FO, consuming
55 in 2002 about 2.685 mmt of FM and 0.666 mmt of FO, that represents 42.1 and 78.7%
56 of the total global FM and FO production, respectively (Tacon, 2004). Moreover, the
57 largest proportion of the total FM and FO used in aquafeeds in 2002, 26.9 and 52.4%,
58 respectively, was consumed by salmonids (Tacon, 2004).

59 Clearly, there is a strong dependence on FM and FO for the salmon industry.
60 This could be risky and even harmful to the viability, growth and profitability of the
61 sector, as it has been estimated that the resources of wild feed grade fisheries will
62 remain static (Pike & Barlow, 2003), while the demand for these commodities by the
63 aquaculture feed industry will grow significantly in the next decade (Sargent &
64 Tacon, 1999; Tidwell & Allan, 2002; Tacon, 2004). Moreover, other issues arise, that
65 make the use of FM and FO for aquafeeds problematic; for instance, FM and FO can
66 contain organic pollutants (e.g. dioxins, PCBs, PBDEs), that are deposited in the fish
67 and, thereby, may limit their inclusion (SCAN, 2000; SCF, 2001; Jacobs *et al.*, 2002a;
68 Jacobs *et al.*, 2002b; Bell *et al.*, 2005). Hence, there is a growing, pressing need for
69 sustainable alternatives to FM and FO and for the reduction of the dependence of FM
70 and FO for fish feeds.

71 Vegetable oils (VO) represent sustainable alternatives to FO. However, the
72 replacement of FO with VO can be challenging, as VO lack the n-3 highly unsaturated
73 fatty acids (HUFA) which are abundant in FO; the n-3 HUFA, especially
74 eicosapentaenoic (20:5n-3; EPA) and docosahexaenoic acids (22:6n-3; DHA), are

essential for optimal growth and development in salmon (Sargent *et al.*, 2002). In addition, the fatty acid (FA) composition of the fish tissues reflects the FA composition of the diets; hence, replacement of FO with VO results in reductions of EPA, DHA and the n-3/n-6 FA ratio, with a direct effect on the nutritional quality of the end product (Bell *et al.*, 2001; Rosenlund *et al.*, 2001; Bell *et al.*, 2003a). This is important for the human consumer as EPA and DHA and the n-3/n-6 FA ratio have been associated with numerous beneficial effects on human health and any reduction in farmed fish would be undesirable (De Deckere *et al.*, 1998; Horrocks & Yeo, 1999; Simopoulos, 1999; Hunter & Roberts, 2000; ISSFAL, 2000; Simopoulos, 2003).

RO is considered to be a good sustainable substitute for FO. It has been used successfully in a number of previous studies with salmon (Bell *et al.*, 2001; Torstensen *et al.*, 2004a; Torstensen *et al.*, 2004b). Moreover, it has a high availability (FAO, 2005), as it is the third largest production of VO in the world, after soy and palm oil (U. S. Department for Agriculture, 2005).

Currently, salmon diets contain high proportions of protein, most of it provided by FM. However, it is crucial for the aquafeed industry to optimise the use of feed protein and to improve the protein utilisation in the salmon diets. This would allow less dependence on FM, reduce the cost of the feed and also reduce the environmental impact through waste output from salmon culture (Halver & Hardy, 2002). Salmon can utilize lipids efficiently, therefore the use of high lipid diets in salmon allows protein sparing (Froyland *et al.*, 1998; Hillestad *et al.*, 1998; Bendiksen *et al.*, 2003) and subsequently improved growth.

Numerous studies have investigated the replacement of FO with RO, and/or other VO in diets of salmonids (Bell *et al.*, 2001; Rosenlund *et al.*, 2001; Tocher *et al.*, 2001; Bell *et al.*, 2002; Bell *et al.*, 2003b; Bell *et al.*, 2003a; Bendiksen *et al.*,

2003; Ng *et al.*, 2004; Tocher *et al.*, 2004; Torstensen *et al.*, 2004a; Torstensen *et al.*, 2004b; Fonseca-Madrigal *et al.*, 2005) and also the use of low protein / high lipid diets (Einen & Roem, 1997; Bendiksen *et al.*, 2003; Azevedo *et al.*, 2004; Solberg, 2004). However, most of these studies were focusing either on the oil source or on the dietary protein/fat level. In addition, very few were conducted at the very low temperatures used in the present study that are common in sites located in high latitudes. It is known that temperature plays a significant role in FA metabolism and, in general, in fish nutrition, physiology and growth (Torstensen *et al.*, 2000; Bendiksen & Jobling, 2003; Bendiksen *et al.*, 2003; Guderley, 2004; Ng *et al.*, 2004; Tocher *et al.*, 2004). In particular, the important role of n-3 HUFAs in low temperature adaptation has been highlighted (Hochachka & Somero, 2002).

The aim of this trial was to elucidate the interactive effects of dietary fat and protein contents and oil source on growth, whole body proximate composition, fatty acid composition and β -oxidation activity of liver and muscle in Atlantic salmon at low water temperatures.

Materials and methods

Fish and facilities

Atlantic salmon (*Salmo salar*) of the NLA strain (03G) with overall mean weight of 1168g were randomly distributed into 18 sea cages of 125 m³ (5x5x5m) with 137 fish in each cage. Prior to the trial the fish were stocked in two trial cages and acclimatised for six weeks. The fish were subjected to artificial light (LD24:0) from the middle of December at decreasing ambient temperature (range; 4-6oC). During this holding period the fish were fed commercial pelleted feed (BioOptimal CPK, 9mm, BioMar AS, Norway) in accordance with the manufacturer's

recommendations. During the experimental period (February –April 2004) the fish were subjected to artificial light (LD24:0), provided from sub-merged light (one 400W bulb shared by four cages). The temperature varied from 2.8 °C to 7.3 °C with an average temperature of 4.2 ± 0.8 °C. Salinity was 34.0 ± 0.8 g L⁻¹. Fish were bulk weighed at the start of the trial, after 6 weeks and at the end of the trial (12 weeks). Mortalities were recorded and dead fish were removed daily.

Experimental diets

Six isoenergetic, practical-type extruded diets (9 mm) were formulated (BioMar TechCentre, Brande, DK) to provide either 390 g kg⁻¹ protein and 320 g kg⁻¹ fat (high protein (HP) diets) or 340 g kg⁻¹ protein and 360 g kg⁻¹ fat (low protein (LP) diets). Within each dietary fat and protein level crude RO comprised 0, 30 or 60% (R0, R30, R60) of the total added oil, the remainder of which was FO (Table 1). The diets were formulated to meet all the known nutritional requirements of salmonid fish (NRC, 1993). The proximate composition of the experimental diets is shown in Table 1 and the fatty acid compositions are shown in Table 2. Each feed was fed daily to satiation by hand to triplicate groups (cages) of fish. When sea temperature was below 5 °C the fish were fed to satiation once a day. Above 5 °C, two daily meals were provided with a minimum of 4 hours between the meals. In order to facilitate accurate calculations of feed intake and FCR, feed wastage was collected using a lift-up system and calculated on a daily basis.

Sampling procedure

Samples were taken from all diets and stored at -20 °C until analyzed. At the start of the experiment an initial sample of six fish was taken to determine baseline values of whole body proximate composition. At the end of the trial (12th week) three

fish per cage were sampled at random from the population in each cage for lipid and fatty acid composition and β -oxidation activities of liver and muscle. Another sample of three fish per cage was used for whole body proximate composition. Fish were killed with a sharp blow to the head and samples of liver were dissected and immediately placed in liquid nitrogen. Viscera, liver and heart weights from four fish per cage were recorded for measurement of viscero-somatic index (VSI), hepato-somatic index (HSI) and cardio-somatic index (CSI), respectively. For whole body analysis fish were minced and homogenate sub-samples of each fish were obtained. Initial whole body samples were pooled in pairs so three samples were finally obtained ($n = 3$) while 12 week whole body samples were pooled so there was one sample per cage. A muscle sample, representative of the edible portion, was obtained by cutting a steak between the dorsal and ventral fins (NQC). This section was then skinned, de-boned and homogenized. All samples were then stored at -20°C until analyzed.

Proximate analysis

Proximate analysis was conducted to determine the nutrient composition of diets and whole body samples. Moisture was determined by thermal drying to constant weight in an oven at 110°C for 24h. Crude protein contents were determined by Kjeldahl analyses (nitrogen $\times 6.25$, Kjeltex Autoanalyser, Tecator). Crude fat was determined in diets by acid hydrolysis using a Soxtec System 1047 hydrolysing unit (Tecator Application note 92/87) followed by exhaustive Soxhlet extraction using petroleum ether ($40-60^{\circ}\text{C}$, BP) on a Soxtec System HT6 (Tecator application note 67/83). Crude fat in whole body samples was determined by the above procedure but without the acid hydrolysis. Ash content was determined by dry ashing in porcelain

crucibles in a muffle furnace at 600 °C overnight. All methods are based on those described in AOAC (1995) and modified as described by Bell et al (2001).

Lipid extraction and fatty acid analyses

Total lipids of flesh, livers and diet samples were extracted by homogenization in 20 volumes of chloroform/methanol (2:1, v/v) containing butylated hydroxytoluene (0.01% w/w, BHT) as antioxidant, according to Folch et al. (1957). Fatty acid methyl esters (FAME) were prepared from total lipid by acid-catalyzed transesterification using 2ml of 1% H₂SO₄ in methanol plus 1 ml toluene as described by Christie (1982) and FAME extracted and purified as described by Tocher & Harvie (1988). FAME were separated and quantified by gas-liquid chromatography (Carlo Erba Vega 8160, Milan, Italy) using a 30 m x 0.32 mm capillary column (CP wax 52CB; Chrompak Ltd., London, U.K.). Hydrogen was used as carrier gas and temperature programming was from 50°C to 150°C at 40°C/min and then to 225°C at 2°C/min. Individual methyl esters were identified by comparison to known standards and by reference to published data (Ackman, 1980).

Total β -oxidation capacity

Liver and red and white muscle were weighed and homogenized in 20% (w/v) ice-cold buffered sucrose solution containing 0.25M sucrose, 0.04M potassium phosphate buffer (pH 7.4), 0.15M KCl, 40mM KF and 1mM N-acetyl cysteine. The resulting total homogenates were then centrifuged at 1880 × g for 10 min at 2°C. The resulting post-nuclear fractions were collected, and portions were used immediately to determine total (mitochondrial and peroxisomal) β -oxidation capacity. The total β -oxidation capacity was determined as acid-soluble products using radiolabelled [1-¹⁴C]-palmitoyl-CoA as a substrate as described by Frøyland *et al.* (1995).

196 *Calculations and statistical analysis*

197 The following formulae were applied to the data:

198 $\text{FCR (Feed Conversion Ratio)} = (\text{feed intake, g}) \times (\text{wet weight gain, g})^{-1}$

199 $\text{SGR (Specific Growth Rate), \% / day} = 100 \times [\ln W_1 - \ln W_0] \times (\text{days})^{-1}$

200 $\text{TGC (Thermal Growth Coefficient), } \times 1000 = 1000 \times [(W_1)^{1/3} - (W_0)^{1/3}] \times (\text{days} \times ^\circ\text{C})^{-1}$
201

202 $\text{K (Condition Factor)} = 100 \times W \times (\text{fork length, cm})^{-3}$

203 $\text{HSI \%} = 100 \times (\text{liver weight, g}) \times W^{-1}$

204 $\text{VSI \%} = 100 \times (\text{viscera weight, g}) \times W^{-1}$

205 $\text{CSI \%} = 100 \times (\text{heart weight, g}) \times W^{-1}$

206 $\text{PRE (Protein retention efficiency, g protein gain} \times \text{g protein ingested}^{-1}), \% = 100 \times$
207 $[(P_1 W_1 - P_0 W_0) \times (P_F \times \text{cumulative feed intake})^{-1}]$

208 In the above formulae W is the weight of the sampled fish in grams, W_0 and
209 W_1 are the initial and the final fish mean weights in grams, P_0 and P_1 are the initial
210 and final protein concentrations of the fish, P_F is the protein concentration of the feed
211 on a dry matter basis, and cumulative feed intake was determined in grams on a dry
212 matter basis.

213 All the data are presented as means \pm SD ($n = 3$) and all statistical analyses
214 were performed using SPSS 13 (SPSS Inc, 2004). The effects of dietary RO, the
215 fat/protein ratio and their interactions on growth, tissue fatty acid compositions, and
216 β -oxidation were analysed by two-way ANOVA. Percentage data and data which
217 were identified as non-homogeneous (Levene's test) were subjected to square root or

log transformation before analysis. Differences were regarded as significant when $P < 0.05$ (Zar, 1996).

Results

There were no significant differences in initial cage mean weights of the fish (Table 3). Following 12 weeks of feeding, the cage mean weight ranged between 1711g and 1784g and the final length of the fish varied from 49.9 to 51.8 cm. No significant effects or interactions of dietary protein level and oil source were identified in final weight and length by two-way ANOVA. However, there was a significant effect of oil source on growth performance (SGR and TGC). Specifically, the inclusion of RO resulted in higher SGR (0.49 vs. 0.56 % day⁻¹) and TGC (4.45 vs. 5.13). Feed conversion ratios (FCR) were good for all treatments and ranged from 0.81 to 0.87. No significant overall effects and interactions of dietary protein level and oil source on FCR were identified, although there was a trend ($P < 0.10$) of lower FCR for the fish fed the HP diets compared to LP diets. K ranged from 1.30 to 1.34 and no significant effects and interactions of dietary protein level and oil source were seen. The organ-to-whole-body indices are shown in Table 3. VSI varied from 11.7% to 12.6%, HSI from 1.4% to 1.5%, and similar CSIs were found for all groups (1.3%). Two-way ANOVA showed no significant effects and interactions of dietary protein level and oil source.

The PRE were high for all groups (42-47%) and there was a significant overall effect of dietary protein and fat level ($P < 0.05$), with higher overall PRE for the LP groups compared to the HP groups.

The proximate composition of whole body is shown in Table 4. Whole body moisture, protein and ash contents were very similar in all groups (approximately 662,

242 162 and 15 g kg⁻¹, respectively), whereas final lipid content ranged from 145 to 164 g
 243 kg⁻¹. Two-way ANOVA did not reveal any significant overall effects or interactions

244 The replacement of increasing proportions of FO with RO in the diets resulted
 245 in significant changes in dietary fatty acid compositions (Table 2). FO diets had
 246 approximately 30% total saturates of which two thirds was 16:0, and about 35% total
 247 monoenes with around 13% 18:1n-9. Long chain monoenes, 20:1 and 22:1,
 248 comprising more than 10% of the diet, and 5% n-6PUFA predominantly 18:2n-6, and
 249 approximately 30% n-3PUFA, with over 20% as the n-3HUFA (mainly EPA and
 250 DHA). Graded inclusion of RO resulted in decreased 16:0, 20:1n-9, 20:4n-6, 20:5n-3
 251 and 22:6n-3 (approximately 10%, 2.5%, 0.2%, 3.5% and 4.5% respectively in diets
 252 containing 60% RO) and increased 18:1n-9, 18:2n-6 and 18:3n-3 (approximately
 253 43%, 14.5% and 6% respectively in diets containing 60% RO) within both dietary
 254 protein levels. The n-3/n-6 ratio decreased from 6.0 in diets containing 100% FO to
 255 1.0 in diets containing 60% RO.

256 The total lipid content and the fatty acid compositions of muscle and liver are
 257 shown in Tables 5 & 6. Total lipid content ranged from 92.6 to 117.8 mg lipid g⁻¹
 258 tissue and from 49.4 to 81.0 mg lipid g⁻¹ tissue for muscle and liver, respectively. RO
 259 inclusion increased the total lipid content in liver but not in muscle. RO inclusion
 260 affected significantly tissue FA compositions. However, there were no significant
 261 interactions between dietary protein level and RO inclusion either in muscle or in liver
 262 and in most cases the overall protein level effect was not significant. Specifically,
 263 both in muscle and liver a reduction was seen in 16:0, total saturates, 20:1n-9, 22:1,
 264 20:4n-6, 20:5n-3, 22:6n-3, total n-3 PUFAs and n-3/n-6 ratio as RO inclusion
 265 increased within both dietary fat and protein levels. Conversely, 18:1n-9, total
 266 monoenes, 18:2n-6, total n-6 PUFAs and 18:3n-3 increased in muscle and liver with

graded RO inclusion. However, 20:2n-6 did not reflect the dietary content, as it increased both in liver and muscle with increased dietary RO inclusion.

The total palmitoyl-CoA oxidation capacity in muscle and liver is shown in Table 7. The values shown for muscle represent the total β -oxidation capacity of a combined red and white muscle fraction. β -oxidation capacity ranged from 0.45 to 0.61 pmol/min/mg protein in muscle and from 3.28 to 5.13 pmol/min/mg protein in liver. However, no significant overall effects of dietary protein/fat level or oil source were shown by two-way ANOVA.

Discussion

This study aimed to investigate the effects and interactions of the replacement of FO with RO at two different protein/lipid ratios at low water temperatures. Several previous studies have shown that replacement of FO with RO, blends of RO and other vegetable oils or other vegetable oils alone, such as linseed oil, (LO) or palm oil, (PO) in diets of salmon, has no negative effects on fish growth (Bell *et al.*, 2001; Rosenlund *et al.*, 2001; Bell *et al.*, 2002; Bell *et al.*, 2003a; Bell *et al.*, 2003b; Bendiksen *et al.*, 2003; Ng *et al.*, 2004; Torstensen *et al.*, 2004a; Torstensen *et al.*, 2004b). Moreover, when low protein feeds were compared to high protein feeds growth was not significantly affected (Azevedo *et al.*, 2004; Solberg, 2004), especially at low temperatures (Hillestad *et al.*, 1998; Bendiksen *et al.*, 2003). It is likely that the effect of low temperature masked any potential effects of feed treatment, and that diet-related growth differences observed at the higher temperature were diminished at the lower temperature (Bendiksen *et al.*, 2003). In line with the previous studies, the current experiment showed no significant effects due to dietary protein level in final weights SGR, TGC and FCRs. Nevertheless, oil source had a significant positive effect on SGR and TGC. Graded inclusion of RO, at the expense

of FO, resulted in increased SGR and TGC. This is in accordance with other studies and could be a result of enhanced protein utilisation, arising from improved use of oil for energy, due to superior fatty acid availability from vegetable oils at low water temperatures, although the exact mechanism of this effect is currently unclear (Bendiksen *et al.*, 2003). Although no significant interactions were found, the effect of oil source seemed to differ between LP and HP series. This indicates that the lower SAFAs RO oil increased the digestible energy content of the feed, resulting in improved growth performance of the fish when the dietary protein and amino acids were in excess (HP feeds).

The protein sparing effect could have also been enhanced by higher dietary oil levels. This is supported by the PRE results, as the overall PRE was significantly improved when the fish were fed the LP diets. Previous studies have also suggested a positive effect of increased dietary lipid content on protein retention and, hence, on protein sparing (Einen & Roem, 1997; Hillestad *et al.*, 1998; Bendiksen *et al.*, 2003).

In this study, the different dietary treatments had no influence on the chemical composition of the whole carcass, either due to dietary protein level or due to the oil source. The moisture, protein and ash content of the carcass were almost constant between the groups and there were only minor differences in the lipid content. This is in agreement with the findings of Hillestad *et al.* (1998) who reported that the fillet and dressed carcass fat content was not influenced by dietary lipid level, although significant differences in tissue fat content were shown, due to the dietary energy content. Other studies have shown that when the dietary oil increases, tissue lipid, and usually moisture, increases, while protein decreases (Hillestad & Johnsen, 1994; Einen & Roem, 1997; Einen & Skrede, 1998; Hemre & Sandnes, 1999; Bendiksen *et al.*, 2003; Azevedo *et al.*, 2004; Solberg, 2004). However, in most of these studies the

lipid content of the fillet, carcass or whole body was possibly influenced by the differences in the dietary energy content, as only Azevedo et al. (2004) and Solberg (2004) used isoenergetic diets. In the present study no significant differences were observed due to dietary oil source. This is in agreement with the results reported by Bendiksen et al. (2003) when FO was replaced with a blend of VO at low water temperatures (2°C). By contrast, other studies have shown that when RO replaced FO in diets of Atlantic salmon, the chemical composition of tissues is significantly influenced, to a small extent, although these studies were of longer duration than the present study (Bell *et al.*, 2001; Torstensen *et al.*, 2004b). It has been shown that in Atlantic salmon fat deposition increases as the fish grow larger (Jobling & Johansen, 2003). This was clearly demonstrated in the present study where an increase in lipid content, along with a decrease in crude protein and ash, was observed between the initial and final sampling.

In the present trial K, VSI, HSI and CSI were not affected by the different dietary treatments. This is in agreement with other studies, as changes in dietary protein / fat ratio in Atlantic salmon have not been associated with changes in K, VSI and HSI (Einen & Roem, 1997; Solberg, 2004) or with the replacement of FO with RO and LO (Rosenlund *et al.*, 2001; Bendiksen *et al.*, 2003). Few reports on diet effects on the CSI are available although a reduced CSI was observed in one study where salmon were fed diets where sunflower oil was used as a FO replacement (Bell *et al.*, 1991).

The fatty acid compositions of tissue lipids of Atlantic salmon are known to be highly influenced by dietary fatty acids (Torstensen *et al.*, 2000; Rosenlund *et al.*, 2001) and linear correlations exist between individual fatty acids in tissue total lipid and their concentrations in dietary lipid (Bell *et al.*, 2001; Bell *et al.*, 2003a; Tocher *et*

al., 2003a). In the present study, the differences in the dietary fatty acid compositions resulting from the graded inclusion of RO, at the expense of FO, were not affected by dietary protein level. Hence, diets including similar proportions of RO had very similar fatty acid compositions irrespective of their protein/lipid ratio. As a result the oil source affected significantly liver and muscle FA compositions while no significant effects on the tissue FA compositions were shown due to the dietary protein level. The results are in line with previous studies showing that dietary fatty acid compositions are reflected in tissue FA compositions.

However, previous studies have shown that although dietary fatty acids correlated to fatty acids deposited in flesh, specific fatty acids were selectively utilized or retained (Bell *et al.*, 2001; 2003a; Torstensen *et al.*, 2004a). This was also demonstrated in the present study. Dietary 18:1n-9 increased more than 3-fold and 18:2n-6 and 18:3n-3 more than 4-fold in diets containing 60% RO compared to FO diets, whereas in muscle and liver these fatty acids increased only around 2-fold. These data confirm that when certain fatty acids are provided to the fish in high concentrations, they are readily metabolised, largely catabolism by β -oxidation, although they may also be subject to limited desaturation and elongation (Bell *et al.*, 2003a). On the contrary, n-3 HUFAs were selectively deposited and retained in flesh. Tissue DHA was reduced only by 30% - 40% and EPA by less than 45% when fish were fed diets containing 60% RO compared to FO groups, whereas in 60% RO diets these HUFAs were only 30% of the concentrations in 0% RO diets. Apart from the selective deposition and retention of these FAs, the moderate reductions in EPA and DHA could have also been affected, even to small extent, by the hepatic desaturation and elongation of dietary α -linolenic acid, which can be increased by inclusion of

vegetable oils in the diets (Tocher *et al.*, 2000; Tocher *et al.*, 2001; Tocher *et al.*, 2003a; Tocher *et al.*, 2003b).

It is well documented that n-3 HUFAs, particularly EPA and DHA and a high n-3 / n-6 ratio, in human diets are beneficial for various aspects of human health including preventive or protective effects in coronary heart disease, rheumatoid arthritis, cancer, neurodevelopmental and mood disorders etc (De Deckere *et al.*, 1998; Horrocks & Yeo, 1999; Simopoulos, 1999; Hunter & Roberts, 2000; ISSFAL, 2000; Simopoulos, 2003). At present, intensive culture of Atlantic salmon uses marine FO resulting in a highly nutritious and healthy product, as it is rich in n-3 HUFAs and has a high n-3 / n-6 ratio (Bell *et al.*, 1998). However, in recent times there has been a desire to investigate more sustainable alternatives to fish meal and fish oil for use in aquaculture feeds. Clearly, any changes towards use of vegetable alternatives to marine FO should not be at the expense of the quality and nutritional value of the final product. In this regard the present study showing moderate reduction of EPA and DHA in fish fed diets containing RO, at levels as high as 60% even at low protein levels, could be significant, although it should be remembered that this was a relatively short trial compared to the whole production cycle for Atlantic salmon.

The total β -oxidation capacity, including mitochondrial and peroxisomal β -oxidation activity, was measured in liver and a combined red and white muscle fraction. No significant effects were observed either in liver or muscle. These results are in line with other studies which showed that when FO was replaced by RO or other VO, in diets for Atlantic salmon, β -oxidation capacity was not affected (Tocher *et al.*, 2003b; Stubhaug *et al.*, 2005).

The results of this study showed no negative effects on growth and feed conversion, no major detrimental effects on lipid and fatty acid metabolism in Atlantic

salmon and an enhanced protein sparing effect, when fish were fed with lower protein feeds where RO replaced FO up to 60% of the total oil. In conclusion, the results of this study suggest that more sustainable, lower protein diets, in which a high proportion of the dietary protein and lipid is of non-marine origin, with high rapeseed oil inclusion, can be used in Atlantic salmon culture at low water temperatures.

Acknowledgements

This work was carried out with support from BioMar AS. V. Karalazos was supported by the State Scholarships Foundation (IKY), Greece.

References

- Ackman, R.G. (1980) Fish Lipids, part 1 In *Advances in Fish Science and Technology* (Connell, J.J. ed.), pp. 86-103. Fishing News Books Ltd., Farnham, U.K.
- AOAC (1995) Official methods of analysis of the Association of Official Analytical Chemists International, 16th edn. Association of Official Analytical Chemists, Arlington, VA, USA.
- Azevedo, P.A., Leeson, S., Cho, C.Y. & Bureau, D.P. (2004) Growth and feed utilization of large size rainbow trout (*Oncorhynchus mykiss*) and Atlantic salmon (*Salmo salar*) reared in freshwater: diet and species effects, and responses over time. *Aquacult. Nutr.*, **10**, 401-411.
- Bell, J.G., Henderson, R.J., Tocher, D.R., McGhee, F., Dick, J.R., Porter, A., Smullen, R.P. & Sargent, J.R. (2002) Substituting fish oil with crude palm oil in the diet of Atlantic salmon (*Salmo salar*) affects muscle fatty acid composition and hepatic fatty acid metabolism. *J. Nutr.*, **132**, 222-230.
- Bell, J.G., McEvoy, J., Tocher, D.R., McGhee, F., Campbell, P.J. & Sargent, J.R. (2001) Replacement of fish oil with rapeseed oil in diets of Atlantic salmon

- 415 (*Salmo salar*) affects tissue lipid compositions and hepatocyte fatty acid
416 metabolism. *J. Nutr.*, **131**, 1535-1543.
- 417 Bell, J.G., McEvoy, J., Webster, J.L., McGhee, F., Millar, R.M. & Sargent, J.R.
418 (1998) Flesh lipid and carotenoid composition of Scottish farmed Atlantic
419 salmon (*Salmo salar*). *J. Agric. Food Chem.*, **46**, 119-127.
- 420 Bell, J.G., McGhee, F., Campbell, P.J. & Sargent, J.R. (2003a) Rapeseed oil as an
421 alternative to marine fish oil in diets of post-smolt Atlantic salmon (*Salmo*
422 *salar*): changes in flesh fatty acid composition and effectiveness of subsequent
423 fish oil "wash out". *Aquaculture*, **218**, 515-528.
- 424 Bell, J.G., McGhee, F., Dick, J.R. & Tocher, D.R. (2005) Dioxin and dioxin-like
425 polychlorinated biphenyls (PCBs) in Scottish farmed salmon (*Salmo salar*):
426 effects of replacement of dietary marine fish oil with vegetable oils.
427 *Aquaculture*, **243**, 305-314.
- 428 Bell, J.G., McVicar, A.H., Park, M.T. & Sargent, J.R. (1991) High dietary linoleic
429 acid affects the fatty acid compositions of individual phospholipids from
430 tissues of Atlantic salmon (*Salmo salar*): association with stress susceptibility
431 and cardiac lesion. *J. Nutr.*, **121**, 1163-1172.
- 432 Bell, J.G., Tocher, D.R., Henderson, R.J., Dick, J.R. & Crampton, V.O. (2003b)
433 Altered fatty acid compositions in Atlantic salmon (*Salmo salar*) fed diets
434 containing linseed and rapeseed oils can be partially restored by a subsequent
435 fish oil finishing diet. *J. Nutr.*, **133**, 2793-2801.
- 436 Bendiksen, E.A., Berg, O.K., Jobling, M., Arnesen, A.M. & Masoval, K. (2003)
437 Digestibility, growth and nutrient utilisation of Atlantic salmon parr (*Salmo*
438 *salar* L.) in relation to temperature, feed fat content and oil source.
439 *Aquaculture*, **224**, 283-299.

- 440 Bendiksen, E.A. & Jobling, M. (2003) Effects of temperature and feed composition on
441 essential fatty acid (n-3 and n-6) retention in Atlantic salmon (*Salmo salar* L.)
442 parr. *Fish Physiol. Biochem.*, **29**, 133-140.
- 443 Christie, W.W. (1982) Lipid Analyses, Vol. 2nd Edition, pp. 52-56. Pergamon Press,
444 Oxford England.
- 445 De Deckere, E.A., Korver, O., Verschuren, P.M. & Katan, M.B. (1998) Health aspects
446 of fish and n-3 polyunsaturated fatty acids from plant and marine origin. *Eur. J.*
447 *Clin. Nutr.*, **52**, 749-753.
- 448 Einen & Skrede (1998) Quality characteristics in raw and smoked fillets of Atlantic
449 salmon, *Salmo salar*, fed high-energy diets. *Aquacult. Nutr.*, **4**, 99-108.
- 450 Einen, O. & Roem, A.J. (1997) Dietary protein/energy ratios for Atlantic salmon in
451 relation to fish size: growth, feed utilization and slaughter quality. *Aquacult.*
452 *Nutr.*, **3**, 115-126.
- 453 FAO (2005) Oilseeds, oils and oilmeals. *Food Outlook*, **2**, 22-25.
- 454 Folch, J., Lees, M. & Sloane-Stanley, G.H. (1957) A simple method for the isolation
455 and purification of total lipides from animal tissues. *J. Biol. Chem.*, **226**, 497-
456 509.
- 457 Fonseca-Madrigal, J., Karalazos, V., Campbell, P.J., Bell, J.G. & Tocher, D.R. (2005)
458 Influence of dietary palm oil on growth, tissue fatty acid compositions, and
459 fatty acid metabolism in liver and intestine in rainbow trout (*Oncorhynchus*
460 *mykiss*). *Aquacult. Nutr.*, **11**, 241-250.
- 461 Froyland, L., Asiedu, D.K., Vaagenes, H., Garras, A., Lie, O., Totland, G.K. & Berge,
462 R.K. (1995) Tetradecylthioacetic acid incorporated into very low density
463 lipoprotein: changes in the fatty acid composition and reduced plasma lipids in
464 cholesterol-fed hamsters. *J. Lipid Res.*, **36**, 2529-2540.

- 465 Froyland, L., Madsen, L., Eckhoff, K.M., Lie, O. & Berge, R.K. (1998) Carnitine
466 palmitoyltransferase I, carnitine palmitoyltransferase II, and acyl-CoA oxidase
467 activities in Atlantic salmon (*Salmo salar*). *Lipids*, **33**, 923-930.
- 468 Guderley, H. (2004) Metabolic responses to low temperature in fish muscle. *Biol.*
469 *Rev.*, **79**, 409-427.
- 470 Halver, J.E. & Hardy, R.E. (2002) Nutrient flow and retention In *Fish Nutrition*, 3rd
471 edn, pp. 768-769. Academic Press, Elsevier Science, San Diego, California,
472 USA.
- 473 Hemre, G.I. & Sandnes, K. (1999) Effect of dietary lipid level on muscle composition
474 in Atlantic salmon *Salmo salar*. *Aquacult. Nutr.*, **5**, 9-16.
- 475 Hillestad, Johnsen, Austreng & Asgard (1998) Long-term effects of dietary fat level
476 and feeding rate on growth, feed utilization and carcass quality of Atlantic
477 salmon. *Aquacult. Nutr.*, **4**, 89-97.
- 478 Hillestad, M. & Johnsen, F. (1994) High-energy/low-protein diets for Atlantic
479 salmon: effects on growth, nutrient retention and slaughter quality.
480 *Aquaculture*, **124**, 109-116.
- 481 Hochachka, P.W. & Somero, G.N. (2002) *Biochemical adaptation: mechanism and*
482 *process in physiological evolution*, Oxford University Press US, New York.
- 483 Horrocks, L.A. & Yeo, Y.K. (1999) Health benefits of docosahexaenoic acid (DHA).
484 *Pharmacol. Res.*, **40**, 211-225.
- 485 Hunter, B.J. & Roberts, D.C.K. (2000) Potential impact of the fat composition of
486 farmed fish on human health. *Nutr. Res.*, **20**, 1047-1058.
- 487 ISSFAL (2000) ISSFAL statement on omega-3 Polyunsaturated fatty acids and heart
488 disease. Retrieved on 3/4/2005 from
489 http://www.issfal.org.uk/heart_statement.htm.

- 490 Jacobs, M., Ferrario, J. & Byrne, C. (2002a) Investigation of polychlorinated dibenzo-
 491 p-dioxins, dibenzo-p-furans and selected coplanar biphenyls in Scottish farmed
 492 Atlantic salmon (*Salmo salar*). *Chemosphere*, **47**, 183-191.
- 493 Jacobs, M.N., Covaci, A. & Schepens, P. (2002b) Investigation of selected persistent
 494 organic pollutants in farmed Atlantic salmon (*Salmo salar*), salmon
 495 aquaculture feed, and fish oil components of the feed. *Environ. Sci. Technol.*,
 496 **36**, 2797-2805.
- 497 Jobling, M. & Johansen, S.J.S. (2003) Fat distribution in Atlantic salmon *Salmo salar*
 498 L. in relation to body size and feeding regime. *Aquacult. Res.*, **34**, 311-316.
- 499 Ng, W.-K., Sigholt, T. & Gordon Bell, J. (2004) The influence of environmental
 500 temperature on the apparent nutrient and fatty acid digestibility in Atlantic
 501 salmon (*Salmo salar* L.) fed finishing diets containing different blends of fish
 502 oil, rapeseed oil and palm oil. *Aquacult. Res.*, **35**, 1228-1237.
- 503 NRC (1993) Nutrient requirements of fish. National Academy Press, Washington
 504 D.C.
- 505 Pike, I.H. & Barlow, S.M. (2003) Impact of fish farming on fish stocks. *Int. Aquafeed*
 506 *Dir.*, 24-29.
- 507 Rosenlund, G., Obach, A., Sandberg, M.G., Standal, H. & Tveit, K. (2001) Effect of
 508 alternative lipid sources on long-term growth performance and quality of
 509 Atlantic salmon (*Salmo salar* L.). *Aquacult. Res.*, **32**, 323-328.
- 510 Sargent, J.R. & Tacon, A.G. (1999) Development of farmed fish: a nutritionally
 511 necessary alternative to meat. *Proc. Nutr. Soc.*, **58**, 377-383.
- 512 Sargent, J.R., Tocher, D.R., Bell, J.G. & Halver, J.E.H.R.E. (2002) The lipids In *Fish*
 513 *Nutrition*, 3rd edn, pp. 181-257. Academic Press, Elsevier Science, San Diego,
 514 California, USA.

- 515 SCAN (2000) Dioxin contamination of feedingstuffs and their contribution to the
 516 contamination of food of animal origin. Opinion of the Scientific Committee
 517 on Animal Nutrition, adopted on November 6th 2000. European Commission
 518 for Health and Consumer Protection Directorate - General, Brussels, Belgium.
 519 Retrieved from http://www.europa.eu.int/comm/food/fs/sc/scan/out55_en.pdf
- 520 SCF (2001) Update of the Risk assessment of dioxins and dioxin-like PCBs in food
 521 based on new scientific information available since adoption of the SCF
 522 opinion of 22nd November 2000. Opinion of the Scientific Committee on
 523 Food, adopted on May 30th 2001. Brussels, Belgium, European Commission
 524 for Health and Consumer Protection Directorate-General. Retrieved from
 525 http://www.europa.eu.int/comm/food/fs/sc/scf/out90_en.pdf.
- 526 Simopoulos, A.P. (1999) Essential fatty acids in health and chronic disease. *Am. J.*
 527 *Clin. Nutr.*, **70**, 560S-569S.
- 528 Simopoulos, A.P. (2003) Omega-3 fatty acids and cancer. *Indoor and Built*
 529 *Environment*, **12**, 405-412.
- 530 Solberg, C. (2004) Influence of dietary oil content on the growth and chemical
 531 composition of Atlantic salmon (*Salmo salar*). *Aquacult. Nutr.*, **10**, 31-37.
- 532 Stubhaug, I., Froyland, L. & Torstensen, B.E. (2005) β -Oxidation capacity of red and
 533 white muscle and liver in Atlantic salmon (*Salmo salar* L.) - effects of
 534 increasing dietary rapeseed oil and olive oil to replace capelin oil. *Lipids*, **40**,
 535 39-47.
- 536 Tacon, A.G.J. (2004) Use of fish meal and fish oil in aquaculture: a global
 537 perspective. *Aquatic Resources, Culture and Development*, **1**, 3-14.

- 538 Tidwell, J.H. & Allan, G.L. (2002) Fish as food: aquaculture's contribution.
 539 Ecological and economic impacts and contributions of fish farming and capture
 540 fisheries. *World Aquaculture*, **33**, 44-48.
- 541 Tocher, D.R., Bell, J.G., Dick, J.R. & Crampton, V.O. (2003a) Effects of dietary
 542 vegetable oil on Atlantic salmon hepatocyte fatty acid desaturation and liver
 543 fatty acid compositions. *Lipids*, **38**, 723-732.
- 544 Tocher, D.R., Bell, J.G., Dick, J.R., Henderson, R.J., McGhee, F., Michell, D. &
 545 Morris, P.C. (2000) Polyunsaturated fatty acid metabolism in Atlantic salmon
 546 (*Salmo salar*) undergoing parr-smolt transformation and the effects of dietary
 547 linseed and rapeseed oils. *Fish Physiol. Biochem.*, **23**, 59-73.
- 548 Tocher, D.R., Bell, J.G., MacGlaughlin, P., McGhee, F. & Dick, J.R. (2001)
 549 Hepatocyte fatty acid desaturation and polyunsaturated fatty acid composition
 550 of liver in salmonids: effects of dietary vegetable oil. *Comp. Biochem. Physiol.*
 551 *B*, **130**, 257-270.
- 552 Tocher, D.R., Bell, J.G., McGhee, F., Dick, J.R. & Fonseca-Madriral, J. (2003b)
 553 Effects of dietary lipid level and vegetable oil on fatty acid metabolism in
 554 Atlantic salmon (*Salmo salar* - L.) over the whole production cycle. *Fish*
 555 *Physiol. Biochem.*, **29**, 193-209.
- 556 Tocher, D.R., Fonseca-Madriral, J., Dick, J.R., Ng, W.-K., Bell, J.G. & Campbell,
 557 P.J. (2004) Effects of water temperature and diets containing palm oil on fatty
 558 acid desaturation and oxidation in hepatocytes and intestinal enterocytes of
 559 rainbow trout (*Oncorhynchus mykiss*). *Comp. Biochem. Physiol. B*, **137**, 49-63.
- 560 Tocher, D.R. & Harvie, D.G. (1988) Fatty acid compositions of the major
 561 phosphoglycerides from fish neural tissues; (n-3) and (n-6) polyunsaturated

- 562 fatty acids in rainbow trout (*Salmo gairdneri*) and cod (*Gadus morhua*) brains
563 and retinas. *Fish Physiol. Biochem.*, **5**, 229-239.
- 564 Torstensen, B.E., Froyland, L. & Lie, O. (2004a) Replacing dietary fish oil with
565 increasing levels of rapeseed oil and olive oil - effects on Atlantic salmon
566 (*Salmo salar* L.) tissue and lipoprotein lipid composition and lipogenic enzyme
567 activities. *Aquacult. Nutr.*, **10**, 175-192.
- 568 Torstensen, B.E., Froyland, L., Ornsrud, R. & Lie, O. (2004b) Tailoring of a
569 cardioprotective muscle fatty acid composition of Atlantic salmon (*Salmo*
570 *salar*) fed vegetable oils. *Food Chem.*, **87**, 567-580.
- 571 Torstensen, B.E., Lie, O. & Froyland, L. (2000) Lipid metabolism and tissue
572 composition in Atlantic salmon (*Salmo salar* L.) - effects of capelin oil, palm
573 oil, and oleic acid-enriched sunflower oil as dietary lipid sources. *Lipids*, **35**,
574 653-664.
- 575 U. S. Department for Agriculture (2005) Vegetable oil production, consumption and
576 imports for top 10 consuming markets - Circular October 2005. Retrieved on
577 11/11/2005 from [http://www.fas.usda.gov/oilseeds/circular/2005/05-](http://www.fas.usda.gov/oilseeds/circular/2005/05-10/table9.pdf)
578 [10/table9.pdf](http://www.fas.usda.gov/oilseeds/circular/2005/05-10/table9.pdf)
- 579 Zar, J.H. (1996) Biostatistical analysis. Prentice-Hall International Editions, London,
580 U.K.
- 581
- 582

Table 1 Feed components and proximate compositions (g kg⁻¹) of the experimental diets

	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60
<i>Components</i>						
Fishmeal	390	390	390	310	310	310
Oil seed and legume seed meals	225	225	225	235	235	235
South American fish oil	280	196	112	318	222	126
Rapeseed oil ^a	0	84	168	0	96	192
Binder	120	120	120	130	130	130
Premixes ^b	18	18	18	23	23	23
<i>Analysed composition ^c</i>						
Moisture	43	47	44	67	58	56
Protein	387	388	391	344	347	342
Lipid	327	323	321	353	347	361
Ash	79	78	78	70	70	69
Gross Energy, kJ g ⁻¹ ^d	25.2	25.1	25.2	25.3	25.6	25.5
Protein/Energy ratio ^e	15.3	15.5	15.3	13.6	13.4	13.2

^a Double-low quality rapeseed oil

^b Vitamin and mineral premixes prepared according to BioMar A/S commercial standards.
Includes crystalline amino acids and Carophyl pink to provide 40mg/kg astaxanthin.

^c Wet weight

^d Estimated from caloric values of 39.5, 23.6 and 17.2 kJ g⁻¹ for fat, protein and carbohydrate, respectively

^e Calculated g protein kJ⁻¹

Table 2 Fatty acid compositions (% by weight of total fatty acids) of the experimental diets

Fatty Acid	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60
14:0	5.9	3.5	1.8	5.4	3.1	1.9
16:0	20.3	15.2	10.4	19.6	14.3	10.4
18:0	3.6	3.0	2.5	3.5	2.9	2.5
Total saturated ^a	31.1	23.1	15.9	29.9	21.6	16.4
16:1n-7	6.6	4.2	2.3	6.1	4.2	2.2
18:1n-9	13.4	29.0	42.5	13.2	28.0	42.9
18:1n-7	2.5	2.8	3.0	2.5	3.1	2.6
20:1n-9	4.4	3.5	2.6	4.6	3.4	2.5
22:1	6.5	4.8	2.5	6.7	4.3	2.5
24:1n-9	0.8	0.6	0.4	0.9	0.9	0.5
Total monoenes ^b	34.6	45.1	53.4	34.7	44.5	53.4
18:2n-6	3.7	9.4	14.5	3.5	9.4	14.3
20:2n-6	0.2	0.2	0.1	0.3	0.2	0.1
20:4n-6	0.5	0.3	0.2	0.6	0.4	0.2
22:5n-6	0.3	0.2	0.1	0.3	0.2	0.1
Total n-6 PUFA ^c	5.0	10.4	15.1	5.0	10.4	14.9
18:3n-3	1.3	3.7	5.8	1.4	3.9	5.9
18:4n-3	2.9	1.8	1.0	3.0	2.0	1.0
20:4n-3	0.7	0.4	0.2	0.7	0.5	0.3
20:5n-3	10.4	6.7	3.6	11.0	7.4	3.5
22:5n-3	2.0	0.7	0.4	1.2	0.8	0.3
22:6n-3	11.9	8.0	4.6	12.8	8.7	4.3
Total n-3 PUFA ^d	29.3	21.4	15.5	30.3	23.4	15.3
Total PUFA	34.3	31.8	30.7	35.4	33.8	30.2
(n-3) / (n-6)	5.9	2.1	1.0	6.0	2.2	1.0

^aIncludes 15:0, 20:0 & 22:0.^bIncludes 16:1n-9 & 20:1n-7.^cIncludes 18:3n-6, 20:3n-6 & 22:4n-6.^dIncludes 20:3n-3 & 22:4n-3.

Table 3 Growth and performance of Atlantic salmon fed the experimental diets for 12 weeks

Parameter	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60	TWO WAY ANOVA <i>P</i>		
							protein	oil	prot x oil
Initial Weight (g)	1168.4 ± 32.3	1184.6 ± 16.1	1152.4 ± 21.0	1162.8 ± 24.4	1171.9 ± 14.6	1168.4 ± 24.4			
Final Weight (g)	1711.3 ± 79.8	1772.0 ± 36.5	1784.3 ± 48.5	1721.7 ± 47.6	1760.3 ± 27.0	1767.7 ± 57.6	0.812	0.149	0.894
Final Length (cm)	50.5 ± 0.4	50.9 ± 1.1	50.7 ± 1.0	50.7 ± 0.3	49.9 ± 0.6	51.8 ± 0.6	0.829	0.130	0.077
Mortalities ¹	5	1	2	3	0	5			
FCR	0.86 ± 0.01	0.84 ± 0.02	0.81 ± 0.01	0.86 ± 0.02	0.87 ± 0.04	0.85 ± 0.03	0.077	0.160	0.349
SGR	0.49 ± 0.03	0.52 ± 0.01	0.56 ± 0.01	0.50 ± 0.02	0.52 ± 0.01	0.53 ± 0.03	0.723	0.002	0.128
TGC	4.45 ± 0.27	4.75 ± 0.16	5.13 ± 0.11	4.59 ± 0.20	4.78 ± 0.07	4.87 ± 0.28	0.742	0.005	0.227
PRE	42.32 ± 3.45	42.55 ± 1.16	44.03 ± 1.44	45.56 ± 1.85	46.33 ± 1.63	46.08 ± 5.26	0.045	0.799	0.868
K (%)	1.31 ± 0.01	1.30 ± 0.02	1.32 ± 0.04	1.31 ± 0.06	1.30 ± 0.01	1.34 ± 0.01	0.609	0.274	0.941
VSI (%)	12.03 ± 1.00	12.01 ± 0.69	12.41 ± 0.98	12.57 ± 0.27	11.67 ± 0.28	11.90 ± 0.49	0.749	0.515	0.395
HSI (%)	1.34 ± 0.12	1.32 ± 0.02	1.45 ± 0.20	1.36 ± 0.10	1.36 ± 0.07	1.31 ± 0.07	0.646	0.822	0.371
CSI (%)	0.15 ± 0.03	0.13 ± 0.01	0.13 ± 0.00	0.13 ± 0.01	0.14 ± 0.03	0.12 ± 0.01	0.358	0.556	0.522

All values are mean ± S.D. (n=3).

¹ Total number

Table 4 Proximate composition (g kg⁻¹ of wet weight) of whole body from Atlantic salmon fed the experimental diets for 12 weeks

	<i>Start</i> ¹	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60	TWO WAY ANOVA <i>P</i>		
								protein	oil	prot x oil
Moisture	666.1 ± 16.5	664.7 ± 12.6	661.7 ± 12.9	665.3 ± 11.5	666.5 ± 6.2	665.8 ± 13.9	650.8 ± 10.9	0.602	0.522	0.350
Protein	176.4 ± 2.2	162.2 ± 4.9	162.1 ± 4.8	162.9 ± 2.5	162.2 ± 2.5	162.2 ± 4.2	162.3 ± 2.0	0.925	0.976	0.983
Lipid	128.9 ± 16.1	150.3 ± 10.8	155.1 ± 6.9	151.9 ± 11.9	145.2 ± 6.7	149.5 ± 13.2	164.5 ± 10.5	0.901	0.250	0.259
Ash	17.5 ± 0.7	14.9 ± 0.3	15.4 ± 1.0	15.0 ± 0.4	16.0 ± 0.7	15.4 ± 1.2	15.1 ± 1.0	0.376	0.672	0.421

All values are mean ± S.D. (n=3).

¹Values not included in the two-way ANOVA

Table 5 Total lipid (mg lipid g⁻¹ tissue) and fatty acid compositions (% by weight of total fatty acids) of muscle from Atlantic salmon fed the experimental diets for 12 weeks

	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60	TWO WAY ANOVA <i>P</i>		
							protein	oil	prot x oil
Total Lipid	117.8 ± 8.5	101.9 ± 6.4	102.8 ± 4.1	92.6 ± 13.9	101.7 ± 17.1	110.0 ± 9.6	0.259	0.748	0.057
<i>Fatty acid</i>									
14:0	4.7 ± 0.4	4.1 ± 0.1	3.4 ± 0.3	5.1 ± 0.4	4.3 ± 0.1	3.6 ± 0.4	0.064	0.000	0.739
16:0	16.0 ± 0.8	15.2 ± 0.9	13.5 ± 1.3	16.2 ± 1.4	14.6 ± 1.1	13.3 ± 1.1	0.649	0.004	0.857
18:0	3.1 ± 0.2	3.1 ± 0.3	3.1 ± 0.4	3.1 ± 0.3	2.9 ± 0.3	2.8 ± 0.3	0.299	0.635	0.825
Total saturated ¹	24.5 ± 1.4	23.2 ± 1.5	21.0 ± 2.5	25.3 ± 2.4	22.7 ± 1.7	20.8 ± 2.1	0.971	0.015	0.874
16:1n-7	6.6 ± 0.1	5.5 ± 0.2	4.4 ± 0.1	6.8 ± 0.0	5.6 ± 0.1	4.6 ± 0.1	0.019	0.000	0.474
18:1n-9	15.7 ± 0.5	23.3 ± 1.1	29.7 ± 0.9	15.9 ± 0.3	22.9 ± 0.4	27.9 ± 1.9	0.197	0.000	0.340
18:1n-7	3.2 ± 0.1	3.4 ± 0.1	3.1 ± 0.1	3.2 ± 0.2	3.3 ± 0.1	3.1 ± 0.3	0.766	0.070	0.613
20:1n-9	7.8 ± 0.5	7.2 ± 0.3	6.7 ± 0.2	8.0 ± 0.2	7.2 ± 0.2	6.8 ± 0.2	0.577	0.000	0.968
22:1	8.3 ± 0.4	7.4 ± 0.5	6.0 ± 0.6	8.5 ± 0.1	7.3 ± 0.2	6.4 ± 0.4	0.573	0.000	0.566
24:1n-9	0.8 ± 0.0	0.8 ± 0.0	0.7 ± 0.1	0.9 ± 0.1	0.8 ± 0.0	0.7 ± 0.0	0.407	0.002	0.531
Total monoenes ²	42.9 ± 1.2	48.0 ± 1.4	51.0 ± 0.5	43.8 ± 0.3	47.4 ± 0.2	49.9 ± 1.0	0.426	0.000	0.270
18:2n-6	3.7 ± 0.1	6.3 ± 0.1	8.2 ± 0.9	3.6 ± 0.1	6.3 ± 0.5	8.2 ± 0.6	0.922	0.000	0.942
20:2n-6	0.4 ± 0.0	0.5 ± 0.0	0.8 ± 0.2	0.4 ± 0.0	0.5 ± 0.0	0.6 ± 0.1	0.179	0.000	0.076
20:3n-6	0.1 ± 0.0	0.1 ± 0.0	0.1 ± 0.0	0.2 ± 0.0	0.1 ± 0.0	0.1 ± 0.0	0.870	0.274	0.481
20:4n-6	0.5 ± 0.0	0.3 ± 0.0	0.3 ± 0.0	0.4 ± 0.0	0.3 ± 0.0	0.3 ± 0.0	0.843	0.000	0.756
Total n-6 PUFA ³	5.1 ± 0.1	7.5 ± 0.2	9.6 ± 0.8	5.0 ± 0.2	7.6 ± 0.6	9.5 ± 0.7	0.817	0.000	0.912
18:3n-3	1.2 ± 0.0	2.2 ± 0.1	3.0 ± 0.4	1.2 ± 0.1	2.2 ± 0.2	3.0 ± 0.3	0.743	0.000	0.968
18:4n-3	1.8 ± 0.1	1.3 ± 0.2	1.0 ± 0.1	1.8 ± 0.1	1.4 ± 0.1	1.1 ± 0.0	0.302	0.000	0.418
20:4n-3	1.5 ± 0.1	1.1 ± 0.1	0.9 ± 0.1	1.4 ± 0.1	1.2 ± 0.1	1.0 ± 0.0	0.936	0.000	0.166
20:5n-3	7.4 ± 0.4	5.4 ± 0.9	4.2 ± 0.4	7.2 ± 0.7	5.7 ± 0.3	4.6 ± 0.2	0.485	0.000	0.679
22:5n-3	2.5 ± 0.1	1.9 ± 0.2	1.6 ± 0.2	2.5 ± 0.3	1.9 ± 0.2	1.7 ± 0.0	0.505	0.000	0.734
22:6n-3	12.9 ± 0.9	9.3 ± 1.2	7.5 ± 0.7	11.9 ± 1.1	9.7 ± 0.5	8.2 ± 0.1	0.898	0.000	0.252
Total n-3 PUFA ⁴	27.5 ± 1.1	21.4 ± 2.6	18.4 ± 1.9	26.0 ± 2.3	22.3 ± 1.3	19.8 ± 0.5	0.668	0.000	0.422
Total PUFA	32.6 ± 1.2	28.8 ± 2.8	28.0 ± 2.7	31.0 ± 2.6	29.8 ± 1.9	29.3 ± 1.2	0.766	0.081	0.535
(n-3) / (n-6)	5.4 ± 0.1	2.9 ± 0.3	1.9 ± 0.1	5.2 ± 0.2	2.9 ± 0.1	2.1 ± 0.1	0.557	0.000	0.263

Values are mean ± S.D. (n=3).

¹Includes 15:0, 20:0 & 22:0.²Includes 16:1n-9 & 20:1n-7.³Includes 18:3n-6, 20:3n-6 & 22:4n-6.⁴Includes 20:3n-3 & 22:4n-3.

Table 6 Total lipid (mg lipid g⁻¹ tissue) and fatty acid compositions (% by weight of total fatty acids) of liver from Atlantic salmon fed the experimental diets for 12 weeks

	HP-R0	HP-R30	HP-R60	LP-R0	LP-R30	LP-R60	TWO WAY ANOVA <i>P</i>		
							protein	oil	prot x oil
Total lipid	62.7 ± 16.6	49.4 ± 6.6	81.0 ± 37.0	52.5 ± 16.7	54.9 ± 7.9	69.3 ± 16.2	0.422	0.037	0.519
<i>Fatty acid</i>									
14:0	2.4 ± 0.4	1.8 ± 0.0	1.2 ± 0.1	2.7 ± 0.3	1.8 ± 0.3	1.2 ± 0.1	0.543	0.000	0.714
16:0	15.2 ± 1.7	12.6 ± 1.1	10.0 ± 0.9	14.2 ± 1.0	13.2 ± 1.1	9.1 ± 1.0	0.418	0.000	0.450
18:0	6.1 ± 0.6	5.1 ± 0.3	4.8 ± 0.5	5.5 ± 0.1	4.9 ± 0.5	4.0 ± 0.2	0.022	0.000	0.537
Total saturated ¹	24.4 ± 2.5	20.1 ± 1.4	16.5 ± 1.4	23.2 ± 1.2	20.4 ± 1.2	15.1 ± 1.5	0.359	0.000	0.611
16:1n-7	3.7 ± 0.5	2.6 ± 0.2	1.9 ± 0.2	4.1 ± 0.2	2.6 ± 0.3	1.9 ± 0.2	0.327	0.000	0.268
18:1n-9	15.9 ± 4.0	23.0 ± 3.9	35.7 ± 3.9	16.7 ± 2.0	21.3 ± 2.1	37.1 ± 4.8	0.928	0.000	0.729
18:1n-7	3.1 ± 0.3	2.9 ± 0.2	3.1 ± 0.1	3.4 ± 0.3	3.0 ± 0.3	3.2 ± 0.5	0.234	0.300	0.946
20:1n-9	4.4 ± 0.6	4.2 ± 0.8	5.0 ± 0.2	4.7 ± 0.3	4.0 ± 0.5	5.3 ± 0.7	0.692	0.025	0.615
22:1	1.4 ± 0.2	1.1 ± 0.1	0.9 ± 0.0	2.1 ± 0.4	1.4 ± 0.3	1.0 ± 0.0	0.007	0.000	0.238
24:1n-9	1.0 ± 0.3	0.9 ± 0.2	0.6 ± 0.2	0.9 ± 0.1	1.1 ± 0.1	0.7 ± 0.1	0.737	0.011	0.299
Total monoenes ²	29.8 ± 5.2	35.0 ± 5.1	47.3 ± 4.0	32.1 ± 2.8	33.6 ± 3.4	49.3 ± 6.2	0.661	0.000	0.746
18:2n-6	2.1 ± 0.4	4.9 ± 0.5	8.2 ± 0.4	2.5 ± 0.3	4.9 ± 0.5	9.0 ± 0.5	0.124	0.000	0.396
20:2n-6	0.6 ± 0.1	1.3 ± 0.2	2.1 ± 0.1	0.6 ± 0.1	1.2 ± 0.1	2.1 ± 0.2	0.695	0.000	0.600
20:3n-6	0.1 ± 0.0	0.1 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.2 ± 0.0	0.853	0.001	0.215
20:4n-6	2.2 ± 0.5	1.9 ± 0.4	1.0 ± 0.3	1.9 ± 0.2	2.0 ± 0.3	1.0 ± 0.2	0.524	0.000	0.413
Total n-6 PUFA ³	5.7 ± 0.1	8.6 ± 0.4	11.9 ± 0.3	5.7 ± 0.3	8.7 ± 0.1	12.5 ± 0.4	0.100	0.000	0.094
18:3n-3	0.7 ± 0.1	1.6 ± 0.2	2.7 ± 0.1	0.9 ± 0.1	1.6 ± 0.2	2.9 ± 0.1	0.047	0.000	0.242
18:4n-3	0.3 ± 0.1	0.2 ± 0.0	0.1 ± 0.0	0.4 ± 0.1	0.2 ± 0.0	0.2 ± 0.0	0.032	0.000	0.879
20:4n-3	1.4 ± 0.2	1.1 ± 0.1	0.8 ± 0.0	1.7 ± 0.1	1.1 ± 0.1	0.8 ± 0.0	0.208	0.000	0.096
20:5n-3	10.4 ± 1.8	8.8 ± 1.3	5.2 ± 0.9	10.3 ± 0.8	9.4 ± 0.8	4.9 ± 1.4	0.921	0.000	0.808
22:5n-3	3.3 ± 0.7	2.9 ± 0.3	1.5 ± 0.2	3.7 ± 0.3	2.5 ± 0.2	1.4 ± 0.3	0.671	0.000	0.370
22:6n-3	23.7 ± 4.6	21.2 ± 3.4	13.4 ± 3.1	21.9 ± 2.1	22.0 ± 2.0	12.1 ± 3.7	0.621	0.000	0.763
Total n-3 PUFA ⁴	40.2 ± 5.9	36.3 ± 4.1	24.3 ± 4.2	39.0 ± 3.1	37.3 ± 2.3	23.0 ± 5.2	0.814	0.000	0.877
Total PUFA	45.9 ± 6.0	45.0 ± 3.7	36.2 ± 4.2	44.7 ± 3.1	46.0 ± 2.1	35.5 ± 4.8	0.899	0.002	0.893
(n-3) / (n-6)	7.0 ± 1.0	4.2 ± 0.6	2.1 ± 0.4	6.9 ± 0.7	4.3 ± 0.3	1.8 ± 0.5	0.757	0.000	0.911

Values are mean ± S.D. (n=3).

¹Includes 15:0, 20:0 & 22:0.

²Includes 16:1n-9 & 20:1n-7.

³Includes 18:3n-6, 20:3n-6 & 22:4n-6.

⁴Includes 20:3n-3 & 22:4n-3.

Table 7 Total β -oxidation capacity (pmol/min/mg protein) of muscle and liver from Atlantic salmon fed the experimental diets for 12 weeks

Tissue	HP-R0		HP-R30		HP-R60		LP-R0		LP-R30		LP-R60		TWO WAY ANOVA <i>P</i>		
													protein	oil	prot x oil
Muscle [†]	0.49	± 0.02	0.61	± 0.10	0.57	± 0.12	0.48	± 0.09	0.45	± 0.07	0.49	± 0.07	0.062	0.575	0.340
Liver	3.28	± 1.19	4.72	± 1.33	5.13	± 1.77	3.86	± 0.16	4.09	± 0.61	4.42	± 0.79	0.631	0.196	0.544

Values are mean \pm S.D. (n=3).

[†]Includes red and white muscle