

1 **Title:** Short term changes in moisture content drive strong changes in Normalized Difference  
2 Vegetation Index and gross primary productivity in four Arctic moss communities

3  
4 Jeremy L. May<sup>1</sup>, Thomas Parker<sup>2,3</sup>, Steven Unger<sup>1,2</sup>, and Steven F. Oberbauer<sup>1</sup>

5 <sup>1</sup>Florida International University, 11200 SW 8<sup>th</sup> Street, Miami, FL 33199, USA

6 <sup>2</sup>The Ecosystems Center, Marine Biological Laboratory, 7 MBL St, Woods Hole, MA 02543, USA

7 <sup>3</sup>Biological and Environmental Sciences, School of Natural Sciences, University of Stirling,  
8 Stirling, UK, FK9 4LA.

9  
10  
  
Accepted refereed manuscript of:

May JL, Parker T, Unger S & Oberbauer SF (2018) Short term changes in moisture content drive strong changes in Normalized Difference Vegetation Index and gross primary productivity in four Arctic moss communities, *Remote Sensing of Environment*, 212, pp. 114-120.

DOI: <https://doi.org/10.1016/j.rse.2018.04.041>.

© 2018, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International <http://creativecommons.org/licenses/by-nc-nd/4.0/>

## Abstract

Climate change is currently altering temperature and precipitation totals and timing in Arctic regions. Moss communities constitute much of the understory in Arctic vegetation, and as poikilohydric plants moss are highly sensitive to timing and duration of moisture levels. Here we investigate the role of moisture content on NDVI, red and near-infrared reflectance, and gross primary productivity (GPP) of two sphagnum and two pleurocarpus moss community types during two separate drying experiments. For both experiments, blocks of moss were collected near Imnavait Creek, Alaska, saturated to full water capacity, and then allowed to air dry before being re-saturated. Drying of blocks was conducted in a translucent outdoor tent during the first experiment and under indoor climate-controlled conditions during the second. Community NDVI (experiment 1 and 2), and GPP (experiment 2) were measured at regular intervals during the dry-down and after rewetting. In both experiments, moss NDVI sharply declined between 80% and 70% moisture content for sphagnum moss communities (NDVI change = -0.17 to -0.2), but less so for pleurocarpus moss communities (NDVI change = -0.06 to -0.12). Changes in NDVI were largely the result of increases in reflectance in red wavelengths. Peak GPP for all community types in the second experiment ( $1.31$  to  $2.08 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) occurred at 80% moisture content and declined significantly as moisture content decreased. Rates of GPP continued to decline below 80% moisture content until near zero as moss reached a steady weight (air dry) over a period of 84 hours, while NDVI values declined slowly between 70% hydration and fully air dry. Re-saturation caused NDVI to increase in both sphagnum (NDVI change = +0.18 to +0.23) and pleurocarpus (NDVI change = +0.10 to +0.17) communities. Only sphagnum communities showed GPP resuming ( $0.824 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) after 24 hours. The strong

33 changes in NDVI and mismatch of moss NDVI values and GPP with moisture content  
34 fluctuations indicate that using NDVI as a proxy for productivity in Arctic vegetation  
35 communities may be problematic and underscores the need for quantification of moss  
36 community coverage, composition, and moisture content.

37

38

## Introduction

Arctic regions have experienced significant warming over the past several decades (Overland 2002, Chapin et al. 2005, IPCC 2013) with substantial ecosystem consequences (Hinzman et al. 2005). In some regions of the Arctic, changes in precipitation patterns, increased evapotranspiration, and falling water tables associated with climate change have reduced pond size and number (Riordan et al. 2006, Andreson and Loughheed 2015) and affected soil moisture available to tundra plants (Roulet et al. 1992, IPCC 2013). While precipitation models for the Arctic generally suggest an increase in precipitation, most of this increase is distributed over winter and fall (as snow), while summer would remain relatively unchanged (Kattsov et al. 2007). Arctic plant communities are shifting to vegetation types and resulting carbon dynamics that reflect drier conditions (Oechel et al. 1992, Chapin et al. 1995, Mack et al. 2011). Mosses constitute an important component of Arctic vegetation communities, particularly in the understory, and may contribute substantially to ecosystem carbon fluxes (Shaver and Chapin 1991, Douma et al. 2007, Campioli et al. 2009, Turestky et al. 2010, Olivas et al. 2011, Zona et al. 2011). As a result of warming, recent vegetation measurements in the Arctic have shown a decrease in moss cover (Chapin et al. 1995, Molau and Alattalo 1998, Elmendorf et al. 2012, Hollister et al. 2015, Hobbie et al. 2017).

Decreases in available water have a particularly marked effect on the carbon balance in moss-dominated communities (Titus et al. 1983, Rydin and McDonald 1985, Alm et al. 1999, Komulainen et al. 1999). As poikilohydric plants, mosses do not have the ability to actively control water loss and therefore are highly susceptible to changes in water availability (Van Breemen 1995, Proctor and Tuba 2002). Consequently the moisture content of mosses varies

widely and rapidly compared to that of tundra vascular plants, and most mosses are more resilient to periodic drying, or even complete desiccation, than vascular plants (Levitt 1956, Proctor and Tuba 2002). Mosses have an optimal water content for peak photosynthetic rates that does not necessarily occur at full saturation (Ueno and Kanda 2006, Van Gaalen et al. 2007, Harris 2008). Photosynthetic rates of moss are particularly sensitive to drying because cellular water content is a crucial limiting factor in the light reactions (Skre and Oechel 1981). As water loss becomes more severe, photosynthesis declines significantly (Schipperges and Rydin 1998). Even after re-saturation there can be a substantial lag before activity resumes (Oliver and Bewley 1984, Green and Lange 1995, Charron and Quatrano 2009, de Carvalho et al. 2012). Multiple drying and wetting cycles have a negative effect on the photosynthesis of moss due to lengthy recovery times (McNeil and Waddington 2003). As a result, prolonged periods of warmer, drier conditions have the potential to adversely affect moss growth and photosynthesis (Potter et al. 1995, Dorrepaal et al. 2004, Proctor et al. 2007) and therefore ecosystem productivity (Turetsky et al. 2012).

Accompanying variation in moss moisture content are changes in apparent coloration of some mosses. For example, some sphagnum species exhibit a markedly lighter appearance as a result of moisture loss (Van Breeman 1995), suggesting the possibility that desiccation may have implications for plant reflectance and remote sensing indices. Remote sensing has provided an effective method for determination of important environmental parameters on large spatial and temporal scales that, using conventional methods, would be otherwise cost and time prohibitive (Kerr and Ostrovsky, 2003). These resource limitations are exacerbated in Arctic regions as a result of the remoteness and scale of study areas. One metric commonly

used is the Normalized Difference Vegetation Index (NDVI) that takes advantage of the strong absorbance of the red and strong reflectance in the near-infrared region of the electromagnetic spectrum by green plants (Kriegler et al. 1969). Changes in growing season length and available moisture associated with climate change have been shown to alter remotely sensed NDVI, a measure strongly correlated with green biomass (Jia et al. 2003, Reidel et al., 2005, Gamon et al. 2013), shrub cover (Stow et al. 2007, Walker et al. 2012), and community productivity (Harris 2008). Increases in Arctic vegetation cover have been associated with increases in peak season NDVI measurements (Laidler et al. 2008, Chen et al. 2009, Kushida et al. 2009), particularly when moisture is high (Riedel et al. 2005, Huemmrich et al. 2010). However, if short-term changes in water content result in significant changes in reflectance of such an important ecosystem component as mosses, considerable uncertainty will be introduced into remote sensing-derived estimates of green biomass and productivity. The role of moisture content on photosynthesis and spectral reflectance has been investigated for a few moss species (Potter et al. 1995, Van Breeman 1995, Dorrepaal et al. 2003, Proctor et al. 2007); however no studies have addressed the role of water content on both of these properties simultaneously for different Arctic bryophyte communities

Here, we investigate how variation in plant water content affects NDVI, red and near-infrared reflectance, and gross primary productivity (GPP) of four moss communities through a full cycle from saturation, dry down, and re-saturation. We hypothesize that NDVI will be greatest at high but not fully saturated water contents and decrease with drying at rates specific to each community type. Re-saturation will quickly re-establish initial NDVI values. Gross Primary Productivity will peak at levels below full saturation, similar to those reported in

previous studies (Ueno and Kanda 2006, Van Gaalen et al. 2007, Harris 2008), and decrease strongly with drying. We also expect that air drying of moss to constant weight will cause a delay in re-establishment of initial GPP values after re-saturation.

## **Methods**

### ***Sample handling***

In two separate drying experiments, conducted in July (Exp 1) and August (Exp 2) 2016, monoliths of four moss communities were collected from the low Arctic tundra near Toolik Field Station (TFS) at Imnavait Creek, Alaska, USA (68.635° N, -149.349° W). The four communities included two sphagnum communities, >95% *Sphagnum angustifolium* (green in color), >95% *Sphagnum capilliofolium* (red in color), and two pleurocarpous communities, >95% *Hylocomium splendens* and a mixed community (~50% *Aulacomnium* spp., ~30% *Hylocomium splendens*, and ~10% *Polytrichum* spp.). Four replicate 20x20x8 cm (length x width x depth) blocks of each community were collected (16 total) for both experiments. Each replicate was prepared by removing vascular plants and soil prior to placing them in a tray of distilled water (3 cm depth) to hydrate. Moss blocks were soaked for two hours until they reached full saturation and then allowed to drain for one hour to remove excess water. The vertical faces of each moss block were wrapped in cellophane and then placed in a Styrofoam tray to prevent uneven drying from the sides.

During Exp1, blocks were allowed to dry gradually in a translucent white outdoor tent to allow temperatures (range 3-29°C) to track daily temperature changes. During Exp2, blocks were allowed to dry gradually in the TFS Incubation Facility maintained at 23°C to minimize

temperature variability during the drying process. This temperature was determined to be a suitable analog of natural peak season conditions (TFS Environmental Data Center, 2016). Drying in the incubation facility took place under constant lighting using Hydrofarm® 1000 watt lamps at a height of 1.5 m and  $400 \mu\text{mol m}^{-2} \text{s}^{-1}$  (Hydrofarm, Inc., Petaluma, California, USA). For both experiments, moss blocks reached ecologically dry status (air dry) after 84 (Exp2) and 96 (Exp1) hours and then were re-saturated with distilled water to determine drying resilience.

### **Measurements**

All measurements during the drying process were taken at approximately 12 h intervals. After rehydration, measurements were conducted at 4, 12 and 24 h intervals. Blocks were then dried further at 50°C for 48 hours to achieve 0% water content allowing for calculation of percent saturation at each measurement. To monitor water loss, block weight was measured to 1 mg using an electronic balance (Ohaus Corporation, Parsippany, New Jersey, USA) throughout the study period of both drying experiments. During Exp1, community reflectance (350 – 1100 nm) was measured in ambient light conditions using a single channel Unispec® (PP Systems, Amesbury Massachusetts, USA) at a height of 10 cm with NDVI calculated afterward (see below). During the second experiment, NDVI was measured using a Trimble self-illuminated, handheld GreenSeeker® crop sensing system (Trimble Navigation, Ltd., Sunnyvale, California, USA) at a height of 15 cm, the minimum recommended distance. The GreenSeeker® system with an internal light source was ideal for these measurements by providing accurate NDVI values despite being under artificial light conditions. NDVI for both experiments was calculated using the reflectance of near-infrared light ( $R_{774}$ ) and red light ( $R_{656}$ ) as  $(R_{774} - R_{656}) / (R_{774} + R_{656})$ . Community  $\text{CO}_2$  exchange for Exp2 was measured using a custom-made



transparent acrylic chamber (32 x 32 x 32 cm) with the moss block positioned on a hard flat surface that formed a gas-tight seal with the chamber using weather stripping. One 12V fan fixed inside the chamber insured full mixing of chamber air. The chamber was attached to a LI-6400XT Portable Photosynthesis System (LI-COR Inc., Lincoln, Nebraska, USA) in closed system mode. Gas exchange measurements were conducted under the Hydrofarm® 1000 watt lighting systems at  $900 \mu\text{mol m}^{-2} \text{s}^{-1}$ . A good seal between the chamber and the flat surface was determined if gas concentrations showed a steady rate of change with no fluctuations for 20 s. Following the methods of Shaver et al. (2007), when a stable change in  $\text{CO}_2$  concentrations was observed,  $\text{CO}_2$  concentration, chamber air temperature and PAR were logged every 2 s for 40 s. Flux ( $\mu\text{mol CO}_2 \text{s}^{-1}$ ) was calculated as a linear change in  $\text{CO}_2$  concentration over time multiplied by the air density ( $\text{mol m}^{-3}$ ). Flux was expressed on an ecosystem area basis using the moss surface area ( $0.04 \text{ m}^2$ ). For each moss block, measurements were taken in the light (Net Primary Production (NPP)) and in the dark under an opaque tarpaulin (Ecosystem Respiration (ER)). Gross Primary Production (GPP) was calculated as the difference between NPP and ER, assigning positive values to GPP and negative values to ER.

### **Data Analysis**


Replicates of moss GPP and NDVI measurements were grouped by measurement time and separately by 10% moisture content increments for statistical analysis. Time and moisture content groupings were compared using a repeated measures analysis of variance with Tukey's post-hoc analysis. In the second experiment, resilience of each moss community to re-hydration after drying was determined using a drying response index (DRI) calculated as the proportion of

NDVI or GPP measured after re-saturation ( $V_{\text{re-sat}}$ ) compared with the initial saturation values ( $V_{\text{initial}}$ ),  $\text{DRI} = -(V_{\text{re-sat}}/V_{\text{initial}})$ . All statistical tests were performed using the R statistical environment (R Core Team, Vienna, Austria).

## Results

During Exp1, all moss communities were fully air dried after 96 h with the largest changes in NDVI occurring between 12 and 24 h of drying (Figure 1). Values of NDVI began to decline with drying between 80 and 70% water content for all communities, but communities differed in the magnitude of NDVI change. The sphagnum communities, *S. capilliofolium* and *S. angustifolium*, showed the largest decline in NDVI with drying ( $-0.190$ ,  $p < 0.001$  and  $-0.230$ ,  $p < 0.001$  respectively). Mixed pleurocarpus and *H. splendens* communities decreased in NDVI to a lesser extent compared to the sphagnum communities but were still significant ( $-0.112$ ,  $p = 0.021$  and  $-0.140$ ,  $p < 0.001$  respectively). All community NDVI values rebounded to near initial saturation levels upon re-saturation.


Decreases in NDVI were largely driven by increases in reflectance of red light (Figure 1), with the largest increases in the two sphagnum communities (*S. capilliofolium*  $+0.058$   $p = 0.025$  and *S. angustifolium*  $+0.067$   $p = 0.017$ ) compared with *H. splendens* ( $+0.039$   $p = 0.231$ ) and the mixed pleurocarpus ( $+0.038$   $p = 0.234$ ). Near-infrared reflectance for all communities was mixed with drying; some communities increased and some decreased, but no changes were significant ( $0.009$ - $0.045$ ). Red and near-infrared reflectance returned to near initial saturation levels upon re-saturation after only a few minutes.



**Figure 1:** NDVI (solid line), near-infrared (dotted line), and red (dashed line) of four communities by percent moisture content (left panels) and hours after initial saturation (right panels) during drying and after re-saturation during experiment 1.

In Exp2, all communities took approximately 84 hours to reach air dry and the lowest NDVI values (0.29 to 0.47, Figure 2). The largest decrease in NDVI observed as water contents declined occurred between 12 and 36 h post-saturation for all communities. *Sphagnum capilliofolium* and *H. splendens* communities had the highest NDVI measurements (0.70 and 0.64 respectively) at initial full saturation, while the *S. angustifolium* and mixed pleurocarpus were lower (0.55 and 0.57, Figure 2, Table 1). The NDVI of all community types was generally stable from 100% to 80% saturation, followed by an abrupt decline between 80 and 70%, after which there was steady but slow decline in NDVI to fully air dry. The largest decreases in NDVI between 80% and 70% saturation were found for the two sphagnum communities (*S.*

*capilliofolium* -0.19,  $P < 0.001$  and *S. angustifolium* -0.16,  $P < 0.001$ ), while the decreases for the mixed pleurocarpus and *H. splendens* communities were less, albeit still significant (-0.06,  $P = 0.038$  and -0.09,  $P = 0.014$  respectively). NDVI of all communities increased strongly upon re-saturation (+0.17 to +0.23, all  $p < 0.001$ ).




**Figure 2:** Gross primary productivity and NDVI of four communities by percent moisture content during drying and three measurement times after re-saturation during experiment 2.

**Table 1:** Gross primary productivity and NDVI of four communities by percent moisture content during drying and three measurement times after re-saturation compared using repeated measures analysis of variance with Tukey's post-hoc analysis, along with the initial saturated and final oven dry weight for each of the four community types during experiment 2. Letters denote statistically significant differences ( $p < 0.05$ ) between moisture content measurements.

	Moisture Content										ReWet (100% Moisture)			Initial	Dry
	100	99-90	89-80	79-70	59-50	49-40	39-30	29-20	19-10	9-0	Hours After			Saturated	Weight
											4	12	24	Weight	Weight
														(g)	(g)
Mixed Pleurocarpus	0.57 a	0.56 a	0.56 a	0.50 ab	0.47 b	0.45 b	0.46 b	0.45 b	0.45 b	0.44 b	0.55 a	0.56 a	0.56 a	578.5	113.8
<i>S. angustifolium</i>	0.55 a	0.54 a	0.51 a	0.35 b	0.32 b	0.32 b	0.30 b	0.30 b	0.30 b	0.29 b	0.47 a	0.43 a	0.43 a	584.0	101.5
<i>H. splendens</i>	0.64 a	0.63 a	0.62 a	0.51 b	0.49 b	0.48 b	0.48 b	0.47 b	0.47 b	0.47 b	0.62 a	0.61 a	0.60 a	403.3	116.2
<i>S. capillifolium</i>	0.70 a	0.68 a	0.67 a	0.46 b	0.45 b	0.45 b	0.45 b	0.38 c	0.40 bc	0.38 c	0.61 a	0.62 a	0.62 a	753.3	144.6
Gross Primary Productivity ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )															
Mixed Pleurocarpus	0.749 a	1.472 b	2.081 c	1.207 b	0.727 a	0.443 d	0.175 e	0.000 f	0.202 e	0.000 f	0.000 f	0.000 f	0.000 f		
<i>S. angustifolium</i>	0.195 a	0.970 b	1.671 c	0.913 b	0.352 ad	0.526 d	0.521 d	0.000 e	0.033 e	0.000 e	0.000 e	0.034 e	0.000 e		
<i>H. splendens</i>	0.883 a	1.652 b	2.042 c	1.608 b	0.002 d	0.000 d	0.374 e	0.000 d	0.206 e	0.111 d	0.000 d	0.296 e	0.000 d		
<i>S. capillifolium</i>	1.046 a	1.320 b	1.311 b	0.571 c	0.565 c	0.277 d	0.235 d	0.184 de	0.031 e	0.000 e	0.000 e	0.552 c	0.824 f		

219  
220 All moss communities were photosynthesizing at full water saturation (0.195 to 1.046  
221  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and initially increased as drying began (Figure 2, Table 1). *Sphagnum capilliofolium*  
222 community GPP peaked at approximately 90% saturation ( $1.320 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) while all other  
223 communities peaked at 80% saturation (mixed pleurocarpus  $1.332 \mu\text{mol m}^{-2} \text{s}^{-1}$ , *S.*  
224 *angustifolium*  $1.476 \mu\text{mol m}^{-2} \text{s}^{-1}$ , *H. splendens*  $1.159 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Rates of GPP for all  
225 communities decreased precipitously below 80% saturation with little or no GPP occurring at  
226 fully air dry (0 to  $0.111 \mu\text{mol m}^{-2} \text{s}^{-1}$ ). Rates of GPP peaked for all communities after 12 hours of  
227 drying following the initial saturation and then continued to decline until a total of 84 hours of  
228 drying. Twelve hours after re-saturation, *S. angustifolium* ( $0.034 \mu\text{mol m}^{-2} \text{s}^{-1}$ ), *H. splendens*  
229 ( $0.296 \mu\text{mol m}^{-2} \text{s}^{-1}$ ), and *S. capilliofolium* ( $0.552 \mu\text{mol m}^{-2} \text{s}^{-1}$ ) had regained some GPP (Figure 3).  
230 Twenty four hours post saturation only the *S. capilliofolium* community showed any GPP ( $0.824$   
231  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ).




**Figure 3:** Gross primary productivity and NDVI of four communities by hours of drying and three measurement times after re-saturation during experiment 2.

Measuring drying resilience by means of the DRI showed that NDVI and GPP decreased in all community types after re-saturation compared with the initial comparison (Figure 4).

Drying response index for NDVI of the mixed pleurocarpus and *H. splendens* communities

decreased only slightly ( $-0.023$ ,  $P = 0.101$  and  $-0.044$ ,  $P = 0.081$  respectively), returning to near original saturation values. The DRI for NDVI for both sphagnum communities had significant declines in response to full drying (*S. angustifolium*  $-0.197$ ,  $P = 0.013$  and *S. capilliofolium*  $-0.122$ ,  $P = 0.042$ ). The DRI for GPP of all communities decreased after drying with mixed pleurocarpus ( $-1.000$ ,  $P < 0.001$ ), *S. angustifolium* ( $-0.942$ ,  $P < 0.001$ ), and *H. splendens* ( $-0.888$ ,  $P < 0.001$ ) declining the most and the *S. capilliofolium* community being the most resilient to drying ( $-0.562$ ,  $P = 0.013$ ).



**Figure 4:** Drying response indexes of four communities for GPP and NDVI during experiment 2 compared using a one-way analysis of variance. Statistical significance denoted with \*.

## Discussion

Previous studies have shown that drying alters spectral reflectance (Riedel et al. 2005, Huemmrich et al. 2012) and moss productivity (Skre and Oechel 1981, Ueno and Kanda 2006,



Harris 2008). Here we show that the reflectance index, NDVI, and GPP decline strongly with moss desiccation, but they do not occur at the same rate and magnitude, resulting in a mismatch between NDVI levels and productivity. Reductions in NDVI of the moss communities with desiccation were very large, approaching 50% of maximum values and driven mostly by increases in red light reflectance during drying. With rewetting after the strong dry down, NDVI values were near peak levels while GPP was near zero.

#### ***Moss water content effect on NDVI values***

In recent decades, measured NDVI values have been increasing in the Arctic as temperatures warm and ecosystem productivity increases (Jia et al. 2003). These changes in peak season NDVI values, however, have not been increasing uniformly across spatial and temporal time scales, with evidence of a slowing of the rate of increase (Bhatt et al. 2013). The heterogeneity of changes in peak season NDVI values may be a result of the non-uniformity of the well-documented community dominance and moss decline in the Arctic (Shaver and Chapin 1991, Douma et al. 2007, Campioli et al. 2009, Elmendorf et al. 2012, Hollister et al. 2015). Moss communities are often an important component of Arctic understories and have been shown to play a large role in community production (Duoma et al. 2007, Campioli et al. 2009) and remotely-sensed spectral measurements (Walker et al 2003). Our results show that even small changes in the water content of the moss understory may play a role in the slowing of changes in landscape scale NDVI. Warmer, drier conditions during peak growing season may artificially decrease estimates of peak season landscape scale NDVI estimates.

These results have important implications for remote sensing of plant biomass and productivity in regions where mosses are important components of the vegetation. A general

assumption in the use of NDVI to estimate green biomass of plants is that NDVI is not strongly affected by short-term changes in leaf water content. While this assumption is generally the case for vascular plants, our results show that changes in moss water content can induce rapid and large changes in NDVI with no change in biomass. Furthermore, the relationship between NDVI and water content is markedly nonlinear. Variation in the water content of moss may be an important source of error in models using NDVI to estimate green biomass or leaf area (Oechel et al. 2000, Vourlitis et al. 2000, Shaver et al. 2007) that is then used in ecosystem photosynthesis models.

All of the moss communities in this study followed similar patterns of NDVI reductions with drying, although the magnitude of NDVI change in response to drying was community-specific. As predicted, all communities had the highest NDVI values at full, initial saturation (80-100% moisture content) with marked NDVI declines with drying. NDVI of all communities declined sharply over a relatively narrow range of water content from 80-70% moisture content, with the most substantial declines found for the two sphagnum communities. The lower NDVI values and higher levels of red reflectance at lower moisture content levels may act as a mechanism to minimize absorption of irradiance to prevent further evaporative water loss or cellular damage (Charron and Quatrano 2009, de Carvalho et al. 2012). The *H. splendens*-dominated and pleurocarpus mixed communities showed moderate increases in NDVI upon re-saturation. In contrast, both sphagnum communities had abrupt (<2 minutes), significant increases in NDVI values upon re-saturation. The rapidity of NDVI increases upon re-saturation of sphagnum communities suggests that changes in NDVI with drying and rehydration are in part a physical rather than biological response. Despite the rapid recovery of NDVI values of

sphagnum communities upon rewetting, values did not attain those of initial saturation, unlike the pleurocarpus mixed and *H. splendens* dominated communities that recovered fully. This lack of full rebound in sphagnum communities may be a result of, at least temporary, physiological damage occurring in response to desiccation to fully air dry (Oliver et al. 2005; Hájek and Beckett, 2008).

#### ***Moss water content effect on GPP***

Rapid changes in NDVI of moss communities with water content are associated with large changes in GPP, albeit nonlinearly. This variability is in addition to the already substantial difference in photosynthesis rates between vascular plants and mosses (Longton 1988). At Barrow, Alaska, production rates of mosses are on the order 10% of that of vascular plants (Oechel and Sveinbjornsson 1978), which means that photosynthesis per unit NDVI are very different for vascular plants compared to those of mosses. Our results show that moss communities may have relatively high NDVI values (0.55- 0.70), that if interpreted as vascular plant biomass would lead to large overestimates in productivity. These mismatches compromise the use of remotely-sensed NDVI data to estimate productivity in communities where mosses are abundant, but information on local moisture content or precipitation are lacking. Models using NDVI as a measure of productivity through estimating productivity by metric such as leaf area index (LAI) are highly effective across a range of spatial scales (Shaver et al. 2007, Loranty et al. 2011, Stoy et al. 2013). As spatial scale and vascular plant cover increases, the proportion of moss contribution to community spectral measurements is likely to decrease.

The magnitudes of changes in moss community GPP rates with drying were also community specific. However, all communities showed moderate rates at initial saturation and increased with drying to around 70-80% moisture content. This pattern of lower productivity at full saturation and increasing productivity after initial drying begins is similar to results found by Van Gaalen et al. (2007). A moderate amount of drying allows for air space within the plant while allowing cells to retain adequate moisture for full function. All communities had a peak GPP at 70-80% moisture content. Drying below 70-80% moisture content caused incremental decreases in GPP dropping to near zero in all of the communities when they reached air dry. Re-saturation had minimal effects on GPP, a finding consistent with previous findings that showed delayed recovery of moss physiological activity with rewetting after drying (Van Breeman 1995). Only the *S. capilliofolium* community showed a recovery of GPP during the 24 hours after re-saturation.

Moss communities such as those in this study are often intermixed at relatively small spatial scales across Arctic terrestrial ecosystems, implying a heterogeneous matrix of drying and recovery responses. While all four communities showed strong reduction in NDVI at the 80% drying threshold, the responses of both sphagnum communities were substantially greater than those in the pleurocarpus moss communities. To use remotely-sensed, reflectance-based productivity monitoring of Arctic ecosystems, further investigation is needed on the effects of intra-seasonal drying and rehydration on productivity and spectral reflectance of different moss communities.

These results in response in moss moisture content highlight the need for repeated remote sensing measurements over the same study regions with monitoring of a region's

recent precipitation events. Because of the remoteness and scale of Arctic regions, remotely sensed data are currently the best means to investigate seasonal productivity and vegetation composition shifts associated with climate change (Raynolds et al. 2008, Bhatt et al. 2010, Stow et al. 2007, Walker et al. 2012). This issue is crucial in Arctic regions where mosses comprise a major vegetation component, contribute substantially to ecosystem productivity (Olivas et al 2011), and are often a large component of total community reflectance (Hope et al. 1993). Our results show that periods of little or no precipitation combined with clear skies, high temperature and windy conditions have the potential to rapidly (<24 hours) lower moss water content sufficiently to reduce ecosystem NDVI values that would imply low predictions of ecosystem productivity even though vascular plant productivity may remain high. Remotely-sensed NDVI values measured for the same area shortly before and after a precipitation event may differ simply in response to moss moisture content.

Conditions conducive to moss desiccation are expected to increase with climate warming as temperatures increase, driving greater evapotranspiration. These changes will increase the frequency of moisture-induced changes in NDVI. Mosses grow in many different conditions ranging from on the surface of mineral soil or even on bare rock to areas that remain nearly continually wet or submerged. The frequency at which mosses desiccate is dependent in part on the microtopographic conditions where they are growing as well as weather conditions. Those growing on well-drained mineral soil or rock surfaces and hummocks are likely to desiccate frequently, whereas others may rarely if ever desiccate. Species colonizing conditions subject to frequent desiccation are likely to tolerate desiccation better than species in areas that rarely dry out (Longton 1988). Sites where mosses are continually wet are less likely to

show rapid NDVI changes in response to drying, but species from these conditions may be more susceptible to climate change-related drying in the long term.

## **Conclusion**

This study reinforces the importance of understanding the moisture content of moss when using remotely-sensed, reflectance techniques for monitoring productivity in Arctic terrestrial systems. Reflectance measures of different communities of moss revealed species-specific variation in response and resiliency to drying, therefore complicating the aggregation of moss as a uniform understory in Arctic ecosystems. At similar NDVI values, GPP varied depending on moss moisture content, demonstrating that moss NDVI is not an accurate proxy for physiological activity of some important Arctic mosses. This study underscores the need for monitoring and understanding the composition, spatial coverage, and moisture content of mosses for remote sensing-based monitoring of Arctic terrestrial ecosystems. Methodologies for remotely monitoring surface water content (e.g. Normalized Difference Water Index (NDWI) (Gao 1996), Normalized Difference Infrared Index (NDII) (Serrano et al. 2000), among others) are improving and could be useful for addressing these issues

## **Acknowledgements**

This work is based in part on support from National Science Foundation grants PLR-1432982, PLR-1417763, and PLR-1417645 for the support of this study. We thank the staff and management at Toolik Field Station and TFS Incubation Facility for support and logistics. Also, we would like thank Dr. Jessica Schedlbauer, Katherine Hood, Claudia Garcia-Estiu, and Salvatore Curasi for their insights and assistance during this study.

384

385 **Literature Cited**

- 386 Alm, J., L. Schulman, J. Walden, H. Nykänen, P. J. Martikainen and J. Silvola. 1999 Carbon  
387 balance of a boreal bog during a year with an exceptionally dry summer. *Ecology*.  
388 80:161–174.
- 389 Andresen, C. G. and V. L. Loughheed. 2015. Disappearing Arctic tundra ponds: Fine-scale analysis  
390 of surface hydrology in drained thaw lake basins over a 65 year period (1948–2013).  
391 *Journal of Geophysical Research: Biogeosciences*, 120(3): 466-479.
- 392 Bhatt, U. S., D. A. Walker, M. K. Reynolds, J. C. Comiso, H. E. Epstein, G. Jia, R. Gens, J. E. Pinzon,  
393 C. J. Tucker, C. E. Tweedie, and P. J. Webber, 2010. Circumpolar Arctic tundra vegetation  
394 change is linked to sea ice decline. *Earth Interactions*. 14(8): 1-20.
- 395 Bhatt, U.S., D. A. Walker, M. K. Reynolds, P. A. Bieniek, H. E. Epstein, J. C. Comiso, J. E. Pinzon, C.  
396 J. Tucker, I. V. Polyakov. 2013. Recent declines in warming and vegetation greening  
397 trends over pan-Arctic tundra. *Remote Sensing*. 5(9):4229-4254.
- 398 Campioli, M., R. Samson, A. Michelsen, S. Jonasson, R. Baxter, and R. Lemeur. 2009.  
399 Nonvascular contribution to ecosystem NPP in a subarctic heath during early and late  
400 growing season. *Plant Ecology* 202: 41–53.
- 401 Chapin, F. S., G. R. Shaver, A. E. Giblin, K. J. Nadelhoffer, J. A. and Laundre. 1995. Responses of  
402 arctic tundra to experimental and observed changes in climate. *Ecology*. 76(3): 694-711.
- 403 Chapin, F. S., M. Sturm, M. C. Serreze, J. P. McFadden, J. R. Key, A. H. Lloyd, A. D. McGuire, T. S.  
404 Rupp, A. H. Lynch, J. P. Schimel. and J. Beringe. 2005. Role of land-surface changes in  
405 Arctic summer warming. *Science*. 310(5748): 657-660
- 406 Charron, A. J. and R. S. Quatrano. 2009. Between a rock and a dry place: the water-stressed  
407 moss. *Molecular Plant*. 2(3): 478-486.
- 408 Chen, W., J. Li, Y. Zhang, F. Zhou, K. Khoeler, S. LeBlanc, R. Fraser, I. Olthof, Y. S. Zhang, and J.  
409 Wang. 2009. Relating biomass and leaf area index to non-destructive measurements in  
410 order to monitor changes in Arctic vegetation. *Arctic* 62:281-294.
- 411 de Carvalho, R. C., M. Catalá, J. M. da Silva, C. Branquinho. and E. Barreno. 2012. The impact of  
412 dehydration rate on the production and cellular location of reactive oxygen species in an  
413 aquatic moss. *Annals of Botany*. p.mcs. 180.
- 414 Dorrepaal, E., R. Aerts, J. H. Cornelissen, T. V. Callaghan, and R. S. Van Logtestijn. 2004. Summer  
415 warming and increased winter snow cover affect *Sphagnum fuscum* growth, structure  
416 and production in a sub-arctic bog. *Global Change Biology*. 10(1): 93-104.
- 417 Douma, J. C., M. T. Van Wijk, S. I. Lang, and G. R. Shaver. 2007. The contribution of mosses to  
418 the carbon and water exchange of arctic ecosystems: quantification and relationships  
419 with system properties. *Plant, Cell and Environment* 30: 1205–1215.

- Elmendorf, S. C., G. H. R. Henry, R. D. Hollister, R. G. Bjork, N. Boulanger-Lapointe, E. J. Cooper, J. H. C. Cornelissen, T. A. Day, E. Dorrepaal, T. G. Elumeeva, M. Gill, W. A. Gould, J. Harte, D. S. Hik, A. Hofgaard, D. R. Johnson, J. F. Johnstone, I. S. Jonsdottir, J. C. Jorgenson, K. Klanderud, J. A. Klein, S. Koh, G. Kudo, M. Lara, E. Levesque, B. Magnusson, J. L. May, J. A. Mercado-Diaz, A. Michelsen, U. Molau, I. H. Myers-Smith, S. F. Oberbauer, V. G. Onipchenko, C. Rixen, N. M. Schmidt, G. R. Shaver, M. J. Spasojevic, P. E. Porhallsdottir, A. Tolvanen, T. Troxler, C. E. Tweedie, S. Villareal, C.-H. Wahren, X. Walker, P. J. Webber, J. M. Welker, and S. Wipf. 2012. Plot-scale evidence of tundra vegetation change and links to recent summer warming. *Nature Climate Change* 2:453–457.
- Gamon, J. A., K. F. Huemmrich, R. S. Stone, C. E. Tweedie. 2013. Spatial and temporal variation in primary productivity (NDVI) of coastal Alaskan tundra: Decreased vegetation growth following earlier snowmelt. *Remote Sensing of Environment*. 129: 144-153.
- Gao, B.C., 1996. NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote sensing of environment*. 58(3):257-266.
- Green, T. G. A. and O. L. Lange. 1995. Photosynthesis in poikilohydric plants: a comparison of lichens and bryophytes. In *Ecophysiology of Photosynthesis*. 319-341. Springer Berlin Heidelberg.
- Hájek, T., and R. P. Beckett. 2008. Effects of water content components on desiccation and recovery of *Sphagnum* mosses. *Annals of Botany*. 101(1): 165-173.
- Harris, A. 2008. Spectral reflectance and photosynthetic properties of *Sphagnum* mosses exposed to progressive drought. *Ecohydrology*. 1(1): 35-42.
- Hinzman, L. D., N. D. Bettez, W. R. Bolton, F. S. Chapin, M. B. Dyurgerov, C. L. Fastie, B. Griffith, R. D. Hollister, A. Hope, H. P. Huntington, and A. M. Jensen. 2005. Evidence and implications of recent climate change in northern Alaska and other arctic regions. *Climatic Change*. 72(3): 251-298.
- Hobbie, J. E., G. R. Shaver, E. B. Rastetter, J. E. Cherry, S. J. Goetz, K. C. Guay, W. A. Gould, and G. W. Kling. 2017. Ecosystem responses to climate change at a Low Arctic and a High Arctic long-term research site. *Ambio* 46:160–173.
- Hollister, R. D., J. L. May, K. S. Kremers, C. E. Tweedie, S. F. Oberbauer, J. A. Liebig, T. F. Botting, R. T. Barrett, and J. L. Gregory. 2015. Warming experiments elucidate the drivers of observed directional changes in tundra vegetation. *Ecology and evolution*. 5(9): 1881-1895.
- Hope, A. S., J. S. Kimball, and D. A. Stow. 1993. The relationship between tussock tundra spectral reflectance properties and biomass and vegetation composition. *International Journal of Remote Sensing*. 14: 1861-1874.
- Hope, A. S., J. B. Fleming, G. Vourlitis, D. A. Stow, W. C. Oechel. T. and Hack. 1995. Relating CO<sub>2</sub> fluxes to spectral vegetation indices in tundra landscapes: importance of footprint definition. *Polar Record*. 31(177): 245-250.



- Huemmrich, K. F., J. A. Gamon, C. E. Tweedie, S. F. Oberbauer, G. Kinoshita, S. Houston, A. Kuchy, R. D. Hollister, H. Kwon, M. Mano, and Y. Harazono. 2010. Remote sensing of tundra gross ecosystem productivity and light use efficiency under varying temperature and moisture conditions. *Remote Sensing of Environment*. 114(3): 481-489.
- Intergovernmental Panel on Climate Change Fifth Assessment Report: Climate Change. 2013. The physical basis. Working group I contribution to the fifth assessment report of the intergovernmental panel on climate change. T. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, P. Midgley (eds). Cambridge University Press. Cambridge, UK.
- Jia, G.J.; H. E. Epstein, H. E.; Walker, D. A. 2003. Greening of Arctic Alaska, 1981-2001. *Geophysical Research Letters*. 30(20):2067.
- Kattsov, V. M., Walsh, J. E., Chapman, W. L., Govorkova, V. A., Pavlova, T. V., & Zhang, X. (2007). Simulation and projection of Arctic freshwater budget components by the IPCC AR4 global climate models. *Journal of Hydrometeorology*, 8(3), 571-589.
- Kerr, J. T., & Ostrovsky, M. (2003). From space to species: ecological applications for remote sensing. *Trends in Ecology & Evolution*, 18(6), 299-305.
- Komulainen, V.-M., E.-S. Tuittila, H. Vasander and J. Laine. 1999. Restoration of drained peatlands in southern Finland: initial effects on vegetation change and CO<sub>2</sub> balance. *Journal of Applied Ecology*. 36: 634–648.
- Kriegler, F.J., W. A. Malila, R. F. Nalepka, W. Richardson. 1969. Preprocessing transformations and their effects on multispectral recognition. *Proceedings of the Sixth International Symposium on Remote Sensing of Environment*. University of Michigan, Ann Arbor, MI. 97-131.
- Kushida, K., Y. Kim, S. Tsuyuzaki, and M. Fukuda. 2009. Spectral vegetation indices for estimating shrub cover, green phytomass and leaf turnover in a sedge-shrub tundra. *International Journal of Remote Sensing* 30:1651-1658.
- Laidler, G. J., P. M. Treitz, and D. M. Atkinson. 2008. Remote sensing of arctic vegetation: relations between the NDVI, spatial resolution and vegetation cover on Boothia Peninsula, Nunavut. *Arctic*. 61:1-13.
- Levitt, J. 1956. The hardiness of plants. *Soil Science*. 82(4): 346.
- Longton, R. E. 1988. Adaptations and strategies of polar bryophytes. *Botanical journal of the Linnean Society*. 98(3): 253-268.
- Lorant, M.M., S. J. Goetz, P. S. Beck, 2011. Tundra vegetation effects on pan-Arctic albedo. *Environmental Research Letters*. 6(2):024014.
- Mack, M. C., M. S. Bret-Harte, T. N. Hollingsworth, R. R. Jandt, E. A. Schuur, G. R. Shaver, and D. L. Verbyla. 2011. Carbon loss from an unprecedented Arctic tundra wildfire. *Nature*. 475(7357): 489-492.
- McNeil, P. and J. M. Waddington. 2003. Moisture controls on *Sphagnum* growth and CO<sub>2</sub>

exchange on a cutover bog. *Journal of Applied Ecology*. 40: 354-367.

Molau, U. and J. M. Alatalo. 1998. Responses of subarctic-alpine plant communities to simulated environmental change: biodiversity of bryophytes, lichens, and vascular plants. *Ambio*. 322-329.

Oechel, W. C. and B. Sveinbjornsson. 1978. Primary production processes in Arctic bryophytes at Barrow, Alaska. In L. L. Tieszen (Ed.), *Vegetation Production and Production Ecology of an Alaskan Arctic Tundra*: 269-298. New York: Springer-Verlag.

Oechel, W. C., S. J. Hastings, G. Vourlitis, M. Jenkins, G. Riechers, and N. Grulke. 1992. Recent change of Arctic tundra ecosystems from a net carbon dioxide sink to a source. *Nature*. 361(6412): 520-523.

Oechel, W. C., G. L. Vourlitis, J. Verfaillie, T. Crawford, S. Brooks, E. Dumas, A. Hope, D. Stow, B. Boynton, V. Nosov, and R. Zulueta. 2000. A scaling approach for quantifying the net CO<sub>2</sub> flux of the Kuparuk River Basin, Alaska. *Global Change Biology*. 6(S1): 160-173.

Olivas, P. C., S. F. Oberbauer, C. E. Tweedie, W. C. Oechel, D. Lin, and A. Kuchy. 2011. Effects of fine-scale topography on CO<sub>2</sub> flux components of Alaskan coastal plain tundra: Response to contrasting growing seasons. *Arctic, Antarctic, and Alpine Research*. 43(2): 256-266.

Oliver, M. J. and J. D. Bewley. 1984. Plant Desiccation and Protein Synthesis V. Stability of Poly (A)<sup>-</sup> and Poly (A)<sup>+</sup> RNA during desiccation and their synthesis upon rehydration in the desiccation-tolerant moss *Tortula ruralis* and the intolerant moss *Cratoneuron filicinum*. *Plant Physiology*. 74(4): 917-922.

Oliver, M. J., J. Velten, B. D. Mishler. 2005. Desiccation tolerance in bryophytes: A reflection of the primitive strategy for plant survival in dehydrating habitats? *Integrative and Comparative Biology*. 45(5): 788-799.

Overland J. E., M. Wang, and N. A. Bond. 2002. Recent Temperature Changes in the Western Arctic during Spring. *Journal of Climate*. 15(13): 1702-1716.

Potter, J. A., M. C. Press, T. V. Callaghan, and J. A. Lee. 1995. Growth responses of *Polytrichum commune* and *Hylocomium splendens* to simulated environmental change in the subarctic. *New Phytologist*. 131(4): 533-541.

Proctor, M.C. and Z. Tuba. 2002. Poikilohydry and homoihydry: antithesis or spectrum of possibilities?. *New Phytologist*. 156(3): 327-349.

Proctor, M. C., M. J. Oliver, A. J. Wood, P. Alpert, L. R. Stark, N. L. Cleavitt, and B. D. Mishler. 2007. Desiccation-tolerance in bryophytes: a review. *The Bryologist*. 110(4): 595-621.

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.

Raynolds, M. K., J. C. Comiso, D. A. Walker and D. Verbyla. 2008. Relationship between satellite-derived land surface temperatures, arctic vegetation types, and NDVI. *Remote Sensing of Environment*. 112(4): 1884-1894.

534 Riedel, S. M., H. E. Epstein, and D. A. Walker. 2005. Biotic controls over spectral reflectance of  
535 arctic tundra vegetation. *International Journal of Remote Sensing*. 26(11): 2391-2405.

536 Riordan, B., D. Verbyla, and A. D. McGuire. 2006. Shrinking ponds in subarctic Alaska based on  
537 1950–2002 remotely sensed images. *Journal of Geophysical Research:*  
538 *Biogeosciences*, 111(G4).

539 Roulet, N., T. I. M. Moore, J. Bubier, and P. Lafleur. 1992. Northern fens: methane flux and  
540 climatic change. *Tellus B*. 44(2): 100-105.

541 Rydin, H. and A. J. S. McDonald. 1985. Photosynthesis in *Sphagnum* at different water contents.  
542 *Journal of Bryology*. 13: 579-584.

543 Schipperges, B. and H. Rydin. 1998. Response of photosynthesis of *Sphagnum*  
544 species from contrasting microhabitats to tissue water content and repeated  
545 desiccation. *New Phytologist*. 149: 677-684.

546 Serrano, L., S. L. Ustin, D. A. Roberts, J. A. Gamon, J. Penuelas, 2000. Deriving water content of  
547 chaparral vegetation from AVIRIS data. *Remote Sensing of Environment*, 74(3):570-581.

548 Shaver, G. R., and F. S. Chapin III. 1991. Production: biomass relationships and element cycling  
549 in contrasting Arctic vegetation types. *Ecological Monographs* 61: 1–31.

550 Shaver, G. R., L. E. Street, E. B. Rastetter, M. T. Van Wijk, and M. Williams. 2007. Functional  
551 convergence in regulation of net CO<sub>2</sub> flux in heterogeneous tundra landscapes in Alaska  
552 and Sweden. *Journal of Ecology*. 95(4): 802-817.

553 Skre, O. and W. C. Oechel. 1981. Moss functioning in different taiga ecosystems in interior  
554 Alaska. *Oecologia*. 48(1): 50-59.

555 Stow, D., A. Petersen, A. Hope, R. Engstrom, and L. Coulter. 2007. Greenness trends of  
556 Arctictundra vegetation in the 1990s: comparison of two NDVI data sets from NOAA  
557 AVHRRsystems. *International Journal of Remote Sensing*. 28(21): 4807-4822.

558 Stoy, P.C., M. Williams, J. G. Evans, A. Prieto-Blanco, M. Disney, T. C. Hill, H. C. Ward, T. J. Wade,  
559 L. E. Street. 2013. Upscaling tundra CO<sub>2</sub> exchange from chamber to eddy covariance  
560 tower. *Arctic, Antarctic, and Alpine Research*. 45(2):275-284.

561 Titus, J. E., D. J. Wagner, and M. D. Stephens. 1983. Contrasting water relations of  
562 photosynthesis for two *Sphagnum* mosses. *Ecology*. 64(5): 1109-1115.

563 Toolik Field Station, Environmental Data Center. 2016.  
564 [https://toolik.alaska.edu/edc/about/conditions\\_of\\_use.php?page=/edc/abiotic\\_monitoring/data\\_query.php](https://toolik.alaska.edu/edc/about/conditions_of_use.php?page=/edc/abiotic_monitoring/data_query.php)  
565

566 Turetsky M. R., M. C. Mack, T. N. Hollingsworth, and J. W. Harden. 2010. The role of mosses in  
567 ecosystem succession and function in Alaska's boreal forest. *Canadian Journal of Forest*  
568 *Research*. 40: 1237–1264.

569 Turetsky, M. R., B. Bond-Lamberty, E. Euskirchen, J. Talbot, S. Frolking, A. D. McGuire, and E. S.  
570 Tuittila. 2012. The resilience and functional role of moss in boreal and arctic ecosystems.  
571 *New Phytologist*, 196(1): 49-67.

- Ueno, T. and H. Kanda. 2006. Photosynthetic response of the arctic semi-aquatic moss *Calliergon giganteum* to water content. *Aquatic Botany*. 85: 241-243.
- Van Breemen, N. 1995. How Sphagnum bogs down other plants. *Trends in Ecology & Evolution*. 10(7): 270-275.
- Van Gaalen, K. E., L. B. Flanagan, D. R. Peddle. 2007. Photosynthesis, chlorophyll, and spectral reflectance in Sphagnum moss at varying water contents. *Oecologia*. 153(1): 19-28.
- Viereck, L. A. 1975. Forest ecology of the Alaska taiga. In *Proceedings, Circumpolar Conference on Northern Ecology*, Sept 15-18, 1975. 11.
- Vourlitis, G. L., W. C. Oechel, A. Hope, D. Stow, B. Boynton, J. Verfaillie, R. Zulueta, and S. J. Hastings. 2000. Physiological models for scaling plot measurements of CO<sub>2</sub> flux across an arctic tundra landscape. *Ecological Applications*. 10(1): 60-72.
- Walker, D. A., H. E. Epstein, G. J. Jia, A. Balser, C. Copass, E. J. Edwards, W. A. Gould, J. Hollingsworth, J. Knudson, H. A. Maier, and A. Moody, A., 2003. Phytomass, LAI, and NDVI in northern Alaska: Relationships to summer warmth, soil pH, plant functional types, and extrapolation to the circumpolar Arctic. *Journal of Geophysical Research: Atmospheres*. 108(D2).
- Walker, D. A., H. E. Epstein, M. K. Reynolds, P. Kuss, M. A. Kopecky, G. V. Frost, F. J. A. Daniëls, M. O. Leibman, N. G. Moskalenko, G. V. Matyshak, and O. V. Khitun. 2012. Environment, vegetation and greenness (NDVI) along the North America and Eurasia Arctic transects. *Environmental Research Letters*. 7(1): 015504.
- Zona, D., W. C. Oechel, J. H. Richards, S. Hastings, I. Kopetz, H. Ikawa, and S. Oberbauer. 2011. Light-stress avoidance mechanisms in a Sphagnum-dominated wet coastal Arctic tundra ecosystem in Alaska. *Ecology*, 92(3): 633-644.

596 List of Figures:

597 **Figure 1:** NDVI (solid line), near-infrared (dotted line), and red (dashed line) of four  
598 communities by percent moisture content (left panels) and hours after initial saturation (right  
599 panels) during drying and after re-saturation during experiment 1.

600 **Figure 2:** Gross primary productivity and NDVI of four communities by percent moisture  
601 content during drying and three measurement times after re-saturation during experiment 2.

602 **Figure 3:** Gross primary productivity and NDVI of four communities by hours of drying and three  
603 measurement times after re-saturation during experiment 2.

604 **Figure 4:** Drying response indexes of four communities for GPP and NDVI during experiment 2  
605 compared using a one-way analysis of variance. Statistical significance denoted with \*.

606 List of Tables:

607 **Table 1:** Gross primary productivity and NDVI of four communities by percent moisture content  
608 during drying and three measurement times after re-saturation compared using repeated  
609 measures analysis of variance with Tukey's post-hoc analysis, along with the initial saturated  
610 and final oven dry weight for each of the four community types during experiment 2. Letters  
611 denote statistically significant differences ( $p < 0.05$ ) between moisture content measurements.

612