

Monitoring neonicotinoid exposure for bees in rural and peri-urban areas of the UK during the transition from pre- to post-moratorium.

Elizabeth Nicholls, CRISTINA BOTÍAS, Ellen L Rotheray, Penelope Whitehorn,
Arthur David, Robert Fowler, Thomas David, Hannah Feltham, Jennifer L.
Swain, Patricia Wells, Elizabeth M Hill, Juliet L Osborne, and Dave Goulson

Environ. Sci. Technol., **Just Accepted Manuscript** • DOI: 10.1021/acs.est.7b06573 • Publication Date (Web): 28 Jun 2018

Downloaded from <http://pubs.acs.org> on July 30, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

This document is the Accepted Manuscript version of a Published Work that appeared in final form in *Environmental Science and Technology*, copyright © American Chemical Society after peer review and technical editing by the publisher.

To access the final edited and published work see <https://doi.org/10.1021/acs.est.7b06573>



ACS Publications

is published by the American Chemical Society, 1155 Sixteenth Street N.W.,
Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society.
However, no copyright claim is made to original U.S. Government works, or works
produced by employees of any Commonwealth realm Crown government in the course
of their duties.

Monitoring neonicotinoid exposure for bees in rural and peri-urban areas of the UK during the transition from pre- to post-moratorium.

Elizabeth Nicholls^{a*}, Cristina Botías^{a‡}, Ellen L. Rotheray^a, Penelope Whitehorn^{b†}, Arthur David^{a‡}, Robert Fowler^a, Thomas David^{a,c}, Hannah Feltham^{b§}, Jennifer L. Swain^c, Patricia Wells^c, Elizabeth M. Hill^a, Juliet L. Osborne^d & Dave Goulson^a

^aSchool of Life Sciences, University of Sussex, Falmer, BN1 9QG, UK


^bSchool of Natural Sciences, University of Stirling, Stirling, FK9 4LA, UK

^cRothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK

^dEnvironment and Sustainability Institute, University of Exeter, Penryn, TR10 9FE, UK

[*e.nicholls@sussex.ac.uk](mailto:e.nicholls@sussex.ac.uk)

ABSTRACT: Concerns regarding the impact of neonicotinoid exposure on bee populations recently led to an EU-wide moratorium on the use of certain neonicotinoids on flowering crops. Currently evidence regarding the impact, if any, the moratorium has had on bees' exposure is limited. We sampled pollen and nectar from bumblebee colonies in rural and peri-urban



habitats in three UK regions; Stirlingshire, Hertfordshire and Sussex. Colonies were sampled over three years; prior to the ban (2013), during the initial implementation when some seed-treated winter-sown oilseed rape was still grown (2014), and following the ban (2015). To compare species-level differences, in 2014 only, honeybee colonies in rural habitats were also sampled. Over half of all samples were found to be contaminated (n=408), with thiamethoxam being the compound detected at the highest concentrations in honeybee- (up to 2.29 ng/g in nectar in 2014, median ≤ 0.1 ng/g, n=79) and bumblebee-collected pollen and nectar (up to 38.77 ng/g in pollen in 2013, median ≤ 0.12 ng/g, n=76). Honeybees were exposed to higher concentrations of neonicotinoids than bumblebees in 2014. While neonicotinoid exposure for rural bumblebees declined post-ban (2015), suggesting a positive impact of the moratorium, the risk of neonicotinoid exposure for bumblebees in peri-urban habitats remained largely the same between 2013 and 2015.

31 INTRODUCTION

32 Neonicotinoids are the most commonly used insecticides worldwide¹. Their systemic nature
33 means that, following seed-application to crops such as oilseeds or cereals, neonicotinoids become
34 incorporated into the tissues of a plant as it grows, including pollen and nectar, the main source of
35 food for economically important pollinators, such as honeybees and bumblebees². Multiple studies
36 have raised concerns regarding the negative impacts of neonicotinoid exposure on bees³. Whitehorn *et al.* (2012)⁴ found that exposure of bumblebees to pollen and nectar containing 6 ng/g and 0.7 ng/g of
37 imidacloprid respectively, resulted in slower colony growth and the production of fewer new queens,
38 relative to unexposed colonies. Other studies have observed detrimental impacts on foraging and
39 navigation^{5,6}, immunity⁷ and worker mortality⁸. Based on these findings, in 2013 the European
40 Commission instated a EU-wide moratorium on the use of three types of neonicotinoid,
41 thiamethoxam, clothianidin and imidacloprid on bee-attractive flowering crops such as oilseed rape⁹.
42 In 2018, this ban was subsequently expanded to include all field crops^{10–12}.

44 Criticism has been levied against studies cited in support of the moratorium, mainly for using
45 neonicotinoid concentrations purported to exceed those routinely experienced by foraging bees¹³,
46 sparking demand for further evidence as to what constitutes a ‘field-realistic’ dose. Several studies
47 have screened bee-collected pollen and nectar^{14–19} for neonicotinoid residues, quantifying the
48 ‘exposure landscape’ by incorporating multiple chemicals from several forage sources.
49 Concentrations have been shown to vary considerably across studies, depending on location, time of
50 year and species. Pollen sampled from rural bumblebee colonies in Sussex, England, prior to the
51 implementation of the moratorium in 2013, was found to contain 18 ng/g of thiamethoxam on
52 average, with pollen collected from nests in nearby peri-urban areas containing up to 20 ng/g
53 imidacloprid¹⁵ (mean=6.5 ng/g), well above the 6 ng/g used by Whitehorn *et al.*⁹. A large scale
54 Swedish field study found clothianidin concentrations averaging 5.4 ng/g in nectar sampled from
55 bumblebees foraging in fields of seed-treated oilseed rape (range 1.4–14 ng/g)¹⁶. In contrast, a study
56 conducted in Germany found considerably lower average concentrations (0.88 ng/g) in pollen
57 collected from bumblebee nests adjacent to neonicotinoids treated winter-sown oilseed rape²⁰, and a

58 more recent study conducted across the UK, Hungary and Germany reported that concentrations
59 detected in pollen and nectar collected by honeybees, bumblebees and the solitary bee *Osmia bicornis*
60 rarely exceeded 1.5 ng/g²¹. The wide ranging values reported by these studies highlights the need for
61 further data to determine the actual exposure risk, particularly for wild bees.

62 Here we monitored bees' risk of neonicotinoid exposure during the period from pre- to post-
63 moratorium, by screening pollen and nectar collected from bumblebee colonies located in several
64 regions; Sussex (2013-2015) and Hertfordshire (2014 only) in the south of England and Stirling,
65 Scotland (2013 only) in the north of the UK. Given the total weight of neonicotinoids applied in
66 Scotland is much lower compared to the south of England (FERA PUS STATS database²²), we
67 expected the exposure risk to be lowest for bees in this region. There is currently limited data on the
68 exposure risk for wild bees from foraging on ornamental plants grown using neonicotinoids^{15,23,24} and
69 the use of neonicotinoid-based garden sprays, therefore we monitored bumblebees in both rural and
70 peri-urban habitats (Sussex and Stirling only), the latter consisting of domestic gardens located on the
71 outskirts of urban areas. For bees in rural areas, we expected neonicotinoid concentrations in pollen
72 and nectar collected in 2015 to be lower than those collected in 2013, before the implementation of
73 the moratorium. In 2014, the impact of the ban may not have fully come into effect, as any winter-
74 sown oilseed crops would have been drilled prior to the implementation of the ban in December 2013
75 and therefore may still have been seed-treated with neonicotinoids. To compare species-level
76 differences in exposure risk during this transitional year (2014), we also screened pollen and nectar
77 from rural honeybee colonies located in Sussex and Hertfordshire.

78

79 MATERIALS AND METHODS

80 **Site Information** Bumblebee colonies (*B.terrestris audax*) were obtained from Agralan Ltd.,
81 Swindon, UK (originating from Biobest, Belgium) and in late spring (late May to early June, see
82 Table 1 for exact dates) were placed into the field:

i) to monitor exposure risk over the course of the implementation of the ban for both rural and peri-urban habitats, bumblebee colonies were placed in rural (n=135, n=32-47/year) and peri-urban (n=42, 12-15/year) locations across Sussex each year between 2013 and 2015. While the UK granted a derogation to use neonicotinoids on oilseed rape in 2015, this was limited to a portion of East England and did not affect the study area;

ii) to assess regional differences in neonicotinoid exposure between the north and south of the UK, prior to the implementation of the ban (2013), bumblebee colonies were also placed in rural (n=10) and peri-urban (n=20) locations in Stirling. In 2014 only, bumblebees were also placed in rural locations across Hertfordshire (n=30) for comparison with Sussex colonies;

iii) to compare species-level differences in exposure risk, 15 rural bumblebee colonies were each paired with a honeybee colony (located within 10m distance and placed into the field at the beginning of April) in both Sussex and Hertfordshire in 2014 only. Queenright honeybee colonies were obtained from experimental stocks at the University of Sussex and Rothamsted Research, which at the beginning of the experiment consisted of a single brood box and a super containing frames of fresh foundation wax, with additional space for bees to store pollen and nectar added as necessary. We also mapped which crops were grown in ten, 5 km² surrounding the experimental colonies in Sussex and Hertfordshire in 2014 (Fig. S4) and, where possible, asked farmers growing winter-sown oilseed rape which seed treatments they had used (Table S4).

Sampling Pollen and nectar was collected from bumblebee colonies following four, eight and ten weeks of foraging in the field. Pollen was scraped out of the colony using a stainless steel micro-spoon, which was cleaned using methanol to avoid cross-contamination. From each colony, we aimed to collect enough pollen to fill a 1.5 ml micro-centrifuge tube, to ensure enough material for chemical analysis. Concurrently, 1.5 ml of nectar was obtained from nectar pots using disposable glass pipettes. However, care was taken not to completely deplete bumblebee colony stores. Where stores were low, no sample was collected (Table 2).

For honeybees, samples were collected once per month in April, May and June 2014, with the last two sampling dates coinciding with sample collection from adjacent bumblebee colonies. Samples were obtained from freshly drawn comb, where possible, to minimise contamination from previous years. Enough pollen to fill a 1.5 ml micro-centrifuge tube was scraped out of ~10 cells using a stainless steel micro-spoon as described above, and 1.5 ml of recently stored nectar was obtained from uncapped and newly drawn comb using disposable glass pipettes. Freshness was determined by first shaking the frame to ensure nectar dripped easily out of the comb. All pollen and nectar samples were stored in individually labelled tubes and put on ice during transport back to the lab, and were then frozen at -20°C until residue analysis was performed.

Chemical analyses: Pollen and nectar samples were extracted using the QuEChERS method¹⁴ and screened for five neonicotinoids: thiamethoxam (TMX), clothianidin (CLO), imidacloprid (IMC), acetamiprid (ACT) and thiacloprid (THC), using ultra high-performance liquid chromatography tandem mass spectrometry (UHPLC-MS/MS). Pollen samples collected in Sussex in 2013 were not screened for acetamiprid.

Sample preparation: Pollen samples were extracted as described by Botias *et al.* (2015)¹⁴. Briefly, 100 mg of pollen was weighed into an Eppendorf tube and 400 pg of deuterated pesticides in ACN were added. The extraction was performed by the addition of 400 µl of water, 500 µl of ACN, 125 mg of magnesium sulphate: sodium acetate mix (4:1) and 125 mg of PSA/C18/ENVI-Carb for the dispersive solid phase extraction (dSPE) step (QuEChERS method). After the first extraction, the aqueous phase and re-suspended pellet were extracted again with 400 µl of ACN and the supernatants combined. Extracts were mixed with PSA/C18/ENVI-Carb and centrifuged. The supernatant was evaporated to dryness under vacuum, reconstituted with 120 µl ACN:H₂O (10:90) and spin filtered (0.22 µm).

Nectar samples were centrifuged at 13,000 relative centrifugal force (RCF) for 10 min to remove plant debris and the supernatant transferred into a clean eppendorf tube. Nectar samples were very viscous and were therefore weighed for more accuracy (175 ± 50 mg depending on availability) and the volume then increased to 400 µl with water. Four hundred pg of deuterated pesticide standard

mixture was added to the nectar and the samples were extracted using the same QuEChERS method described for pollen.

UHPLC-MS/MS analyses. The ultra high-performance liquid chromatography tandem mass spectrometry (UHPLC-MS/MS) method described by Botias *et al.* (2015)¹⁴ was used for the analysis of samples. UHPLC-MS/MS analyses were carried out using a Waters Acquity UHPLC system coupled to a Quattro Premier triple quadrupole mass spectrometer from Micromass (Waters, Manchester, UK). Data were acquired using MassLynx 4.1 and the quantification was carried out by calculating the response factor of neonicotinoid compounds to their respective internal standards. Concentrations were determined using a least-square linear regression analysis of the peak area ratio *versus* the concentration ratio (native to deuterated). Method detection and quantification limits (MDL and MQL, respectively) as well as recoveries were determined as described by Botias *et al.* (2015)¹⁴ (Table S1-3).

Quality control. One blank workup sample (*i.e.* solvent without matrix) per batch of eleven samples was included and injected on the UHPLC-MS/MS to ensure that no contamination occurred during the sample preparation. Solvent samples were also injected between sample batches to ensure that there was no carryover in the UHPLC system that might affect adjacent results in analytical runs. Samples were analysed in a random order and quality control samples (*i.e.* standards) were injected during runs every ten samples to check the sensitivity of the machine. Identities of detected neonicotinoids were confirmed by comparing ratio of MRM transitions in samples and pure standards.

Statistical Analysis. All analyses were performed using R-3.3.3. Residue concentrations that were above the MDL but below the MQL were assigned the MDL (Tables 2-3, range 0.03-0.10 ng/g). Concentrations below the MDL were assumed to be zero¹⁴. Shapiro-Wilk tests, combined with inspection of *q-q* plots, confirmed that residue data were not normally distributed. Therefore we compared the frequency of neonicotinoid contamination using contingency tables and either χ^2 or Fisher's exact tests (where expected frequencies were <5). To compare total neonicotinoid concentrations between regions (Sussex *vs.* Stirling; Sussex *vs.* Herts), habitats (Rural *vs.* Peri-Urban) and years of the study (2013 *vs.* 2015) we used non-parametric Mann-Whitney tests. For honeybee

data, where frequencies of contamination and residue concentrations were compared between samples from the same hive over several months, we used Cochran's Q test (with McNemar's test for post-hoc comparisons) and the Wilcoxon Signed-Rank test, with Bonferroni corrections to account for multiple comparisons. Given the relatively small number of pollen and nectar samples collected from each bumblebee colony, for analyses involving bumblebees we pooled samples collected after four and eight weeks in the field.

RESULTS

Bumblebees: In total, 233 pollen and nectar samples were collected from bumblebee colonies placed in rural and peri-urban habitats in the regions of Stirling, Sussex and Hertfordshire between 2013 and 2015. Forty percent of all samples screened were found to be contaminated with neonicotinoids, predominantly thiamethoxam (23%), thiacloprid (15%) and imidacloprid (10%). Pollen samples were more often contaminated (62% samples) than nectar (25% samples) and the mean combined total residues detected in pollen (Pollen N=132, 62% samples, mean \pm standard deviation (SD) =1.44 \pm 5.44 ng/g, median <MDL, max= 38.77 ng/g) were more than ten times higher (Nectar N=101, mean \pm SD= 0.12 \pm 0.44 ng/g, median <MDL, max=3.58 ng/g).

Differences in exposure by habitat and year: In 2013, the frequency of neonicotinoid contamination was similar for pollen (Table 1, $\chi^2_1=0$, $p=1.000$, Rural =58%; Peri-urban= 59%) and nectar ($\chi^2_1=0$, $p=1.000$, Rural=14%, Peri-urban =14%) sampled from peri-urban (PU) and rural (R) bumblebee colonies across the regions of Sussex (SU) and Stirling (ST) (Table 1). Concentrations of neonicotinoids were very similar in nectar (Mann-Whitney $U_{21, 21}=225$, $p=0.867$, mean $_{PU}\leq 0.10$, median $_{PU}\leq 0.10$, mean $_R\pm SD=0.22\pm 0.55$ ng/g, median $_R$ <MDL), and though higher in pollen from rural colonies, this difference was not significant ($U_{36, 32}=603.5$, $p=0.73$; mean $_R=3.37\pm 9.36$ ng/g, median $_R\leq 0.12$, mean $_{PU}= 1.28\pm 3.62$ ng/g, median $_{PU}\leq 0.12$). While nectar from both habitats contained only one type of neonicotinoid, predominantly thiamethoxam, over a quarter of pollen samples from bumblebee colonies in rural (28%) and peri-urban (26%) habitats contained more than one residue. Thiamethoxam (up to 38.77 ng/g, median <0.12, mean $\pm SD= 2.08\pm 7.47$ ng/g) and clothianidin (up to 2.08 ng/g, mean ≤ 0.12 ng/g, median <0.12 ng/g) were present at the highest concentrations in rural

colonies. While thiamethoxam was also present in a high percentage of pollen samples collected from peri-urban colonies in Sussex (79% samples), thiacloprid was found at the highest concentration in these samples (up to 14.8 ng/g, mean ≤ 0.04 ng/g, median < 0.04 ng/g).

In 2014, less than 10% of pollen (n=13) and nectar (n=13) samples from rural bumblebee colonies in Sussex contained neonicotinoids, all thiamethoxam and below the method quantification limit, whereas a significantly higher proportion of both pollen (85%, $\chi^2_1=8.987$, $p=0.003$, n=7) and nectar samples (80%, Nectar $\chi^2_1=6.152$, $p=0.013$, n=5) from peri-urban nests were contaminated (N=12), frequently with multiple residues (40% nectar samples, 29% of pollen). Again, thiacloprid (up to 9.32 ng/g in pollen, mean $= 1.34 \pm 3.52$ ng/g, median ≤ 0.04 ng/g) and thiamethoxam (up to 3.48 ng/g in pollen, mean $= 0.76 \pm 1.52$, median $= 0.10$ ng/g) and were detected at the highest concentrations.

In 2015, the frequency of neonicotinoid detection was similar for nectar collected from rural and peri-urban bumblebee colonies in Sussex ($\chi^2_1=0.158$, $p=0.691$, Rural=47%, Peri-urban=33%) as were the concentrations present (Mann-Whitney $U_{19, 12}=130.5$, $p=0.469$, mean_R $=0.10 \pm 0.15$ ng/g, median_R $< \text{MDL}$, mean_{PU} $=0.08 \pm 0.17$ ng/g, median_{PU} $< \text{MDL}$). While the frequency of detection (Rural=35%, Peri-urban=64%), proportion of samples with multiple residues (Rural=9% vs. Peri-urban=18%) and mean concentration of neonicotinoids were higher in pollen from peri-urban nests, the difference was not significant ($\chi^2_1=1.238$, $p=0.266$, $U_{22, 11}=75.5$, $p=0.06$, mean_R $=0.06 \pm 0.14$ ng/g, median_R $< \text{MDL}$, mean_{PU} $=1.29 \pm 3.30$ ng/g, median_{PU} $< \text{MDL}$). Both habitats were contaminated predominantly with thiacloprid (up to 0.44 ng/g, mean $\pm \text{SD} = 0.04 \pm 0.11$ ng/g, median $< \text{MDL}$), and imidacloprid (up to 11.16 ng/g in peri-urban nests, mean $\pm \text{SD} = 0.21 \pm 1.40$ ng/g, median < 0.14), though a small proportion of peri-urban samples also contained acetamiprid (4% up to 1.4 ng/g, mean ≤ 0.03 ng/g, median $< \text{MDL}$).

To compare the changing risk of exposure to peri-urban and rural bees over the transitional period from pre- to post- moratorium, we compared residue concentrations in 2013 and 2015 for Sussex bumblebee colonies only. For pollen collected from rural colonies there was a significant decrease in overall combined residue concentrations between years (Mann-Whitney $U_{23, 22}=385$, $p=0.002$, mean₂₀₁₃ $= 5.10 \pm 11.40$ ng/g, median ≤ 0.12 ng/g, mean₂₀₁₅ $= 0.06 \pm 0.14$ ng/g, median $< \text{MDL}$), but not for

nectar ($U_{14, 19}=98$, $p=0.134$; $\text{mean}_{2013}=0.20\pm0.51$ ng/g, median <MDL, $\text{mean}_{2015}=0.10\pm0.15$ ng/g, median <MDL). When considering just those neonicotinoids affected by the moratorium (thiamethoxam, clothianidin and imidacloprid), the same effect is observed, with a significant decrease in residue concentrations in pollen ($U_{23, 22}=389$, $p<0.001$, $\text{mean}_{2013}=5.02\pm11.32$ ng/g, median ≤ 0.12 ng/g, $\text{mean}_{2015}=0.05\pm0.14$ ng/g, median <MDL) but not nectar between 2013 and 2015 ($U_{14, 19}=140$, $p=0.676$; $\text{mean}_{2013}=0.20\pm0.51$ ng/g, median <MDL, $\text{mean}_{2015}<\text{MDL}$, median <MDL). In contrast, concentrations of thiacloprid, which was unaffected by the ban, increased significantly in nectar between 2013 and 2015 ($U_{14, 19}=84$, $p=0.013$, $\text{mean}_{2013}<\text{MDL}$, median <MDL, $\text{mean}_{2015}=0.09\pm0.15$ ng/g, median <MDL). Concentrations of thiacloprid in pollen remained unchanged over this period ($U_{23, 22}=267$, $p=0.627$, $\text{mean}_{2013}=0.08\pm0.31$ ng/g, median <MDL, $\text{mean}_{2015}<\text{MDL}$, median <MDL).

For peri-urban nests, there was no significant difference in overall residue concentrations in either pollen ($U_{19, 11}=124$, $p=0.408$, $\text{mean}_{2013}=2.11\pm4.56$ ng/g, median=0.12 ng/g, $\text{mean}_{2015}=1.29\pm0.14$ ng/g, median ≤ 0.04 ng/g) or nectar ($U_{13, 12}=62.5$, $p=0.276$, $\text{mean}_{2013}=0.02\pm0.05$ ng/g, median <MDL, $\text{mean}_{2015}=0.08\pm0.17$ ng/g, median <MDL), samples collected between 2013 and 2015. When considering either the banned neonicotinoids only (Pollen, $U_{19, 11}=134.5$, $p=0.188$; $\text{mean}_{2013}=0.63\pm1.64$ ng/g, median ≤ 0.12 , $\text{mean}_{2015}=1.14\pm3.33$ ng/g, median <MDL; Nectar $U_{13, 12}=76$, $p=0.898$, $\text{mean}_{2013}<\text{MDL}$, median <MDL, $\text{mean}_{2015}<\text{MDL}$, median <MDL) or thiacloprid, which was unaffected by the ban (Pollen $U_{19, 11}=104$, $p=1$, $\text{mean}_{2013}=1.47\pm4.41$ ng/g, median <MDL, $\text{mean}_{2015}<\text{MDL}$, median <MDL, Nectar $U_{13, 12}=58.5$, $p=0.067$, $\text{mean}_{2013}<\text{MDL}$, median <MDL, $\text{mean}_{2015}=0.05\pm0.13$ ng/g, median <MDL), again there was no difference in the concentrations detected in pollen and nectar collected from peri-urban nests between 2013 and 2015.

Regional differences in exposure In 2013, pollen collected from bumblebee colonies in Sussex (SU) was more frequently contaminated ($\chi^2_1=15.62$, $p<0.001$, Sussex=79%; Stirling=27%), with significantly higher concentrations of neonicotinoids than pollen collected from colonies in Stirling (ST) (Mann-Whitney $U_{42,26}=276$, $p<0.001$; $\text{mean}_{\text{SU}}\pm\text{SD}=3.74\pm9.01$ ng/g, $\text{median}_{\text{SU}}\leq 0.12$ ng/g, $\text{mean}_{\text{ST}}\pm\text{SD}=0.20\pm0.49$ ng/g, $\text{median}_{\text{ST}}<\text{MDL}$). Nectar was contaminated at similar frequencies

(Fisher's Exact Test $p=1.00$, Sussex=14%; Stirling 12.5%) and concentrations ($U_{27,15}=200$, $p=0.931$; $\text{mean}_{\text{SU}}=0.11\pm0.37$ ng/g, $\text{median}_{\text{SU}} < \text{MDL}$, $\text{mean}_{\text{ST}}=0.13\pm0.47$ ng/g, $\text{median}_{\text{ST}} < \text{MDL}$).

Pollen sampled from Sussex colonies was more frequently contaminated with multiple residues (Peri-urban=37%, Rural=35%) compared to Stirling samples (Peri-urban=8%, Rural=15%), and the concentrations of thiamethoxam detected in pollen were considerably higher ($\text{mean}_{\text{SU}}=0.58\pm1.64$ ng/g, $\text{median}=0.12$ ng/g vs. $\text{mean}_{\text{ST}}\leq0.12$ ng/g, $\text{median} < 0.12$ ng/g). Sussex peri-urban colonies in particular also contained higher concentrations of thiacloprid compared to Stirling ($\text{mean}_{\text{SU}}=1.47\pm4.41$ ng/g $\text{median} < 0.03$ ng/g vs. $\text{mean}_{\text{ST}}=0.07\pm0.22$ ng/g, $\text{median} < 0.03$ ng/g). Imidacloprid was also frequently detected in pollen from Sussex nests in 2013, but was not detected in any samples from Stirling. Clothianidin was not detected in any Sussex nests, but accounted for the highest residue concentrations detected in nests in Stirling ($\text{mean}_{\text{ST}}=0.16\pm0.58$ g/g, $\text{median} < \text{MDL}$, $\text{max}_{\text{ST}}=2.08$ ng/g).

In 2014, residues detected in pollen and nectar samples collected from bumblebee colonies placed in rural habitats in Hertfordshire (H) and Sussex (SU) were all below the limits of quantification (<0.04 - 0.1 ng/g). Though there was a higher frequency of contamination of both pollen (H=36%, SU=7%) and nectar (H=20%, SU=8%) from Hertfordshire colonies, this difference was not significant (Nectar: Fisher's Exact Test $p=0.560$; $N_{\text{SU}}=13$, $N_{\text{H}}=10$; Pollen $p=0.142$, $N_{\text{SU}}=13$, $N_{\text{H}}=11$). A small proportion of pollen from Sussex (10%), and nectar from both regions was contaminated with thiamethoxam (SU=10%; H=20%). Pollen from Hertfordshire colonies also contained acetamiprid (10%) and, more frequently, thiacloprid (40%).

Honeybees: In total, 175 pollen and nectar samples were collected from honeybee hives in Sussex and Hertfordshire between April and June May 2014, with over two thirds (68%) found to be contaminated with neonicotinoids. Total residue concentrations in nectar ($N=85$, $\text{mean}\pm\text{SD}=0.64\pm0.84$ ng/g, $\text{median}=0.20$ ng/g, $\text{max}=4.23$ ng/g) were approximately three times the concentrations detected in pollen ($N=90$, $\text{mean}\pm\text{SD}=0.20\pm0.32$ ng/g, $\text{median}\leq0.12$ ng/g, $\text{max}=1.74$ ng/g), with 40% of nectar samples containing more than one residue, compared to just 9% of pollen samples. Alongside thiamethoxam, which was highly prevalent in both pollen (61% of samples) and nectar

(69%), clothianidin was also frequently detected in nectar collected from honeybee hives (40%), but only once in pollen (Table 2). Imidacloprid and thiacloprid were detected in a very small percentage of samples (4-5%) and acetamiprid was not detected.

Seasonal differences: Frequency of neonicotinoid detection in pollen (Cochran's $Q=24.67$, $df=2$, $p<0.001$) and nectar ($Q=20.38$, $df=2$, $p<0.001$) sampled from honeybee colonies in 2014 changed significantly across the season. The highest frequency and concentration of neonicotinoid residues were detected in April (Fig. 3), when nearly all nectar samples collected from hives in Hertfordshire (H) and Sussex (SU) were contaminated with neonicotinoids ($H=100\%$, $\text{mean}_H \pm \text{SD} = 1.46 \pm 0.66$ ng/g; median = 1.17 ng/g; $SU=93\%$, $\text{mean}_{SU} = 0.95 \pm 1.13$ ng/g, median ≤ 0.12 ng/g). Likewise, almost all pollen samples contained neonicotinoid residues ($H=80\%$, $\text{mean}_H = 0.41 \pm 0.47$ ng/g, median ≤ 0.12 ng/g; $SU=100\%$, $\text{mean}_{SU} = 0.23 \pm 0.19$ ng/g, median ≤ 0.12 ng/g) in April.

Between April and May, there was a similar frequency of neonicotinoid detection in both pollen (April = 90%, May = 73%, McNemar test, $p=0.287$) and nectar (April = 81%, May = 80%, $p=0.760$). While the concentration of neonicotinoid residues in pollen remained the same as the previous month (Wilcoxon signed-rank test, $Z_{30}=0.28$, $p=0.120$, $\text{mean}_{\text{April}} = 0.32 \pm 0.37$ ng/g, median ≤ 0.12 ng/g; $\text{mean}_{\text{May}} = 0.22 \pm 0.33$, median ≤ 0.12 ng/g), neonicotinoid concentrations in nectar, previously high in comparison to pollen, declined significantly between April and May ($Z_{26}=0.75$, $p<0.001$; $\text{mean}_{\text{April}} = 1.20 \pm 0.95$ ng/g, median = 1.06 ng/g, $\text{mean}_{\text{May}} = 0.65 \pm 0.72$, median = 0.27 ng/g).

At the final sampling point in June, neonicotinoid concentrations detected in samples from both regions were below the limit of quantification, and were significantly lower than in May (Pollen $Z_{30}=0.55$, $p=0.003$; Nectar $Z_{27}=0.73$, $p<0.001$). The frequency of neonicotinoid detection in both pollen (30% samples, McNemar test, $p=0.002$) and nectar (34% samples, $p=0.002$) was also significantly lower than the previous month (Table 2)

Regional differences: While overall neonicotinoid concentrations in pollen contamination did not differ between Hertfordshire and Sussex (Mann-Whitney $U_{45, 45}=1014$, $p=0.100$, $\text{mean}_H = 0.23 \pm 0.36$, median ≤ 0.12 ng/g, $\text{mean}_{SU} = 0.17 \pm 0.27$, median ≤ 0.12 ng/g), concentrations in

nectar were significantly higher in Hertfordshire hives ($U_{44, 42}=1301$, $p\leq 0.001$, $\text{mean}_H=0.88\pm 0.81$, median=0.75 ng/g, $\text{mean}_{SU}=0.40\pm 0.80$ ng/g, median ≤ 0.10 ng/g). Crop mapping of the five 5 km² study areas in each region in 2014, showed that arable crops accounted for 55% of land cover in Hertfordshire (9% oilseed rape), and 32% in Sussex (5% oilseed rape, Figure S4).

Species-specific differences: A comparison of residue concentrations in pollen and nectar collected from adjacent honeybee (HB) and bumblebee (BB) nests located in rural habitats in Hertfordshire and Sussex revealed significantly higher concentrations of neonicotinoid exposure for honeybees compared to bumblebees (Table 1, 2, $U_{18, 18}= 112$, $p=0.04$; $\text{mean}_{HB}=0.17\pm 0.39$ ng/g, median <MDL, max=1.38 ng/g; $\text{mean}_{BB}\leq 0.12$ ng/g, median <MDL, max ≤ 0.12 ng/g).

DISCUSSION

In December 2013, an EU-wide moratorium on the use of certain neonicotinoids on bee-attractive flowering crops was implemented by the European Commission, which in early 2018 was subsequently expanded to include all field crops. To monitor bees' exposure to neonicotinoids during the initial transitional period from pre- to post-ban, between 2013 and 2015 we collected more than 400 pollen and nectar samples from bumblebee and honeybee colonies located in rural and peri-urban habitats in three regions across the UK, finding just over half of all samples to be contaminated with neonicotinoids. While combined total concentrations of neonicotinoids in pollen collected by rural bumblebees declined post-ban from an average of 5.1 ng/g in 2013, to 0.06 ng/g in 2015, suggesting a positive impact of the moratorium, neonicotinoid concentrations detected in samples collected from peri-urban bumblebee colonies remained largely unchanged between 2013 and 2015, indicating that the risk of exposure for peri-urban bees was not altered during the transitional period, and that more could be done to mitigate the risk for bees foraging in such habitats.

Across all samples, the highest neonicotinoid residue concentrations were detected in 2013, in pollen samples collected from rural bumblebee colonies in Sussex. Concentrations of up to 38.77 ng/g of thiamethoxam were detected, with the average total neonicotinoid concentrations of 5.1 ng/g

322 similar to that detected by previous studies conducted prior to the moratorium^{25,15,26}, and within the
323 range demonstrated to have negative impacts on bumblebee physiology^{27,28}, foraging efficiency²⁹ and
324 colony growth²⁸. Pre-ban (2013), the frequency of neonicotinoid contamination was extremely high
325 for pollen sampled from bumblebee colonies in both rural and peri-urban habitats in Sussex (74% and
326 84% of pollen samples respectively, mean=3.74 ng/g). As predicted, pollen samples collected from
327 nests near Stirling in 2013 were contaminated to a lesser degree (23-30% of pollen samples), and with
328 lower concentrations (mean=0.20 ng/g). This likely reflects the fact that across Scotland,
329 neonicotinoid use in 2013/2014 was approximately four times lower than in South East England (4,
330 186 kg, over 78, 345 ha vs. 16, 820 kg, over 197,507 ha²²), though differences in the growth season
331 and therefore timing of neonicotinoid application between regions may also have played a role.

332 Pollen and nectar samples collected from honeybee colonies in 2014, post-implementation of
333 the ban, but when any winter-sown oilseed rape may still have been seed-treated with neonicotinoids,
334 also had a high prevalence of neonicotinoid contamination (68% samples). Contamination was highest
335 in April when oilseed rape was flowering (93% samples), and declined throughout the season, a
336 phenomenon observed in several earlier studies^{14,15,23,30}, and hypothesised to arise from temperature
337 increases and photo-degradation of neonicotinoid residues in plant tissues as the season progresses³¹.
338 During this early part of the year, concentrations detected in honeybee-collected nectar averaged 1.2
339 ng/g, close to the average maximum concentration detected in seed-treated crop nectar, as reported by
340 Godfray *et al.*³² (1.9 ng/g, averaged from 20 published studies). Concentrations in pollen were
341 considerably lower (0.32 ng/g, average maximum concentration in seed-treated crop pollen=6.1
342 ng/g³²), likely reflecting honeybees' preference for collecting nectar from oilseed rape. For both
343 bumblebees and honeybees, early spring is a period when the colony might be expected to be
344 particularly vulnerable^{33,34}, and levels detected in pollen were within the range known to impair
345 honeybee foraging performance³⁵, immune function⁷ and alter gene expression pathways³⁶.
346 Furthermore, as observed in several previous studies^{15,17,18}, many of the samples we screened were
347 found to contain more than one neonicotinoid residue, which gives rise to the potential for additive or
348 synergistic effects. Tosi *et al.*¹⁷ found when screening honeybee pollen collected from multiple

apiaries across Italy for 66 different pesticides, that the frequency of detection actually peaked in summer months. Though here we did not screen for the presence of other chemical classes such as fungicides, there is evidence to suggest that exposure to certain fungicides can make bees more susceptible to the adverse effects of neonicotinoids³⁷.

Although the concentration of neonicotinoids in pollen and nectar sampled from rural bumblebee colonies declined between 2013 and 2015, bumblebees from both rural and peri-urban habitats were nevertheless still exposed to neonicotinoids following the implementation of the ban. Indeed 47% of nectar and 36% of pollen samples collected from rural colonies in 2015 contained neonicotinoid residues, a similar frequency as observed for peri-urban nests (33% nectar, 64% pollen), albeit at lower concentrations (mean concentration detected in pollen from rural nests = 0.06 ng/g vs. 1.29 ng/g detected in peri-urban pollen in 2015). This echoes the findings of Woodcock *et al.*³⁰ who screened honey samples submitted by beekeepers across the UK, and found that while samples harvested in 2014 were more likely to be contaminated (52% samples), 22.9% of samples harvested post-ban in 2015 also contained neonicotinoids. Similarly, a worldwide study of honey contamination spanning five years between 2012 and 2016, found 75% of 198 samples to contain neonicotinoids, with the highest prevalence in honey from North America, Asia and Europe³⁸.

Not only did exposure to neonicotinoids change for rural bees between 2013 and 2015, so did the chemical type. Across all samples, thiamethoxam was the most frequently detected, which is unsurprising given that, prior to the moratorium, it was the active ingredient in the mostly commonly used seed dressing on oilseed rape across Great Britain. Indeed, of fifteen farmers growing winter-sown oilseed rape within a 5 km radius of our experimental bee colonies that we interviewed in 2014, twelve had used seeds dressed with a thiamethoxam-based formulation (Cruiser®). Clothianidin, a metabolite of thiamethoxam and still in use as a seed-dressing on non-flowering cereal crops, was also frequently detected in honeybee nectar (69% samples), but only once in pollen, and was rarely detected in any samples collected from bumblebee colonies. Post-ban, acetamiprid and thiacloprid, the use of which is unaffected by the moratorium, were detected more often and at higher levels than thiamethoxam. For nectar samples collected from rural bumblebee colonies, thiacloprid

376 concentrations actually significantly increased between 2013 and 2015. Thiacloprid is an active
377 ingredient in many bug sprays sold in garden centres, and a recent study in which ornamental ‘bee-
378 friendly’ plants were screened for multiple pesticide and fungicide residues found more than 70% of
379 plants contained neonicotinoids, with thiacloprid present in almost half²⁴.

380 Imidacloprid was detected in a moderate proportion (10%) of samples collected from
381 bumblebee nests throughout the duration of the study. Considering that use of imidacloprid in arable
382 farming has dramatically declined in the UK (50% and 90% decline in weight of imidacloprid applied
383 to cereals and oilseeds respectively between 2012 and 2014, PUS Stats database, Table S6), having
384 been replaced by thiamethoxam and clothianidin, it is somewhat concerning that it was detected to
385 such an extent. Woodcock et al.³⁰ also noted that imidacloprid was present in honey samples
386 harvested in 2014 at a rate ‘disproportional to its use’ and Tosi et al.¹⁷ detected imidacloprid in 9.1%
387 of honeybee-collected pollen sampled from multiple apiaries across Italy in 2014 at mean
388 concentrations of 2 ng/g, raising concerns about the persistence of this chemical in agro-
389 environments. As previously observed when screening pollen from bumblebee colonies¹⁵ and wild
390 bumblebees collected in peri-urban areas²³, the highest concentrations of imidacloprid were detected
391 in peri-urban colonies, at levels up to 11.16 ng/g in 2015 (mean=1.13 ng/g). Again, this may originate
392 from use by the horticulture industry, since screening of ornamental plants detected imidacloprid in
393 38% of samples²⁴. An alternative, yet untested source, is the use of imidacloprid for flea control in
394 domestic pets and as ant poison.

395 Honeybees in Hertfordshire were exposed to significantly higher neonicotinoid concentrations
396 in nectar compared to Sussex honeybees, which is most likely explained by the fact that, in 2014,
397 there was almost double the percentage cover of treated oilseed crops (9% land cover in Hertfordshire
398 vs. 5% in Sussex), and generally a higher percentage of arable land cover (55%) compared to Sussex
399 (32%).

400 Overall, honeybee samples had higher concentrations of neonicotinoids compared to
401 bumblebees. This contrasts with findings from an earlier study conducted in 2013 where the reverse
402 was found to be true¹⁵. However in the previous study, colonies of each species were not placed in

identical locations, therefore in addition to differences in foraging range and flower preferences^{39,40}, colonies may simply have been in proximity to a different range of plant species. Clearly more paired sampling of both species is required to establish whether there are consistent differences in exposure.

On the basis of evidence published post-2013, the European Food Standards Agency recently concluded that neonicotinoids do indeed pose a risk to bees⁴¹, and in 2017 the EU commission proposed extending the moratorium to include all field crops (barring permanent greenhouse crops), which was passed by the European Union in early 2018¹⁰⁻¹². Here we have shown for the first time how exposure to neonicotinoids has changed for bees foraging in rural and peri-urban areas across the UK, since the initial implementation of the moratorium on their usage in December 2013. The exposure of rural bumblebees appears to have declined post-ban, suggesting that continued limitation of their use on flowering crops could have a positive impact on the risk for bees and other pollinators in rural areas. However, exposure for peri-urban bees remains largely unaffected, presumably as a result of contaminated ornamental plants sold in garden centres and ongoing domestic usage of neonicotinoid-based bug sprays. This is concerning given the growing interest in encouraging pollinators in urban areas; more research is needed to understand the sources of exposure and find ways to reduce it.

FIGURES




Figure 1 Map of the UK showing the regions in which honeybee (Hertfordshire and Sussex, 2014) and bumblebee (Stirling, 2013; Hertfordshire, 2014; Sussex 2013-2015) colonies were placed in rural (honeybees and bumblebees) and peri-urban (bumblebees only) habitats (see Fig. S1-3 for exact locations).

Moratorium Status	Year	Region	Bee Species	Habitat	N Colonies	Sampling Dates
Pre-ban	2013	Stirling	Bumblebee	Rural	10	12 th June; 11 th July; 18 th July
				Peri-urban	20	6 th June; 4 th July; 17 th July
		Sussex	Bumblebee	Rural	32	30 th May; 9 th June; 23 rd June
				Peri-urban	12	30 th May; 9 th June; 23 rd June
During ban (Winter-sown crops still seed-treated)	2014	Sussex	Bumblebee	Rural	47	28 th May; 25 th June; 9 th July
				Peri-urban	15	28 th May; 25 th June; 9 th July
			Honeybee	Rural	15	16 th April; 28 th May; 25 th June
		Herts	Honeybee	Rural	15	16 th April; 28 th May; 25 th June
			Bumblebee	Rural	30	28 th May; 25 th June; 9 th July
During ban	2015	Sussex	Bumblebee	Rural	45	15 th June; 13 th July; 27 th July
				Peri-urban	15	15 th June; 13 th July; 27 th July

Table 1 Number of honeybee and bumblebee colonies placed in each habitat type (Peri-urban vs. Rural), in each region (Sussex, Stirling, Hertfordshire (Herts)) across the three years of the study (2013-2015). The specific dates colonies were sampled for pollen and nectar are listed.

Year	Region	Location	N Colonies	N Samples	ng/g	NECTAR					TOTAL	% Multi-residue	POLLEN					TOTAL	% Multi-residue	
						Method Quantification Limit (ng/g)							MQL							
						0.3	0.3	0.4	0.08	0.08			0.12	0.12	0.16	0.04	0.04			
Method Detection Limit (ng/g)						0.1	0.1	0.14	0.03	0.03	MDL									
						TMX	CLO	IMC	ACT	THC			N	TMX	CLO	IMC	ACT	THC		
468	2013	STIRLING	Peri-Urban	20	8	Frequency %	12.5%				12.5%	0%	13	7.7%	7.7%		15.4%	23.1%	8.3%	
469						Mean ±SD	≤0.10				≤0.10			≤0.12	≤0.12		0.06±0.22	0.08±0.21		
470						Median	≤0.10				≤0.10			≤0.12	≤0.12		≤0.04	≤0.12		
471						Max	≤0.10				≤0.10			≤0.12	≤0.12		0.76	0.76		
472						Frequency %	12.5%				12.5%	0%		7.7%	7.7%		30.8%	30.8%	15.3%	
473		SUSSEX	Rural	10	7	Mean ±SD	0.26±0.68				0.26±0.68		13	≤0.12	0.16±0.58		0.15±0.36	0.32±0.65		
474						Median	≤0.12				≤0.10			≤0.12	≤0.10		≤0.03	≤0.12		
475						Max	1.81				1.81			≤0.12	2.08		1.15	2.08		
476						Frequency %	7.7%		7.7%		15.4%	0%		79%	5.26%	26.3%	nt	15.8%	84.2%	36.8%
477						Mean ±SD	≤0.10		≤0.14		≤0.10			0.58±1.64	≤0.10	≤0.16		1.47±4.41	2.11±4.56	
478	2014	Peri-Urban	15	5	Median	≤0.10		≤0.14		≤0.10		7	≤0.12	≤0.10	≤0.16		≤0.04	≤0.16		
479					Max	≤0.10		≤0.14		≤0.14			7.1	≤0.10	≤0.16		14.68	14.8		
480					Frequency %	14.3%				14.3%	0%		60.9%	4.35%	39.1%	nt	17.4%	74%	34.8%	
481					Mean ±SD	0.2±0.51				0.20±0.51			4.96±11.29	≤0.12	≤0.16		0.08±0.31	5.10±11.41		
482					Median	≤0.10				≤0.10			≤0.12	≤0.12	≤0.16		≤0.04	≤0.12		
483	2015	Peri-Urban	15	12	Max	1.49				1.49		11	38.77	≤0.12	≤0.16		1.5	38.93		
484					Frequency %	80.0%	40.0%			80.0%	40%		57.1%		14.3%	14.3%	42.9%	85.7%	28.6%	
485					Mean ±SD	0.76±1.52	≤0.10			0.80±1.56			≤0.12		0.31±0.82	≤0.04	1.34±3.52	1.73±3.43		
486					Median	0.10	≤0.10			0.1			≤0.12		≤0.16	≤0.04	≤0.04	≤0.16		
487					Max	3.48	≤0.10			3.58			≤0.12		2.18	≤0.04	9.32	9.32		
488	2016	Rural	47	13	Frequency %	8.3%				8.3%	0%	13	7.7%					7.7%	0.0%	
489					Mean ±SD	≤0.10				≤0.10			≤0.12		≤0.16	≤0.04	≤0.04	≤0.16		
490					Median	≤0.10				≤0.10			≤0.12		≤0.16	≤0.04	≤0.04	≤0.16		
491					Max	≤0.10				≤0.10			≤0.12		≤0.16	≤0.04	≤0.04	≤0.16		
492					Frequency %	20%				20%	0%					9.1%	36.4%	36.4%	9.1%	
493	2017	Rural	30	10	Mean ±SD	≤0.10				≤0.10		11	≤0.04	≤0.04	≤0.04	≤0.04	≤0.04			
494					Median	≤0.10				≤0.10			≤0.04	≤0.04	≤0.04	≤0.04	≤0.04			
495					Max	≤0.10				≤0.10			≤0.04	≤0.04	≤0.04	≤0.04	≤0.04			
496					Frequency %			16.7%		25%	33.3%		8.3%	9.1%		27.3%	36.35	18.2%	63.6%	18.2%
497					Mean ±SD			≤0.14		0.05±0.13	0.08±0.17			≤0.12		1.13±3.34	0.14±0.42	≤0.04	1.29±3.30	
498	2018	Peri-Urban	15	12	Median			≤0.14		≤0.03	≤0.10		11	≤0.12		≤0.16	≤0.04	≤0.04	≤0.04	
499					Max			≤0.14		0.44	0.44			≤0.12		11.16	1.40	≤0.04	11.16	
500					Frequency %	5.3%		5.3%		36.8%	47.4%	0%		13.6%		9.1%	9.1%	13.6%	36.4%	9.1%
501					Mean ±SD	≤0.10		≤0.14		0.09±0.15	0.10±0.15			≤0.12		≤0.16	≤0.04	≤0.04	0.06±0.14	
502					Median	≤0.10		≤0.14		≤0.03	≤0.10			≤0.12		≤0.16	≤0.04	≤0.04	≤0.12	
503	2019	Rural	45	19	Max	≤0.10		≤0.14		0.42	0.42		≤0.12		0.60	≤0.04	≤0.04	0.60		

Table 2 Frequency of detection (% samples), mean (± standard deviation (SD)), median and maximum concentrations of five neonicotinoids (TMX=thiamethoxam, CLO= clothianidin, IMC= imidacloprid, ACT=acetamiprid, THC=thiacloprid) and the combined total concentration of neonicotinoids detected in pollen and nectar sampled from bumblebee colonies located in rural and peri-urban habitats in three different regions; Stirling, Hertfordshire (Herts) and Sussex. Samples were collected across three years (2013-2015). Multi-residue samples are those where more than one type of neonicotinoid was detected. *MQL*= Method quantification limit, *MDL*=Method detection limit, *nt*= not tested, ≤ less than or equal to.

489				NECTAR						POLLEN								
	Method Quantification Limit (ng/g)			0.3	0.3	0.4	0.08	0.08			MQL	0.36	0.36	0.48	0.12	0.12		
490	Method Detection Limit (ng/g)			0.1	0.1	0.14	0.03	0.03			MDL	0.12	0.12	0.16	0.04	0.04		
	Month	Region	N	TMX	CLO	IMC	ACT	THC	TOTAL	% Multi-residue	N	TMX	CLO	IMC	ACT	THC	TOTAL	% Multi-residue
491	APRIL	HERTS	15	Frequency of detection %	100%	73.3%	6.7%		100%	80.0%		80%		6.6%		13.3%	80%	20.0%
492				Mean ± SD (ng/g)	0.83 ± 0.48	0.63 ± 0.51	≤0.14		1.46±0.66		15	0.26±0.28		≤0.16		0.14±0.42	0.41±0.47	
				Median (ng/g)	0.77	0.66	≤0.14		1.17			0.12		≤0.16		≤0.04	0.12	
493				Max (ng/g)	1.83	1.38	≤0.14		1.83			0.94		≤0.16		1.62	1.62	
494		SUSSEX	15	Frequency of detection %	93%	47%	7%	7%	93.3%	60.0%		100%					100%	0%
				Mean ± SD (ng/g)	0.56± 0.14	0.37±0.18	≤0.14	≤0.03	0.95 ±1.13		15	0.23±0.19					0.23±0.19	
				Median (ng/g)	0	≤0.1	≤0.14	≤0.03	0.58			0.12					0.12	
495				Max (ng/g)	1.76	2.47	≤0.03	≤0.03	2.47			0.6					0.60	
496	MAY	HERTS	15	Frequency of detection %	86.6%	73.3%			93.3%	66.7%		80%					80%	0%
				Mean ± SD (ng/g)	0.60±0.16	0.38±0.11			1.04±0.74		15	0.19±0.24					0.19±0.24	
497				Median (ng/g)	0.45	0.10			1.08			0.12					0.12	
				Max (ng/g)	2.29	1.26			2.29			0.92					0.92	
498		SUSSEX	12	Frequency of detection %	66.7%	16.7%		16.70%	66.7%	25.0%		53.3%	6.7%	6.7%		20%	66.7%	20%
				Mean ± SD (ng/g)	0.12±0.05	≤0.10		≤0.03	0.19±0.34		15	≤0.12	≤0.12	≤0.16		0.16±0.4	0.24±0.4	
499				Median (ng/g)	0.10	≤0.10		≤0.03	0.10			≤0.12	≤0.12	≤0.16		≤0.04	0.1	
				Max (ng/g)	0.53	0.68		≤0.03	0.68			≤0.12	≤0.12	≤0.16		1.19	1.2	
500	JUNE	HERTS	14	Frequency of detection %	50%	21.4%	7.1%		66.3%	21.4%		26.7%		6.7%			26.7%	8.9%
501				Mean ± SD (ng/g)	≤0.10	≤0.10	≤0.14		0.08±0.08		15	≤0.12		≤0.16			0.09±0.26	
				Median (ng/g)	≤0.10	≤0.10	≤0.14		0.10			≤0.12		≤0.16			≤0.12	
502				Max (ng/g)	≤0.10	≤0.10	≤0.14		≤0.14			≤0.12		0.88			0.88	
503		SUSSEX	15	Frequency of detection %	13.3%				13.3%	0%		26.7%		6.7%		6.7%	33.3%	6.7%
				Mean ± SD (ng/g)	≤0.10				≤0.10		15	≤0.12		≤0.16		≤0.04	0.05±0.07	
				Median (ng/g)	≤0.10				≤0.10			≤0.12		≤0.16		≤0.04	≤0.12	
504				Max (ng/g)	≤0.10				≤0.10			≤0.12		≤0.16		≤0.04	≤0.16	

505

506 Table 3 Frequency of detection (% samples), mean (\pm standard deviation (SD)), median and maximum concentrations of five neonicotinoids
507 (TMX=thiamethoxam, CLO= clothianidin, IMC= imidacloprid, ACT=acetamiprid, THC=thiacloprid) and the combined total concentration of neonicotinoids
508 detected in honeybee nectar and pollen sampled from colonies located in in Sussex (N=15) and Hertfordshire (Herts, N=15) between April and June. Multi-
509 residue samples are those where more than one type of neonicotinoid was detected. *MQL*= Method quantification limit, *MDL*=Method detection limit, *nt*= not
510 tested, \leq less than or equal to.




Figure 2 Total neonicotinoid concentrations (Thiamethoxam, clothianidin, imidacloprid, acetamiprid and thiacloprid combined) detected in A) Pollen and B) Nectar samples collected from bumblebee colonies in Rural (White, N Pollen samples=45; Nectar=33) and Peri-urban (Grey, N Pollen samples=30; Nectar=25) habitats across the region of Sussex in the years 2013 and 2015. Concentrations are plotted on a square-root scale. Black horizontal bars show median values. Box limits denote the first and third quantiles, and boxplot whiskers extend to 1.5 times the interquartile range. Outliers are represented by solid black circles.

541 **ASSOCIATED CONTENT**

542 **Supporting Information**

543 The following file is available free of charge.

544 Additional figures and tables as mentioned in the text (PDF)

545 **AUTHOR INFORMATION**

546 **Corresponding Author**

547 *Email: e.nicholls@sussex.ac.uk. Phone: +44(0)1273 873310

548 **Present Addresses**

549 [‡]Doñana Biological Station (EBD-CSIC), Integrative Ecology Department, Seville, ES-41092, Spain.

550

551 [†]Karlsruhe Institute of Technology (KIT), Institute of Meteorology and Climate Research–
552 Atmospheric Environmental Research (IMK-IFU), 82467 Garmisch-Partenkirchen, Germany

553

554 [§] School of Geosciences, Grant Institute, King's Buildings, Edinburgh, EH9 3JW, UK

555 [‡] French School of Public Health (EHESP), Research Institute for Environmental and Occupational
556 Health (Irset - Inserm UMR 1085), 35043 Rennes, France

557 **Notes**

558 The authors declare no competing financial interest.

559 **ACKNOWLEDGEMENTS**

560 We are grateful to BBSRC (BB/K014498/1; BB/J014915/1) and Defra (Research Project PS2372) for
561 funding this work and for the farmers and land-owners who allowed us to work on their property and
562 shared their pesticide usage data. We thank Bill Hughes, Mark Crumpton, Steve Kennedy and
563 Luciano Scandian for beekeeping equipment, assistance and advice, and Pete Kennedy and Martyn
564 Stenning for useful discussions regarding experimental design and logistics. Thanks also to Julia

Horwood and Daniel Ingram for technical advice and assistance and to Grace Twiston-Davies for digitising crop maps.

REFERENCES

- (1) Simon-Delso, N.; Amaral-Rogers, V.; Belzunces, L. P.; Bonmatin, J. M.; Chagnon, M.; Downs, C.; Furlan, L.; Gibbons, D. W.; Giorio, C.; Girolami, V.; Goulson, D. Kreutzweiser, D. P.; Krupke, C.; H. Liess, M.; Long, E.; McField, M.; Mineau, P.; Mitchell, E.; A. D. Morrissey, C. A.; Noome, D. A.; Pisa, L.; Settele, J.; Stark, J. D.; Tapparo, A.; Van Dyck, H.; Van Praagh, J.; Van der Sluijs, J. P.; Whitehorn, P. R.; Wiemers, M. Systemic Insecticides (Neonicotinoids and Fipronil): Trends, Uses, Mode of Action and Metabolites. *Environ. Sci. Pollut. Res.* **2014**.
- (2) Ollerton, J.; Winfree, R.; Tarrant, S. How Many Flowering Plants Are Pollinated by Animals? *Oikos* **2011**, *120* (3), 321–326.
- (3) Alkassab, A. T.; Kirchner, W. H. Sublethal Exposure to Neonicotinoids and Related Side Effects on Insect Pollinators: Honeybees, Bumblebees, and Solitary Bees. *J. Plant Dis. Prot.* **2017**, *124* (1), 1–30.
- (4) Whitehorn, P. R.; O'Connor, S.; Wackers, F. L.; Goulson, D. Neonicotinoid Pesticide Reduces Bumble Bee Colony Growth and Queen Production. *Science* **2012**, *336* (6079), 351–352.
- (5) Gill, R. J.; Ramos-Rodriguez, O.; Raine, N. E. Combined Pesticide Exposure Severely Affects Individual- and Colony-Level Traits in Bees. *Nature* **2012**, *491* (7422), 105–108.
- (6) Henry, M.; Béguin, M.; Requier, F.; Rollin, O.; Odoux, J.-F.; Aupinel, P.; Aptel, J.; Tchamitchian, S.; Decourtye, A. A Common Pesticide Decreases Foraging Success and Survival in Honey Bees. *Science* **2012**, *336* (6079), 348–350.
- (7) Di Prisco, G.; Cavaliere, V.; Annoscia, D.; Varricchio, P.; Caprio, E.; Nazzi, F.;

- 590 Gargiulo, G.; Pennacchio, F. Neonicotinoid Clothianidin Adversely Affects Insect
591 Immunity and Promotes Replication of a Viral Pathogen in Honey Bees. *Proc. Natl.*
592 *Acad. Sci. U. S. A.* **2013**, *110* (46), 18466–18471.
- 593 (8) Laycock, I.; Cresswell, J. E. Repression and Recuperation of Brood Production in
594 *Bombus Terrestris* Bumble Bees Exposed to a Pulse of the Neonicotinoid Pesticide
595 Imidacloprid. *PLoS One* **2013**, *8* (11), e79872–e79872.
- 596 (9) European Commission. Commission Implementing Regulation (EU) No 485/2013 of
597 24 May 2013 Amending Implementing Regulation (EU) No 540/2011, as Regards the
598 Condition of Approval of the Active Substances Clothianidin, Thiamethoxam and
599 Imidacloprid, and Prohibiting the Use and Sa. *Off. J. Eur. Union* **2013**, *L139/12*.
- 600 (10) European Commission. Commission Implementing Regulation (EU) 2018/783 of 29
601 May 2018 Amending Implementing Regulation (EU) No 540/2011 as Regards the
602 Conditions of Approval of the Active Substance Imidacloprid. *Off. J. Eur. Union* **2018**,
603 *L 132/31*.
- 604 (11) European Commission. Commission Implementing Regulation (EU) 2018/784 of 29
605 May 2018 Amending Implementing Regulation (EU) No 540/2011 as Regards the
606 Conditions of Approval of the Active Substance Clothianidin. *Off. J. Eur. Union* **2018**,
607 *L132/35*.
- 608 (12) European Commission. Commission Implementing Regulation (EU) 2018/785 of 29
609 May 2018 Amending Implementing Regulation (EU) No 540/2011 as Regards the
610 Conditions of Approval of the Active Substance Thiamethoxam. *Off. J. Eur. Union*
611 **2018**, *L 132/40*.
- 612 (13) Carreck, N. L.; Ratnieks, F. L. W. The Dose Makes the Poison: Have “field Realistic”
613 Rates of Exposure of Bees to Neonicotinoid Insecticides Been Overestimated in
614 Laboratory Studies? *J. Apic. Res.* **2014**, *53* (5), 607–614.

- 615 (14) Botías, C.; David, A.; Horwood, J.; Abdul-Sada, A.; Nicholls, E.; Hill, E.; Goulson, D.
616 Neonicotinoid Residues in Wildflowers, a Potential Route of Chronic Exposure for
617 Bees. *Environ. Sci. Technol.* **2015**, *49* (21), 12731–12740.
- 618 (15) David, A.; Botías, C.; Abdul-Sada, A.; Nicholls, E.; Rotheray, E. L.; Hill, E. M.;
619 Goulson, D. Widespread Contamination of Wildflower and Bee-Collected Pollen with
620 Complex Mixtures of Neonicotinoids and Fungicides Commonly Applied to Crops.
621 *Environ. Int.* **2016**, *88*, 169–178.
- 622 (16) Rundlöf, M.; Andersson, G. K. S.; Bommarco, R.; Fries, I.; Hederström, V.;
623 Herbertsson, L.; Jonsson, O.; Klatt, B. K.; Pedersen, T. R.; Yourstone, J.; Smith, Henrik
624 G. Seed Coating with a Neonicotinoid Insecticide Negatively Affects Wild Bees.
625 *Nature* **2015**, *521*, 77–80.
- 626 (17) Tosi, S.; Costa, C.; Vesco, U.; Quaglia, G.; Guido, G. A 3-Year Survey of Italian
627 Honey Bee-Collected Pollen Reveals Widespread Contamination by Agricultural
628 Pesticides. *Sci. Total Environ.* **2018**, *615*, 208–218.
- 629 (18) Colwell, M. J.; Williams, G. R.; Evans, R. C.; Shutler, D. Honey Bee-Collected Pollen
630 in Agro-Ecosystems Reveals Diet Diversity, Diet Quality, and Pesticide Exposure.
631 *Ecol. Evol.* **2017**, *7* (18), 7243–7253.
- 632 (19) Balfour, N. J.; Al Toufalia, H.; Scandian, L.; Blanchard, H. E.; Jesse, M. P.; Carreck,
633 N. L.; Ratnieks, F. L. W. Landscape Scale Study of the Net Effect of Proximity to a
634 Neonicotinoid-Treated Crop on Bee Colony Health. *Environ. Sci. Technol.* **2017**, *51*
635 (18), 10825–10833.
- 636 (20) Rolke, D.; Persigehl, M.; Peters, B.; Sterk, G.; Blenau, W. Large-Scale Monitoring of
637 Effects of Clothianidin-Dressed Oilseed Rape Seeds on Pollinating Insects in Northern
638 Germany: Residues of Clothianidin in Pollen, Nectar and Honey. *Ecotoxicology* **2016**,
639 *25* (9), 1691–1701.

- 640 (21) Woodcock, B. A.; Bullock, J. M.; Shore, R. F.; Heard, M. S.; Pereira, M. G.; Redhead,
641 J.; Ridding, L.; Dean, H.; Sleep, D.; Henrys, P.; Peyton, J.
642 Hulmes, S.; Hulmes, L.; Sárospataki, M.; Saure, C.; Edwards, M.; Genersch, E.;
643 Knäbe, S.; Pywell, R. F. Country-Specific Effects of Neonicotinoid Pesticides on
644 Honey Bees and Wild Bees. *Science* (80-.). **2017**, *356* (6345), 1393–1395.
- 645 (22) FERA PUS STATS for insecticides.
- 646 (23) Botías, C.; David, A.; Hill, E. M.; Goulson, D. Quantifying Exposure of Wild
647 Bumblebees to Mixtures of Agrochemicals in Agricultural and Urban Landscapes.
648 *Environ. Pollut.* **2017**, *222*, 73–82.
- 649 (24) Lentola, A.; David, A.; Abdul-Sada, A.; Tapparo, A.; Goulson, D.; Hill, E. M.
650 Ornamental Plants on Sale to the Public Are a Significant Source of Pesticide Residues
651 with Implications for the Health of Pollinating Insects. *Environ. Pollut.* **2017**, *228*,
652 297–304.
- 653 (25) Scheper, J.; Holzschuh, A.; Kuussaari, M.; Potts, S. G.; Rundlöf, M.; Smith, H. G.;
654 Kleijn, D. Environmental Factors Driving the Effectiveness of European Agri-
655 Environmental Measures in Mitigating Pollinator Loss--a Meta-Analysis. *Ecol. Lett.*
656 **2013**, *16* (7), 912–920.
- 657 (26) Botías, C.; David, A.; Hill, E. M.; Goulson, D. Contamination of Wild Plants near
658 Neonicotinoid Seed-Treated Crops, and Implications for Non-Target Insects. *Sci. Total*
659 *Environ.* **2016**, *566-567*, 269–278.
- 660 (27) Moffat, C.; Pacheco, J. G.; Sharp, S.; Samson, A. J.; Bolland, K. A.; Huang, J.;
661 Buckland, S. T.; Connolly, C. N. Chronic Exposure to Neonicotinoids Increases
662 Neuronal Vulnerability to Mitochondrial Dysfunction in the Bumblebee (*Bombus*
663 *terrestris*). *FASEB J.* **2015**, *29* (5), 2112–2119.
- 664 (28) Moffat, C.; Buckland, S. T.; Samson, A. J.; McArthur, R.; Chamosa Pino, V.; Bolland,

- 665 K. A.; Huang, J. T. J.; Connolly, C. N. Neonicotinoids Target Distinct Nicotinic
666 Acetylcholine Receptors and Neurons, Leading to Differential Risks to Bumblebees.
667 *Sci. Rep.* **2016**, *6*.
- 668 (29) Feltham, H.; Park, K.; Goulson, D. Field Realistic Doses of Pesticide Imidacloprid
669 Reduce Bumblebee Pollen Foraging Efficiency. *Ecotoxicology* **2014**, *23* (3), 317–323.
- 670 (30) Woodcock, B. A.; Ridding, L.; Freeman, S. N.; Gloria Pereira, M.; Sleep, D.; Redhead,
671 J.; Aston, D.; Carreck, N. L.; Shore, R. F.; Bullock, J. M.; Heard, Matthew S.;
672 Pywell, Richard F. Neonicotinoid Residues in UK Honey despite European Union
673 Moratorium. *PLoS One* **2018**, *13* (1).
- 674 (31) Bonmatin, J.-M.; Giorio, C.; Girolami, V.; Goulson, D.; Kreutzweiser, D. P.; Krupke,
675 C.; Liess, M.; Long, E.; Marzaro, M.; Mitchell, E. a. D.; Noome, D. A.; Simon-Delso,
676 N.; Tapparo, A. Environmental Fate and Exposure; Neonicotinoids and Fipronil.
677 *Environ. Sci. Pollut. Res.* **2014**.
- 678 (32) Godfray, H. C. J.; Blacquière, T.; Field, L. M.; Hails, R. S.; Petrokofsky, G.; Potts, S.
679 G.; Raine, N. E.; Vanbergen, A. J.; McLean, A. R. A Restatement of the Natural
680 Science Evidence Base Concerning Neonicotinoid Insecticides and Insect Pollinators.
681 *Proc. Biol. Sci.* **2014**, *281* (1786).
- 682 (33) Seeley, T. D.; Visscher, P. K. Survival of Honeybees in Cold Climates: The Critical
683 Timing of Colony Growth and Reproduction. *Ecol. Entomol.* **1985**, *10* (1), 81–88.
- 684 (34) Wu-Smart, J.; Spivak, M. Sub-Lethal Effects of Dietary Neonicotinoid Insecticide
685 Exposure on Honey Bee Queen Fecundity and Colony Development. *Sci. Rep.* **2016**, *6*.
- 686 (35) Cresswell, J. E. A Meta-Analysis of Experiments Testing the Effects of a
687 Neonicotinoid Insecticide (Imidacloprid) on Honey Bees. *Ecotoxicology* **2011**, *20* (1),
688 149–157.
- 689 (36) Christen, V.; Mittner, F.; Fent, K. Molecular Effects of Neonicotinoids in Honey Bees

- 690 (Apis Mellifera). *Environ. Sci. Technol.* **2016**, acs.est.6b00678.
- 691 (37) Iwasa, T.; Motoyama, N.; Ambrose, J. T.; Roe, R. M. M. Mechanism for the
692 Differential Toxicity of Neonicotinoid Insecticides in the Honey Bee, *Apis Mellifera*.
693 *Crop Prot.* **2004**, 23 (5), 371–378.
- 694 (38) Mitchell, E. A. D.; Mulhauser, B.; Mulot, M.; Mutabazi, A.; Glauser, G.; Aebi, A. A
695 Worldwide Survey of Neonicotinoids in Honey. *Science (80)*. **2017**, 358 (6359), 109–
696 111.
- 697 (39) Leonhardt, S. D.; Blüthgen, N. The Same, but Different: Pollen Foraging in Honeybee
698 and Bumblebee Colonies. *Apidologie* **2011**, 43 (4), 1–16.
- 699 (40) Wood, T. J.; Holland, J. M.; Goulson, D. Pollinator-Friendly Management Does Not
700 Increase the Diversity of Farmland Bees and Wasps. *Biol. Conserv.* **2015**, 187, 120–
701 126.
- 702 (41) EFSA. Evaluation of the Data on Clothianidin, Imidacloprid and Thiamethoxam for
703 the 402 Updated Risk Assessment to Bees for Seed Treatments and Granules in the
704 EU. *EFSA Support. Publ.* **2018**, 15, 1–31.

705

706

707

708

709

710